

STRUCTURE OF GENERAL IDEAL SEMIGROUPS OF MONOIDS AND DOMAINS

ANDREAS REINHART

ABSTRACT. Let H be a monoid (resp. an integral domain) and r an ideal system on H . In this paper we investigate the r -ideal semigroup of H . One goal is to specify monoids such that their r -ideal semigroup possesses semigroup-theoretical properties, like almost completeness, π -regularity and completeness. Moreover if H is an integral domain and $*$ a star operation on H , then we provide conditions on H such that the idempotents of the $*$ -ideal semigroup are trivial or such that H is $\pi*$ -stable.

0. INTRODUCTION

In 1961 E. C. Dade, O. Taussky and H. Zassenhaus [9] investigated the semigroup structure of the ideal class semigroup of one-dimensional noetherian domains with a focus on non-principal orders in algebraic number fields. In the sequel, this paper seems to have fallen into oblivion. It was reconsidered and generalized by F. Halter-Koch [17], who put the results into the context of the structure theory of semigroups as presented in [14]. In recent times, starting with a paper by P. Zanardo and U. Zannier [23], the structure of the ideal class semigroup of an integral domain attracted the interest of several authors. In particular, the Clifford and the Boolean property of ideal class semigroups was investigated. First, this was done for valuation domains in [7]. In the sequel, S. Bazzoni provided a general theory, focused on Prüfer domains [2], [3], [4], [5]. Among others, she proved that a Prüfer domain has a Clifford semigroup if and only if it has finite character, and she highlighted the connection with stable domains. Following S. Bazzoni, S. Kabbaj and A. Mimouni, in [20], [21] and [22], investigated the Clifford and Boolean property of the t -ideal class semigroup. Among others, they characterized Prüfer v -multiplication domains with Clifford t -class semigroup and determined the structure of their constituent groups.

A more general approach, based on the theory of ideal systems on commutative cancellative monoids, was recently presented by F. Halter-Koch [18]. He observed that the semigroup structure of the ideal class semigroup is essentially determined by the semigroup structure of the ideal semigroup itself. Among others, he proved that the semigroup of fractional r -ideals of an r -Prüfer monoid is a Clifford semigroup if and only if it has the local invertibility property (a generalization of Bazzoni's conjecture for Prüfer domains), and he characterized v -domains with a Clifford v -ideal semigroup.

A valuable overview of Clifford and Boolean properties of (t -)ideal class semigroups of integral domains may be found in the survey article by S. Bazzoni and S. Kabbaj [6]. A collection of important results concerning the ideal class semigroups of Prüfer domains can be found in [10].

In this paper, we reconsider the more general semigroup theoretical properties of ideal semigroups as this was done in [9] and [17], but now we depart from the special cases of one-dimensional or Prüfer domains. Following [18], we present our theory as far as possible in the language of ideal systems on commutative monoids [19]. We provide conditions on a monoid or domain which entail nice semigroup-theoretic properties (as almost completeness, π -regularity and completeness) of the ideal class semigroup in question.

2010 *Mathematics Subject Classification.* 13A15, 20M12, 13F05.

Key words and phrases. semigroup, star operation, ideal system, regular, stable.

Our main subject of interest is the v -ideal (class) semigroup of a Mori domain. The main results are in Section 4, and there the case of $* = v$ is of particular interest. The strength of our results in the ring-theoretical case is illustrated by a series of examples and counter-examples in Section 5.

In Section 1 we recall the relevant definitions and facts from the structure theory of commutative semi-groups, essentially following [14]. In Section 2 we introduce ideal systems and r -ideal semigroups, summarize some of their elementary properties and provide first results concerning stability and completeness. In Section 3 we discuss preparatory ring-theoretical results concerning Mori domains and a general prime avoidance lemma (Lemma 3.2) which seems to be new. As already mentioned, Section 4 contains the main results, in particular, criteria for the idempotents of the ideal semigroup to be trivial and criteria for πr -stability.

1. COMMUTATIVE SEMIGROUPS

Throughout this section, let S be a multiplicative commutative semigroup.

Our main reference for the theory of commutative semigroups is [14]. We denote by $\mathbf{E}(S)$ the set of all idempotents of S , endowed with the Rees order \leq , defined by $e \leq f$ if $ef = e$. Note that $ef \leq e$, for all $e, f \in \mathbf{E}(S)$. For $a \in S$, let $\mathbf{E}(a) = \{e \in \mathbf{E}(S) \mid ea = a\}$ be the set of all idempotents belonging to a and set $\mathbf{E}^\infty(a) = \bigcup_{k \in \mathbb{N}} \mathbf{E}(a^k)$. Then $\mathbf{E}(a) \subseteq \mathbf{E}^\infty(a) \subseteq \mathbf{E}(S) \subseteq S$ are subsemigroups. If $I \subseteq \mathbf{E}(S)$ is a subsemigroup, then every minimal element of I (with respect to the Rees-order) is a least element. In particular, if I is finite, then I has a minimum.

An element $a \in S$ is called

- *regular* if there exist elements $e \in \mathbf{E}(a)$ and $b \in S$ such that $ab = e$ (equivalently: There exists some $b \in S$ such that $a^2b = a$).
- π -*regular* if a^n is regular for some $n \in \mathbb{N}$.

The semigroup S is called

- *regular* or a *Clifford semigroup* if every $a \in S$ is regular.
- π -*regular* if every $a \in S$ is π -regular.
- *almost complete* if for every $a \in S$, the set $\mathbf{E}(a)$ possesses a minimum (in the Rees-order).
- *complete* if it is π -regular and almost complete.

For $a \in \mathbf{E}(S)$, let $P_a^* = aS \setminus \bigcup_{b \in \mathbf{E}(S), b <_S a} bS$ be the partial Ponizovski factor of a . Note that the partial Ponizovski factors are essential invariants for the structure of S (see [14] and [17, Theorems 2.4 and 2.5]).

Lemma 1.1. *Let $a \in S$.*

1. *If $n \in \mathbb{N}$ such that a^n is regular, then a^k is regular for all $k \in \mathbb{N}_{\geq n}$.*
2. *If there exist $k, n \in \mathbb{N}$ such that $k \neq n$ and $a^k = a^n$, then a is π -regular.*
3. *If a is π -regular, then $\mathbf{E}^\infty(a)$ has a smallest element in the Rees order.*

Proof. **1.** Let $n \in \mathbb{N}$ be such that a^n is regular and let $k \in \mathbb{N}_{\geq n}$. There exists some $l \in \mathbb{N}$ such that $nl > k$. Since a^n is regular, there exist $e \in \mathbf{E}(a^n)$ and $b \in S$ such that $a^n b = e$. Let $b' = a^{nl-k} b^l \in S$. Then $a^k b' = a^{nl} b^l = e^l = e$. Since $e \in \mathbf{E}(a^k)$, it follows that a^k is regular.

2. Let $k, n \in \mathbb{N}$ be such that $k < n$ and $a^k = a^n$. It is straightforward to show that $a^k = a^{k+i(n-k)}$ for all $i \in \mathbb{N}$. There exists some $j \in \mathbb{N}$ such that $j(n-k) > k$, hence $a^{j(n-k)} = a^k a^{j(n-k)-k} = a^{k+j(n-k)} a^{j(n-k)-k} = a^{2j(n-k)}$. It follows that $a^{j(n-k)}$ is regular, and thus a is π -regular.

3. Let a be π -regular. Then there exist $n \in \mathbb{N}$, $e \in \mathbf{E}(a^n)$ and $b \in S$ such that $a^n b = e$. This implies that $e \in \mathbf{E}^\infty(a)$. We assert that $e = \min \mathbf{E}^\infty(a)$. Let $f \in \mathbf{E}^\infty(a)$. There is some $m \in \mathbb{N}$ such that $fa^m = a^m$. If $l \in \mathbb{N}$ is such that $nl > m$, then $ef = e^l f = a^{nl} b^l f = a^{nl-m} a^m f b^l = a^{nl-m} a^m b^l = a^{nl} b^l = e^l = e$. \square

2. IDEAL SYSTEMS AND r -IDEAL SEMIGROUPS

Next we recall the most important facts about ideal systems. For a basic introduction into ideal systems see [15]. We briefly gather the required terminology. In the following a monoid is always a multiplicative commutative semigroup that possesses an identity element and a zero element and where every non-zero element is cancellative. A submonoid of a monoid H always contains the zero element and the identity element of H . A monoid K is called quotient monoid of a submonoid H if $K \setminus \{0\}$ is a quotient group of $H \setminus \{0\}$. Quotient monoids of monoids are essentially unique. A submonoid T of a quotient monoid of a monoid H is called overmonoid of H if $H \subseteq T$. For a monoid H let $H^\bullet = H \setminus \{0\}$.

Throughout this section, let H be a monoid and K a quotient monoid of H .

A subset $X \subseteq K$ is called fractional if there exists some $c \in H^\bullet$ such that $cX \subseteq H$. Let $\mathbf{F}(H)$ denote the set of all fractional subsets of K . We set $\mathbf{F}^\bullet(H) = \{I \in \mathbf{F}(H) \mid I \setminus \{0\} \neq \emptyset\}$. For arbitrary $X, Y \subseteq K$ let $(X : Y) = \{z \in K \mid zY \subseteq X\}$, $X^{-1} = (H : X)$ and $\mathcal{R}(X) = (X : X)$. Let \widehat{H} denote the complete integral closure of H . Note that for all $X \in \mathbf{F}^\bullet(H)$, $\mathcal{R}(X)$ is a submonoid of \widehat{H} . A subset $X \subseteq H$ is called multiplicatively closed if X contains the identity element and $XX \subseteq X$. A subset $\emptyset \neq P \subsetneq H$ is called prime ideal of H if $HP = P$ and $H \setminus P$ is multiplicatively closed. For a multiplicatively closed subset $T \subseteq H^\bullet$ let $T^{-1}H = \{t^{-1}x \mid t \in T, x \in H\}$ and for a prime ideal P of H let $H_P = (H \setminus P)^{-1}H$. If T and S are submonoids of K such that $T \subseteq S$, then let $\mathcal{F}_{S/T} = (T : S)$. Especially if T is an overmonoid of H , then $\mathcal{F}_{T/H} = T^{-1}$.

A map $r : \mathbf{F}(H) \rightarrow \mathbf{F}(H)$, $I \mapsto I_r$ is called ideal system on H , if for all $X, Y \in \mathbf{F}(H)$ and all $c \in K^\bullet$ it follows that:

- $X \cup \{0\} \subseteq X_r$.
- If $X \subseteq Y_r$, then $X_r \subseteq Y_r$.
- $(cX)_r = cX_r$.
- $H_r = H$.

Throughout this section let r be an ideal system on H .

Observe that for all $X, Y \in \mathbf{F}^\bullet(H)$ it follows that $(XY)_r = (X_r Y)_r = (X_r Y_r)_r$ and $(X_r : Y)_r = (X_r : Y) = (X_r : Y_r)$. (For a proof see [15, Propositions 2.3 and 11.7]). Furthermore it is straightforward to show that if $\mathcal{F}_{\widehat{H}/H} \neq \{0\}$, then $\widehat{H}, \mathcal{F}_{\widehat{H}/H} \in \mathcal{F}_r^\bullet(H)$. Next we introduce the most important ideal systems. Let $s : \mathbf{F}(H) \rightarrow \mathbf{F}(H)$ be defined by $s(\emptyset) = \{0\}$ and $s(I) = IH$ for all $I \in \mathbf{F}(H) \setminus \{\emptyset\}$.

Let $v : \mathbf{F}(H) \rightarrow \mathbf{F}(H)$ be defined by $I_v = (I^{-1})^{-1}$ for all $I \in \mathbf{F}(H)$.

If H is a domain, then let $d : \mathbf{F}(H) \rightarrow \mathbf{F}(H)$ be defined by $I_d = (I)_R$ for all $I \in \mathbf{F}(H)$.

Observe that s , v and d are ideal systems on H . Let $\mathcal{F}_r^\bullet(H) = \{I \in \mathbf{F}^\bullet(H) \mid I_r = I\}$. If $X \in \mathbf{F}(H)$, then let $\mathcal{R}_r(X) = \mathcal{R}(X_r)$. If r' is another ideal system on H , then we set $r \leq r'$ if $\mathcal{F}_{r'}^\bullet(H) \subseteq \mathcal{F}_r^\bullet(H)$ (equivalently: For all $I \in \mathbf{F}^\bullet(H)$ it follows that $I_r \subseteq I_{r'}$). r is called finitary if for all $X \in \mathbf{F}(H)$ it follows that $X_r = \bigcup_{F \subseteq X, |F| < \infty} F_r$. Let $\cdot_r : \mathcal{F}_r^\bullet(H) \times \mathcal{F}_r^\bullet(H) \rightarrow \mathcal{F}_r^\bullet(H)$ be defined by $I \cdot_r J = (IJ)_r$, for all $I, J \in \mathcal{F}_r^\bullet(H)$. Then $(\mathcal{F}_r^\bullet(H), \cdot_r)$ is a commutative semigroup, called the r -ideal semigroup.

We say that H is r -noetherian, if H satisfies one of the following equivalent conditions:

- Every ascending sequence of elements of $\{I \in \mathcal{F}_r^\bullet(H) \mid I \subseteq H\}$ becomes stationary.
- Every subset $\emptyset \neq \mathcal{M} \subseteq \{I \in \mathcal{F}_r^\bullet(H) \mid I \subseteq H\}$ has a maximal element.
- If $I \in \mathbf{F}^\bullet(H)$, then there exists some finite $E \subseteq I$ such that $I_r = E_r$.

If $r = v$, then these conditions are equivalent to each the following conditions:

- Every descending sequence of elements of $\mathcal{F}_r^\bullet(H)$ with non-zero intersection becomes stationary.
- Every subset $\emptyset \neq \mathcal{M} \subseteq \mathcal{F}_r^\bullet(H)$ with $\bigcap_{I \in \mathcal{M}} I \neq \{0\}$ has a minimal element.

For $I, J \in \mathcal{F}_r^\bullet(H)$, we set $I \sim_r J$ if there exists some $c \in K^\bullet$ such that $I = cJ$. It is easily checked that \sim_r is an equivalence relation on $\mathcal{F}_r^\bullet(H)$. Let $\mathcal{S}_r(H) = \mathcal{F}_r^\bullet(H) / \sim_r$. For $I \in \mathcal{F}_r^\bullet(H)$ we denote by $[I]_{\sim_r}$

the equivalence class of I . Let $\bullet_r : \mathcal{S}_r(H) \times \mathcal{S}_r(H) \rightarrow \mathcal{S}_r(H)$ be defined by $[I]_{\sim_r} \bullet_r [J]_{\sim_r} = [(IJ)_r]_{\sim_r}$, for all $I, J \in \mathcal{F}_r^\bullet(H)$. Then $(\mathcal{S}_r(H), \bullet_r)$ is a commutative semigroup, called the r -ideal class semigroup.

From now on we assume that $s \leq r$.

Let $r\text{-spec}(H) = \{P \in \mathcal{F}_r^\bullet(H) \mid P \text{ is a prime ideal of } H\}$ and let $r\text{-max}(H)$ be the set of all maximal elements of $\{I \in \mathcal{F}_r^\bullet(H) \mid I \subsetneq H, IH = I\}$. If r is finitary and $T \subseteq H^\bullet$ is multiplicatively closed, then let $T^{-1}r : \mathbf{F}(T^{-1}H) \rightarrow \mathbf{F}(T^{-1}H)$ be defined by $T^{-1}r(T^{-1}X) = T^{-1}X_r$, for all $X \in \mathbf{F}(H)$. Then $T^{-1}r$ is an ideal system on $T^{-1}H$. If P is a prime ideal of H , then let $r_P = (H \setminus P)^{-1}r$. If $H' \in \mathcal{F}_r^\bullet(H)$ is an overmonoid of H , then let $r[H'] : \mathbf{F}(H) \rightarrow \mathbf{F}(H)$ be defined by $r[H'](X) = (XH')_r$, for all $X \in \mathbf{F}(H)$. Observe that $r[H']$ is an ideal system on H' . Note that this construction is investigated in [16] in the more general context of module systems.

Next we introduce the concept of r -stability and πr -stability. Note that the r -stability of H is investigated in [11] for integral domains and in [18] for arbitrary monoids.

An element $I \in \mathcal{F}_r^\bullet(H)$ is called

- r -invertible if $(II^{-1})_r = H$.
- r -regular if I is regular as an element of $\mathcal{F}_r^\bullet(H)$.
- πr -regular if I is π -regular as an element of $\mathcal{F}_r^\bullet(H)$.
- r -stable if there exists some $J \in \mathcal{F}_r^\bullet(H)$ such that $(IJ)_r = \mathcal{R}(I)$ (equivalently: $(I(\mathcal{R}(I) : I))_r = \mathcal{R}(I)$).
- πr -stable if $(I^n)_r$ is r -stable for some $n \in \mathbb{N}$ (equivalently: $(I^n(\mathcal{R}_r(I^n) : I^n))_r = \mathcal{R}_r(I^n)$, for some $n \in \mathbb{N}$).

Note that every r -invertible element of $\mathcal{F}_r^\bullet(H)$ is r -stable and every r -stable element of $\mathcal{F}_r^\bullet(H)$ is r -regular. The monoid H is called

- r -stable if every $I \in \mathcal{F}_r^\bullet(H)$ is r -stable.
- πr -stable if every $I \in \mathcal{F}_r^\bullet(H)$ is πr -stable.

Lemma 2.1. *Let H be r -noetherian, T a submonoid of H and $H' \in \mathcal{F}_r^\bullet(H)$ an overmonoid of H .*

1. H' is $r[H']$ -noetherian.
2. $T^{-1}H$ is $T^{-1}r$ -noetherian.
3. For all $I, J \in \mathcal{F}_r^\bullet(H)$, $\bigcap_{P \in r\text{-max}(H)} I_P = I$ and $T^{-1}(I : J) = (T^{-1}I : T^{-1}J)$.

Proof. 1. Let $I \in \mathbf{F}^\bullet(H') = \mathbf{F}^\bullet(H)$. Then there exists some finite $E \subseteq I$ such that $E_r = I_r$. This implies that $E_{r[H']} = (EH')_r = (E_r H')_r = (I_r H')_r = (IH')_r = I_{r[H']}$. Therefore it follows that H' is $r[H']$ -noetherian.

2. This follows by [15, Theorem 4.4].

3. Let $I, J \in \mathcal{F}_r^\bullet(H)$. It follows by [15, Theorem 11.3] that $\bigcap_{P \in r\text{-max}(H)} I_P = I$ and it follows by [15, Proposition 11.7] that $T^{-1}(I : J) = (T^{-1}I : T^{-1}J)$. \square

In the following we investigate the r -ideal semigroup of H with respect to the properties defined in Section 1. Note that $\mathcal{R}(I) \in \mathbf{E}(I) \subseteq \mathbf{E}(\mathcal{F}_r^\bullet(H))$ for all $I \in \mathcal{F}_r^\bullet(H)$. Therefore $\{I \in \mathcal{F}_r^\bullet(H) \mid I = \mathcal{R}(I)\} \subseteq \mathbf{E}(\mathcal{F}_r^\bullet(H))$ and it is natural to ask, when equality holds. In this context it is worth remarking, that the structure of the idempotents of the t -ideal class semigroup is studied in [22].

Lemma 2.2. *Let $I \in \mathcal{F}_r^\bullet(H)$.*

1. If $J \in \mathcal{F}_r^\bullet(H)$ is such that $(IJ)_r = I$, then $(II^{-1})_r \subseteq (I(J : I))_r \subseteq J \subseteq \mathcal{R}(I)$ and $\mathcal{F}_{\mathcal{R}(I)/H} \subseteq J^{-1} \subseteq \mathcal{R}(I)$.
2. If $(I^2 I^{-1})_r = I$, then I is r -regular.
3. If $E \in \mathbf{E}(I)$, then $(I(E : I))_r \in \{J \in \mathcal{F}_r^\bullet(H) \mid (II^{-1})_r \subseteq J \subseteq \mathcal{R}(I), E \in \mathbf{E}(J) \text{ and } J^2 \subseteq J\}$.
4. If $\mathbf{E}^\infty(I)$ possesses a least element, then there exists some $n \in \mathbb{N}$ such that $\mathcal{R}_r(I^k) = \mathcal{R}_r(I^n)$ for all $k \in \mathbb{N}_{\geq n}$.

5. Assume that $n \in \mathbb{N}$, $E \in \mathbf{E}((I^n)_r)$, and let $J \in \mathcal{F}_r^\bullet(H)$ be such that $(I^n J)_r = E$. Then it follows that $(I^k(\mathcal{R}_r(I^k):I^k))_r = E$ and $\mathcal{R}((\mathcal{R}_r(I^k):I^k)) = \mathcal{R}_r(I^k) = \mathcal{R}(E)$ for all $k \in \mathbb{N}_{\geq n}$.

Proof. 1. Let $J \in \mathcal{F}_r^\bullet(H)$ be such that $(IJ)_r = I$. Then $IJ \subseteq (IJ)_r = I$, hence $J \subseteq \mathcal{R}(I)$. Of course, $J^{-1} = (H : J) \subseteq (IH : IJ) \subseteq ((IH)_r : (IJ)_r) = (I : I) = \mathcal{R}(I)$. Clearly $I^{-1} = (H : I) \subseteq (HJ : IJ) \subseteq ((HJ)_r : (IJ)_r) = (J : I)$ and hence $(II^{-1})_r \subseteq (I(J : I))_r \subseteq J$. Since $J \subseteq \mathcal{R}(I)$, it follows that $\mathcal{F}_{\mathcal{R}(I)/H} = (H : \mathcal{R}(I)) \subseteq (H : J) = J^{-1}$.

2. Let $(I^2 I^{-1})_r = I$. Since $I^{-1} \in \mathcal{F}_r^\bullet(H)$, it follows that I is r -regular.

3. Let $E \in \mathbf{E}(I)$. Of course, $(I(E : I))_r \in \mathcal{F}_r^\bullet(H)$ and by 1. it follows that $(II^{-1})_r \subseteq (I(E : I))_r \subseteq \mathcal{R}(I)$. Since $E \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$ and $((I(E : I))_r E)_r = (IE(E : I))_r = ((IE)_r(E : I))_r = (I(E : I))_r$, it follows that $E \in \mathbf{E}((I(E : I))_r)$. Finally, it follows that $((I(E : I))_r)^2 \subseteq (((I(E : I))_r)^2)_r = (I^2(E : I)^2)_r \subseteq (I(E : I)E)_r = (I(E : I))_r$.

4. Let E be a least element of $\mathbf{E}^\infty(I)$. There exists some $n \in \mathbb{N}$ such that $(EI^n)_r = (I^n)_r$. Let $k \in \mathbb{N}_{\geq n}$. Then there exists some $l \in \mathbb{N}$ such that $ln \geq k$. Since $\mathcal{R}_r(I^{ln}) \in \mathbf{E}((I^{ln})_r) \subseteq \mathbf{E}^\infty(I)$, it follows that $E \leq \mathcal{R}_r(I^{ln})$. This implies that $(E\mathcal{R}_r(I^{ln}))_r = E$, hence $\mathcal{R}_r(I^{ln}) \subseteq \mathcal{R}(E)$. Therefore it follows that $\mathcal{R}(E) \subseteq \mathcal{R}_r(EI^n) = \mathcal{R}_r(I^n) \subseteq \mathcal{R}_r(I^k) \subseteq \mathcal{R}_r(I^{ln}) \subseteq \mathcal{R}(E)$, hence $\mathcal{R}_r(I^k) = \mathcal{R}_r(I^n)$.

5. Let $n \in \mathbb{N}$, $k \in \mathbb{N}_{\geq n}$, $E \in \mathbf{E}((I^n)_r)$, and $J \in \mathcal{F}_r^\bullet(H)$ be such that $(I^n J)_r = E$. There exists some $l \in \mathbb{N}$ such that $ln \geq k$. It follows that $\mathcal{R}_r(I^n) \subseteq \mathcal{R}_r(I^k) \subseteq \mathcal{R}_r(I^{ln}) \subseteq \mathcal{R}_r(I^{ln} J^l) = \mathcal{R}_r(E^l) = \mathcal{R}(E) \subseteq \mathcal{R}_r(EI^n) = \mathcal{R}_r(I^n)$, hence $\mathcal{R}_r(I^{ln}) = \mathcal{R}_r(I^k) = \mathcal{R}_r(I^n) = \mathcal{R}(E)$. It follows that $E = (I^n J)_r \subseteq (I^n(E : I^n))_r \subseteq E$, hence $(I^n(E : I^n))_r = E$. Since $(E : I^n) = (E : (I^n)_r) = (E : (EI^n)_r) = (E : EI^n) = (\mathcal{R}(E) : I^n) = (\mathcal{R}_r(I^n) : I^n)$, it follows that $(I^n(\mathcal{R}_r(I^n) : I^n))_r = E$. This implies that $E = (E^l)_r = (I^{ln}(\mathcal{R}_r(I^n) : I^n)^l)_r \subseteq (I^{ln}(\mathcal{R}_r(I^n) : I^{ln}))_r \subseteq (I^k(\mathcal{R}_r(I^n) : I^k))_r \subseteq (I^n(\mathcal{R}_r(I^n) : I^n))_r = E$, hence $(I^k(\mathcal{R}_r(I^k) : I^k))_r = (I^k(\mathcal{R}_r(I^n) : I^k))_r = E$. Finally, this implies that $\mathcal{R}((\mathcal{R}_r(I^k) : I^k)) = (\mathcal{R}_r(I^k) : I^k(\mathcal{R}_r(I^k) : I^k)) = (\mathcal{R}(E) : E) = \mathcal{R}(E)$. \square

Theorem 2.3.

1. The following conditions are equivalent:
 - a. If $I \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$, then $I = \mathcal{R}(I)$ (equivalently: $\mathbf{E}(\mathcal{F}_r^\bullet(H)) = \{I \in \mathcal{F}_r^\bullet(H) \mid I = \mathcal{R}(I)\}$).
 - b. For all $E, F \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$, $E \leq F$ implies $F \subseteq E$.
 - c. For all $E, F \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$, $F \subseteq E$ implies $E \leq F$.
 - d. For all $I \in \mathcal{F}_r^\bullet(H)$ we have $\min(\mathbf{E}(I)) = \mathcal{R}(I)$.

If these equivalent conditions are satisfied, then $\mathcal{F}_r^\bullet(H)$ is almost complete.

2. Let $I, E \in \mathcal{F}_r^\bullet(H)$. Then the following conditions are equivalent:
 - a. E is a least element of $\mathbf{E}(I)$ with respect to the Rees-order.
 - b. E is a least element of $\{F \in \mathbf{E}(I) \mid F \leq \mathcal{R}(I)\}$ with respect to inclusion.
 - c. E is a minimal element of $\{F \in \mathbf{E}(I) \mid F \leq \mathcal{R}(I)\}$ with respect to inclusion.
3. The following conditions are equivalent:
 - a. H is πr -stable.
 - b. $\mathcal{F}_r^\bullet(H)$ is π -regular and $\mathbf{E}(\mathcal{F}_r^\bullet(H)) = \{E \in \mathcal{F}_r^\bullet(H) \mid E = \mathcal{R}(E)\}$.
4. Suppose that $\mathbf{E}(\mathcal{F}_r^\bullet(H)) = \{I \in \mathcal{F}_r^\bullet(H) \mid I = \mathcal{R}(I)\}$ and let $I \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$. Then $I \cdot_r \mathcal{F}_r^\bullet(H) = \{J \in \mathcal{F}_r^\bullet(H) \mid I \subseteq \mathcal{R}(J)\}$ and $P_I^* = \{J \in \mathcal{F}_r^\bullet(H) \mid I = \mathcal{R}(J)\}$.

Proof. 1. a. \Rightarrow b. If $E, F \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$ and $E \leq F$, then $(EF)_r = F$. Hence $F \subseteq (F\mathcal{R}(E))_r = (FE)_r = E$.

a. \Rightarrow c. If $E, F \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$ and $E \leq F$, then $(EF)_r \subseteq (E^2)_r = E$ and $E \subseteq (E\mathcal{R}(F))_r = (EF)_r$. This implies that $(EF)_r = E$, hence $E \leq F$.

b. \Rightarrow d. Let $I \in \mathcal{F}_r^\bullet(H)$ and $F \in \mathbf{E}(I)$. Then we have to show that $(F\mathcal{R}(I))_r = \mathcal{R}(I)$. It follows that $(F\mathcal{R}(I))_r \in \mathbf{E}(I)$ and $(F\mathcal{R}(I))_r \leq \mathcal{R}(I)$, hence $\mathcal{R}(I) \subseteq (F\mathcal{R}(I))_r$. Since $F \in \mathbf{E}(I)$, it follows by Lemma 2.2.1. that $F \subseteq \mathcal{R}(I)$, hence $(F\mathcal{R}(I))_r \subseteq (\mathcal{R}(I)^2)_r = \mathcal{R}(I)$. Finally, this implies that $(F\mathcal{R}(I))_r = \mathcal{R}(I)$.

c. \Rightarrow a. Let $E \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$. Then $E \subseteq \mathcal{R}(E)$ by Lemma 2.2.1., hence $\mathcal{R}(E) \leq E$. Since $(E\mathcal{R}(E))_r = E$,

it follows that $E \leq \mathcal{R}(E)$, hence $E = \mathcal{R}(E)$.

d. \Rightarrow a. Let $E \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$. Then $E \in \mathbf{E}(E)$, hence $\mathcal{R}(E) \leq E$. Since $(E\mathcal{R}(E))_r = E$, it follows that $E \leq \mathcal{R}(E)$, hence $E = \mathcal{R}(E)$.

2. a. \Rightarrow b. Of course, $E \in \mathbf{E}(I)$. Since $\mathcal{R}(I) \in \mathbf{E}(I)$, it follows that $E \leq \mathcal{R}(I)$. Let $F \in \mathbf{E}(I)$ be such that $F \leq \mathcal{R}(I)$. It follows that $(EF)_r \in \mathbf{E}(I)$ and $(EF)_r \leq E$, hence $E = (EF)_r$. By Lemma 2.2.1. it follows that $E \subseteq \mathcal{R}(I)$, hence $E = (EF)_r \subseteq (\mathcal{R}(I)F)_r = F$.

b. \Rightarrow c. Trivial.

c. \Rightarrow a. Let $F \in \mathbf{E}(I)$. Then it is sufficient to show that $(EF)_r = E$. It follows that $(EF)_r \leq E \leq \mathcal{R}(I)$ and $(EF)_r \in \mathbf{E}(I)$, hence $(EF)_r \in \{G \in \mathbf{E}(I) \mid G \leq \mathcal{R}(I)\}$. By Lemma 2.2.1. $F \subseteq \mathcal{R}(I)$, hence $(EF)_r \subseteq (E\mathcal{R}(I))_r = E$. This implies that $E = (EF)_r$.

3. a. \Rightarrow b. Let $I \in \mathcal{F}_r^\bullet(H)$. Then there is some $n \in \mathbb{N}$ such that $(I^n(\mathcal{R}_r(I^n) : I^n))_r = \mathcal{R}_r(I^n)$. Since $\mathcal{R}_r(I^n) \in \mathbf{E}((I^n)_r)$, it follows that I is πr -regular. This implies that $\mathcal{F}_r^\bullet(H)$ is π -regular. Let $F \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$. There is some $n \in \mathbb{N}$ such that $(F^n(\mathcal{R}_r(F^n) : F^n))_r = \mathcal{R}_r(F^n)$, hence $F = (F\mathcal{R}(F))_r = (F(\mathcal{R}(F) : F))_r = (F^n(\mathcal{R}_r(F^n) : F^n))_r = \mathcal{R}_r(F^n) = \mathcal{R}(F)$.

b. \Rightarrow a. Let $I \in \mathcal{F}_r^\bullet(H)$. Since I is πr -regular, there exist some $n \in \mathbb{N}$, $E \in \mathbf{E}((I^n)_r)$ and $J \in \mathcal{F}_r^\bullet(H)$ such that $(I^n J)_r = E$. Consequently, Lemma 2.2.5. implies that $(I^n(\mathcal{R}_r(I^n) : I^n))_r = E = \mathcal{R}(E) = \mathcal{R}_r(I^n)$.

4. Let us show the first equality. “ \subseteq ”: Let $J \in I \cdot_r \mathcal{F}_r^\bullet(H)$. Then there exists some $A \in \mathcal{F}_r^\bullet(H)$ such that $J = (IA)_r$. It is clear that $IJ \subseteq (IJ)_r = (I^2 A)_r = ((I^2)_r A)_r = (IA)_r = J$, hence $I \subseteq \mathcal{R}(J)$. “ \supseteq ”: Let $J \in \mathcal{F}_r^\bullet(H)$ be such that $I \subseteq \mathcal{R}(J)$. It follows that $J \subseteq JI \subseteq \mathcal{J}\mathcal{R}(J) = J$, hence $J = J_r = (IJ)_r = I \cdot_r J \in I \cdot_r \mathcal{F}_r^\bullet(H)$. Now we show the second equality. It follows by 1. that $\bigcup_{A \in \mathbf{E}(\mathcal{F}_r^\bullet(H)), A < I} A \cdot_r \mathcal{F}_r^\bullet(H) = \bigcup_{A \in \mathbf{E}(\mathcal{F}_r^\bullet(H)), I \not\subseteq A} \{J \in \mathcal{F}_r^\bullet(H) \mid A \subseteq \mathcal{R}(J)\} = \{J \in \mathcal{F}_r^\bullet(H) \mid \text{there exists some } A \in \mathbf{E}(\mathcal{F}_r^\bullet(H)) \text{ such that } I \not\subseteq A \subseteq \mathcal{R}(J)\} = \{J \in \mathcal{F}_r^\bullet(H) \mid I \not\subseteq \mathcal{R}(J)\}$. This implies that $P_I^* = \{J \in \mathcal{F}_r^\bullet(H) \mid I \subseteq \mathcal{R}(J)\} \setminus \{J \in \mathcal{F}_r^\bullet(H) \mid I \not\subseteq \mathcal{R}(J)\} = \{J \in \mathcal{F}_r^\bullet(H) \mid I = \mathcal{R}(J)\}$. \square

It is not difficult to see that algebraic properties of the r -ideal semigroup $\mathcal{F}_r^\bullet(H)$ induce corresponding properties of the r -ideal class semigroup $\mathcal{S}_r(H)$. Obviously $\mathbf{E}(\mathcal{S}_r(H)) = \{[I]_{\sim_r} \mid I \in \mathbf{E}(\mathcal{F}_r^\bullet(H))\}$ and if $I, J \in \mathbf{E}(\mathcal{F}_r^\bullet(H))$, then $[I]_{\sim_r} \leq [J]_{\sim_r}$ holds if and only if $I \leq J$. Moreover if $I \in \mathcal{F}_r^\bullet(H)$, then

- $P_{[I]_{\sim_r}}^* = \{[J]_{\sim_r} \mid J \in P_I^*\}$.
- I is $(\pi)r$ -regular if and only if $[I]_{\sim_r}$ is a (π) -regular element of $\mathcal{S}_r(H)$.

Consequently,

- $\mathcal{S}_r(H)$ is regular[π -regular, almost complete, complete] if and only if $\mathcal{F}_r^\bullet(H)$ is regular[π -regular, almost complete, complete].

Moreover let r' be an ideal system on H satisfying $r \leq r'$.

- If $\mathcal{F}_r^\bullet(H)$ is regular[π -regular], then $\mathcal{F}_{r'}^\bullet(H)$ is regular[π -regular].
- If H is r -stable[πr -stable], then H is r' -stable[$\pi r'$ -stable].

For an abstract semigroup-theoretic formalism behind these statements, see [18, Lemma 2.2].

Lemma 2.4.

1. If $I \in \mathcal{F}_v^\bullet(H)$, then $(II^{-1})_v = \mathcal{F}_{\mathcal{R}(I)/H}$, and if $I \in \mathbf{E}(\mathcal{F}_v^\bullet(H))$, then $\mathcal{F}_{\widehat{H}/H} \subseteq I \subseteq \widehat{H}$ and $\mathcal{F}_{\widehat{H}/H} \subseteq I^{-1} \subseteq \widehat{H}$.
2. If $I \in \mathcal{F}_v^\bullet(H)$ and $(I\mathcal{F}_{\mathcal{R}(I)/H})_v = I$, then I is v -regular.
3. Let \widehat{H} be factorial and $I \in \mathcal{F}_v^\bullet(H)$. Then there exists some $c \in K^\bullet$ such that $((\mathcal{F}_{\widehat{H}/H})^2)_v \subseteq cI \subseteq \widehat{H}$.

Proof. **1.** If $I \in \mathcal{F}_v^\bullet(H)$, then $(II^{-1})_v = (H : (H : II^{-1})) = (H : ((H : I^{-1}) : I)) = (H : (I : I)) = (H : \mathcal{R}(I)) = \mathcal{F}_{\mathcal{R}(I)/H}$. Let $I \in \mathbf{E}(\mathcal{F}_v^\bullet(H))$. Then it follows by Lemma 2.2.1. that $\mathcal{F}_{\widehat{H}/H} \subseteq \mathcal{F}_{\mathcal{R}(I)/H} = (II^{-1})_v \subseteq I \subseteq \mathcal{R}(I) \subseteq \widehat{H}$ and $\mathcal{F}_{\widehat{H}/H} \subseteq \mathcal{F}_{\mathcal{R}(I)/H} \subseteq I^{-1} \subseteq \mathcal{R}(I) \subseteq \widehat{H}$.

2. Let $I \in \mathcal{F}_v^\bullet(H)$ be such that $(I\mathcal{F}_{\mathcal{R}(I)/H})_v = I$. Then it follows by 1. that $(I^2I^{-1})_v = (I(II^{-1})_v)_v = (I\mathcal{F}_{\mathcal{R}(I)/H})_v = I$. Therefore Lemma 2.2.2. implies that I is v -regular.
3. Since \widehat{H} is factorial, it follows that every $J \in \mathcal{F}_{v\widehat{H}}^\bullet(\widehat{H})$ is principal. Since $(\widehat{H}:I) \in \mathcal{F}_{v\widehat{H}}^\bullet(\widehat{H})$, there exists some $c \in K^\bullet$ such that $(\widehat{H}:I) = c\widehat{H}$. By 1. it follows that $\mathcal{F}_{\widehat{H}/H} \subseteq \mathcal{F}_{\mathcal{R}(I)/H} = (II^{-1})_v \subseteq (I(\widehat{H}:I))_v = (Ic\widehat{H})_v$. This implies that $((\mathcal{F}_{\widehat{H}/H})^2)_v \subseteq ((Ic\widehat{H})_v\mathcal{F}_{\widehat{H}/H})_v = c(I\mathcal{F}_{\widehat{H}/H})_v \subseteq cI \subseteq cI\widehat{H} = I(\widehat{H}:I) \subseteq \widehat{H}$. \square

Lemma 2.4. shows that every idempotent of the v -ideal semigroup contains the conductor. In Section 5 we present several examples of integral domains, where the conductor itself is an idempotent of the v -ideal semigroup. In the following we summarize some properties of the v -ideal semigroup of v -noetherian monoids and consider the extremal case that the conductor is non-trivial and v -idempotent.

Lemma 2.5. *Let H be v -noetherian and $\mathcal{F}_{\widehat{H}/H} \in \mathbf{E}(\mathcal{F}_v^\bullet(H))$.*

1. *Let $I \in \mathcal{F}_v^\bullet(H)$ be such that $\mathcal{F}_{\widehat{H}/H} \subseteq I \subseteq \widehat{H}$. If there exist some $k, l \in \mathbb{N}$ such that $k \neq l$ and $(I^k)_v \subseteq (I^l)_v$, then I is πv -regular.*
2. *Let \widehat{H} be factorial and $I \in \mathcal{F}_v^\bullet(H)$ such that $\mathcal{R}(I) = \widehat{H}$. Then I is πv -regular.*

Proof. 1. Let $k, l \in \mathbb{N}$ be such that $k \neq l$ and $(I^k)_v \subseteq (I^l)_v$.

Case1: $k < l$: Since $\mathcal{F}_{\widehat{H}/H} \neq \{0\}$, we have $\widehat{H} = (\mathcal{F}_{\widehat{H}/H})^{-1} \in \mathcal{F}_v^\bullet(H)$. If $0 \neq c \in \mathcal{F}_{\widehat{H}/H}$ and $r \in \mathbb{N}$, then $c(I^{k+r(l-k)})_v \subseteq c(I^{k+(r+1)(l-k)})_v \subseteq c(\widehat{H}^{k+(r+1)(l-k)})_v = c\widehat{H} \subseteq H$. Since H is v -noetherian, there is some $n \in \mathbb{N}$ such that $c(I^{k+n(l-k)})_v = c(I^{k+(n+1)(l-k)})_v$. This implies that $(I^{k+n(l-k)})_v = (I^{k+(n+1)(l-k)})_v$ and consequently Lemma 1.1.2. implies that I is πv -regular.

Case2: $l < k$: If $r \in \mathbb{N}$, then $(I^{l+(r+1)(k-l)})_v \subseteq (I^{l+r(k-l)})_v$. This implies that $\{0\} \neq \mathcal{F}_{\widehat{H}/H} = ((\mathcal{F}_{\widehat{H}/H})^{l+(r+1)(k-l)})_v \subseteq (I^{l+(r+1)(k-l)})_v \subseteq (I^{l+r(k-l)})_v$. Since H is v -noetherian it follows that there exists some $n \in \mathbb{N}$ such that $(I^{l+(n+1)(k-l)})_v = (I^{l+n(k-l)})_v$. Therefore Lemma 1.1.2. implies that I is πv -regular.

2. Of course, $(I\widehat{H})_v = (I\mathcal{R}(I))_v = I_v = I$. It follows by Lemma 2.4.3. that there exists some $c \in K^\bullet$ such that $\mathcal{F}_{\widehat{H}/H} \subseteq cI \subseteq \widehat{H}$. Since $(cI)^2 \subseteq cI\widehat{H} \subseteq c(I\widehat{H})_v = cI$, it follows by 1. that cI is πv -regular. This implies that I is πv -regular. \square

Theorem 2.6. *Let H be v -noetherian.*

1. *$\mathcal{F}_v^\bullet(H)$ is almost complete.*
2. *If $I \in \mathcal{F}_v^\bullet(H)$ and if there exists some $n \in \mathbb{N}$ such that $\mathcal{R}_v(I^n) = \mathcal{R}_v(I^k)$ for all $k \in \mathbb{N}_{\geq n}$, then $\mathbf{E}^\infty(I)$ has a least element.*
3. *If \widehat{H} is factorial, $\mathcal{F}_{\widehat{H}/H} \in \mathbf{E}(\mathcal{F}_v^\bullet(H))$ and $\{J \in \mathcal{F}_v^\bullet(H) \mid J = \mathcal{R}(J)\} = \{H, \widehat{H}\}$, then $\mathcal{F}_v^\bullet(H)$ is complete.*

Proof. 1. Let $I \in \mathcal{F}_v^\bullet(H)$ and $M = \{F \in \mathbf{E}(I) \mid F \leq \mathcal{R}(I)\}$. Since $\mathcal{R}(I) \in M$, it follows that $M \neq \emptyset$. It follows by Lemma 2.2.1. and Lemma 2.4.1. that $\bigcap_{B \in M} B \supseteq \mathcal{F}_{\mathcal{R}(I)/H} \neq \{0\}$. Since H is v -noetherian, there is some $E \in M$ that is minimal with respect to inclusion. Finally, it follows by Theorem 2.3.2. that E is a least element of $\mathbf{E}(I)$ with respect to the Rees-order.

2. Let $I \in \mathcal{F}_v^\bullet(H)$ and $n \in \mathbb{N}$ such that $\mathcal{R}_v(I^k) = \mathcal{R}_v(I^n)$ for all $k \in \mathbb{N}_{\geq n}$. Let $M = \{F \in \mathbf{E}^\infty(I) \mid F \leq \mathcal{R}_v(I^n)\}$. Since $\mathcal{R}_v(I^n) \in \mathbf{E}((I^n)_v) \subseteq \mathbf{E}^\infty(I)$, it follows that $M \neq \emptyset$. Let us show that $\bigcap_{B \in M} B \supseteq \mathcal{F}_{\mathcal{R}_v(I^n)/H}$. Let $B \in M$. Then there is some $k \in \mathbb{N}_{\geq n}$ such that $B \in \mathbf{E}((I^k)_v)$. It follows by Lemma 2.2.1. and Lemma 2.4.1. that $B \supseteq \mathcal{F}_{\mathcal{R}_v(I^k)/H} = \mathcal{F}_{\mathcal{R}_v(I^n)/H}$. This implies that $\bigcap_{B \in M} B \neq \{0\}$, hence (since H is v -noetherian) there exists some $E \in M$ that is minimal with respect to inclusion. Let $F \in \mathbf{E}^\infty(I)$. Then we have to show that $(EF)_v = E$. There exists some $k \in \mathbb{N}_{\geq n}$ such that $F \in \mathbf{E}((I^k)_v)$, hence Lemma 2.2.1. implies that $(EF)_v \subseteq (E\mathcal{R}_v(I^k))_v = (E\mathcal{R}_v(I^n))_v = E$. It follows that $(EF)_v \in \mathbf{E}^\infty(I)$

and $(EF)_v \leq E \leq \mathcal{R}_v(I^n)$, hence $(EF)_v \in M$ and $(EF)_v = E$.

3. Let \widehat{H} be factorial, $\mathcal{F}_{\widehat{H}/H} \in \mathbf{E}(\mathcal{F}_v^\bullet(H))$ and $\{J \in \mathcal{F}_v^\bullet(H) \mid J = \mathcal{R}(J)\} = \{H, \widehat{H}\}$. By 1. it follows that $\mathcal{F}_v^\bullet(H)$ is almost complete. Let $I \in \mathcal{F}_v^\bullet(H)$. Then $\mathcal{R}(I) \in \{H, \widehat{H}\}$.

Case1: $\mathcal{R}(I) = H$: Since $\mathcal{F}_{\mathcal{R}(I)/H} = H \in \mathbf{E}(I)$, it follows by Lemma 2.4.2. that I is v -regular. Case2: $\mathcal{R}(I) = \widehat{H}$: It follows by Lemma 2.5.2. that I is πv -regular. \square

Note that Theorem 2.6.2. proves the converse of Lemma 2.2.4., if $r = v$ and H is v -noetherian. We were not able to decide whether the r -ideal semigroup of a r -noetherian monoid is almost complete. Theorem 2.6.3. gives a hint how to construct examples of monoids which are not πv -stable but whose v -ideal semigroup is π -regular.

3. PREPARATIONS FOR THE MAIN RESULTS

Let R be an integral domain and K a field of quotients of R . By $\mathcal{F}^\bullet(R)$ we denote the set of all non-zero fractional ideals of R and by \overline{R} we denote the integral closure of R in K . R is called G -domain if $\bigcap_{P \in \text{spec}(R), P \neq \{0\}} P \neq \{0\}$. An ideal system r on R is called (extended) star operation on R if $d \leq r$. Note that an ideal system r on R is a star operation if and only if $r|_{\mathcal{F}^\bullet(R)}$ is a star operation on R in the sense of [13]. Observe that for all $X \in \mathcal{F}^\bullet(R)$, $\mathcal{R}(X)$ is an intermediate ring of R and \widehat{R} .

Lemma 3.1. *Let R be a Mori domain, $x \in R^\bullet$ and $S \subseteq R^\bullet$ a multiplicatively closed set.*

1. $\text{spec}^1(R) \subseteq v\text{-spec}(R)$, and $\{P \in v\text{-spec}(R) \mid x \in P\}$ is finite.
2. $S^{-1}\widehat{R} = \widehat{S^{-1}R}$.
3. If R is not a field, then $R \setminus R^\times = \bigcup_{M \in v\text{-max}(R)} M$.
4. If $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$, then \widehat{R} is a Krull domain and $S^{-1}\mathcal{F}_{\widehat{R}/R} = \mathcal{F}_{\widehat{S^{-1}R}/S^{-1}R}$.
5. If R is local, $\dim(R) = 1$ and \widehat{R} is a Krull domain, then \widehat{R} is a semilocal principal ideal domain.

Proof. By $\text{spec}^1(R)$ we denote the set of height-one prime ideals of R . Observe that R is a Mori domain if and only if R^\bullet is a v -noetherian monoid in the sense of [12]. Note that if \widehat{R}^\bullet denotes the complete integral closure of R^\bullet in its quotient group, then $\widehat{R}^\bullet = \widehat{R}^\bullet$.

1. The first assertion follows by [12, Proposition 2.2.4.2.], and the second assertion follows by [12, Theorem 2.2.5.1.].
2. See [12, Theorem 2.3.5.2.].
3. Let R be not a field. The claim is an immediate consequence of [12, Proposition 2.2.4.1.].
4. Let $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$. The first assertion follows by [12, Theorem 2.3.5.3.], and the second assertion follows by 2. and [12, Proposition 2.2.8.1.].
5. Let R be local, $\dim(R) = 1$ and \widehat{R} a Krull domain and let M be the maximal ideal of R . It is an immediate consequence of [12, Proposition 2.10.5.1.(c)] that $\dim(\widehat{R}) = 1$. Therefore \widehat{R} is a Dedekind domain. Of course, $M' \cap R = M$ for all $M' \in \text{max}(\widehat{R})$, hence $\text{max}(\widehat{R}) = \mathcal{P}(M\widehat{R})$. Since \widehat{R} is noetherian this implies that $\text{max}(\widehat{R})$ is finite. Therefore \widehat{R} is a semilocal Dedekind domain, hence \widehat{R} is a semilocal principal ideal domain. \square

Lemma 3.2 (Prime avoidance lemma). *Let S be a commutative ring with identity and $I \subseteq S$ an additively closed subset.*

For $k \in \mathbb{N}$, let $I^{(k)} = \{\prod_{i=1}^k z_i \mid (z_i)_{i=1}^k \in I^{[1,k]}\}$ and $I^k = \{\sum_{i=1}^m z_i \mid m \in \mathbb{N}, (z_i)_{i=1}^m \in (I^{(k)})^{[1,m]}\}$.

Let $n \in \mathbb{N}$ and $(P_i)_{i=1}^n \in \text{spec}(S)^{[1,n]}$ be such that $I \not\subseteq P_i$ for all $i \in [1, n]$. Then it follows that $I^{(n-1)!} \not\subseteq \bigcup_{i=1}^n P_i$.

Proof. We use induction on n . The assertion is obvious for $n = 1$. $n \rightarrow n + 1$: Let $n \in \mathbb{N}$ and $(P_i)_{i=1}^{n+1} \in \text{spec}(S)^{[1, n+1]}$ be such that $I \not\subseteq P_i$ for all $i \in [1, n+1]$. If $i \in [1, n+1]$, then $I^{(n-1)!} \not\subseteq \bigcup_{j=1, j \neq i}^{n+1} P_j$ (by the induction hypothesis), and thus there exists some $(y_i)_{i=1}^{n+1} \in S^{[1, n+1]}$ such that $y_i \in I^{(n-1)!} \setminus \bigcup_{j=1, j \neq i}^{n+1} P_j$ for all $i \in [1, n+1]$. Now we set $y = y_{n+1}^n + \prod_{i=1}^n y_i \in I^{n!}$ and assume (contrary to our assertion) that $I^{n!} \subseteq \bigcup_{i=1}^{n+1} P_i$. For every $i \in [1, n+1]$ we have $y_i^n \in I^{n!} \subseteq \bigcup_{j=1}^{n+1} P_j$ and thus $y_i \in \bigcup_{j=1}^{n+1} P_j$, hence $y_i \in P_i$. Since $y \in I^{n!} \subseteq \bigcup_{i=1}^{n+1} P_i$, there exists some $m \in [1, n+1]$ such that $y \in P_m$.

Case1: $m \in [1, n]$: Since $y_m \in P_m$ it follows that $\prod_{i=1}^n y_i \in P_m$. Since $y \in P_m$, we have $y_{n+1}^n \in P_m$, hence $y_{n+1} \in P_m$, a contradiction.

Case2: $m = n + 1$: Since $y_{n+1} \in P_{n+1}$ and $y \in P_{n+1}$, it follows that $\prod_{i=1}^n y_i \in P_{n+1}$, hence there exists some $l \in [1, n]$ such that $y_l \in P_{n+1}$, a contradiction. \square

4. MAIN RESULTS

In [9] it was shown that orders in quadratic number fields are (d -)stable. This result is based on two other results. The first one states that the ideal semigroup of a noetherian 1-dimensional integral domain is πd -stable. The second result gives a common upper bound on the "index of stability" of all elements I (i.e. the smallest $n \in \mathbb{N}$ for which I^n is stable) of the ideal semigroup. The main goal of this paper is to extend the first result to arbitrary $*$ -ideal semigroups. As already shown in Section 2 there exists a close connection between $\pi*$ -stability and the structure of idempotents of the $*$ -ideal semigroup. As pointed out in [9] and [17] the idempotents of the ideal semigroup of a noetherian integral domain contain the identity. Unfortunately, there is no analogous result for general $*$ -ideal semigroups. In Section 5 it will be shown that the idempotents of the v -ideal semigroup of a Mori domain need not contain the identity. It is natural to ask for conditions on R that enforce the $*$ -idempotents to be "trivial" (i.e. they contain the identity). The first main result and its corollary deal with this problem.

Theorem 4.1. *Let R be an integral domain, K a field of quotients of R , $*$ a star operation on R and $I, J \in \mathcal{F}_*^\bullet(R)$ such that $(IJ)_* = I$.*

1. *If $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$ and \widehat{R} is a Dedekind domain, then $J\widehat{R} = \widehat{R}$.*
2. *Suppose that one of the following conditions is satisfied:*
 - a. *$\mathcal{F}_{\widehat{R}/R} \neq \{0\}$ and \overline{R} is a semilocal principal ideal domain.*
 - b. *R is a $*$ -noetherian G -domain.*
 - c. *R is $*$ -noetherian and $\dim(R) = 1$.*

Then $J\mathcal{R}(I) = \mathcal{R}(I)$.

3. *If R is $*$ -noetherian and $*$ -max(R) = $\text{spec}^1(R)$, then $(J\mathcal{R}(I))_* = \mathcal{R}(I)$.*

Proof. 1. Let $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$, and \widehat{R} a Dedekind domain. Of course, $I\widehat{R}, IJ\widehat{R} \in \mathcal{F}^\bullet(\widehat{R})$. Since $I\widehat{R}$ and $IJ\widehat{R}$ are invertible this implies that $(I\widehat{R})_{v_{\widehat{R}}} = I\widehat{R}$ and $(IJ\widehat{R})_{v_{\widehat{R}}} = IJ\widehat{R}$. It follows that $IJ\widehat{R} = (\widehat{R} : (\widehat{R} : IJ\widehat{R})) = (\widehat{R} : (\widehat{R} : (IJ\widehat{R})_*)) = (\widehat{R} : (\widehat{R} : (I\widehat{R})_*)) = (\widehat{R} : (\widehat{R} : I\widehat{R})) = I\widehat{R}$. Since $I\widehat{R}$ is invertible, it follows that $J\widehat{R} = \widehat{R}$.

2.a. Let $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$ and let \overline{R} be a semilocal principal ideal domain. Then $\widehat{R} = \overline{R}$. By 1. it follows that $J\mathcal{R}(I)\widehat{R} = J\widehat{R} = \widehat{R}$, hence $J\mathcal{R}(I) \not\subseteq M$ for all $M \in \text{max}(\widehat{R})$. Since \widehat{R} is semilocal, this implies that $J\mathcal{R}(I) \not\subseteq \bigcup_{M \in \text{max}(\widehat{R})} M = \widehat{R} \setminus \widehat{R}^\times$, hence $J\mathcal{R}(I) \cap \widehat{R}^\times \neq \emptyset$. Of course, $\widehat{R}/\mathcal{R}(I)$ is an integral extension, hence $\widehat{R}^\times \cap \mathcal{R}(I) = \mathcal{R}(I)^\times$. This implies that $\emptyset \neq J\mathcal{R}(I) \cap \widehat{R}^\times = J\mathcal{R}(I) \cap \mathcal{R}(I) \cap \widehat{R}^\times = J\mathcal{R}(I) \cap \mathcal{R}(I)^\times$, hence $J\mathcal{R}(I) = \mathcal{R}(I)$.

2.b. Let R be a $*$ -noetherian G -domain. Without restriction we may assume that R is not a field. It follows that $(\mathcal{R}(I) : J\mathcal{R}(I)) = (I : IJ) = (I : (IJ)_*) = \mathcal{R}(I)$, hence $J\mathcal{R}(I) \subseteq (J\mathcal{R}(I))_{v_{\mathcal{R}(I)}} = \mathcal{R}(I)$. Since $\mathcal{R}(I) \in \mathcal{F}_*^\bullet(R)$ is an overring of R , it follows by Lemma 2.1.1. that $\mathcal{R}(I)$ is $*$ $[\mathcal{R}(I)]$ -noetherian, hence

$\mathcal{R}(I)$ is a Mori domain. If $\mathcal{R}(I) = K$, then $K \in \mathcal{F}_*(R)$, hence $R = K$, a contradiction. Consequently, Lemma 3.1.3. implies $\mathcal{R}(I) \setminus \mathcal{R}(I)^\times = \bigcup_{P \in v\text{-max}(\mathcal{R}(I))} P$, and for all $P \in v\text{-max}(\mathcal{R}(I))$, it follows that $P \neq \{0\}$. Therefore $P \cap R \neq \{0\}$ for all $P \in v\text{-max}(\mathcal{R}(I))$, hence $\{0\} \neq \bigcap_{M \in \text{spec}(R), M \neq \{0\}} M \subseteq R \cap \bigcap_{P \in v\text{-max}(\mathcal{R}(I))} P$. If $0 \neq x \in \bigcap_{P \in v\text{-max}(\mathcal{R}(I))} P$, then $\{P \in v\text{-spec}(\mathcal{R}(I)) \mid x \in P\} \supseteq v\text{-max}(\mathcal{R}(I))$, and thus $v\text{-max}(\mathcal{R}(I))$ is finite by Lemma 3.1.1.. Assume that $J\mathcal{R}(I) \subsetneq \mathcal{R}(I)$. Then there is some $Q \in \text{max}(\mathcal{R}(I))$ such that $J\mathcal{R}(I) \subseteq Q$. Since $Q \subseteq \mathcal{R}(I) \setminus \mathcal{R}(I)^\times = \bigcup_{P \in v\text{-max}(\mathcal{R}(I))} P$, it follows that there exists some $P \in v\text{-max}(\mathcal{R}(I))$ such that $Q \subseteq P$, hence $Q = P$ and $\mathcal{R}(I) = (J\mathcal{R}(I))_{v\mathcal{R}(I)} \subseteq Q_{v\mathcal{R}(I)} = Q$, a contradiction.

2.c. Let R be $*$ -noetherian and $\dim(R) = 1$. If $P \in \text{max}(R)$, then R_P is a $*$ -noetherian G -domain by Lemma 2.1.2.. Since $I_P, J_P \in \mathcal{F}_{*P}^\bullet(R_P)$, and $I_P = ((IJ)_*)_P = (I_P J_P)_{*P}$, we obtain $J_P \mathcal{R}(I_P) = \mathcal{R}(I_P)$ by 2.b.. Hence Lemma 2.1.3. implies that $J\mathcal{R}(I) = \bigcap_{P \in \text{max}(R)} (J\mathcal{R}(I))_P = \bigcap_{P \in \text{max}(R)} J_P \mathcal{R}(I_P) = \bigcap_{P \in \text{max}(R)} \mathcal{R}(I_P) = \bigcap_{P \in \text{max}(R)} \mathcal{R}(I)_P = \mathcal{R}(I)$.

3. Let R be $*$ -noetherian and $*$ -max(R) = $\text{spec}^1(R)$. If $P \in \text{spec}^1(R)$, then R_P is $*$ -noetherian and $I_P = ((IJ)_*)_P = (I_P J_P)_{*P}$ by Lemma 2.1.2.. Since $I_P, J_P \in \mathcal{F}_{*P}^\bullet(R_P)$ and $\dim(R_P) = 1$, we obtain $J_P \mathcal{R}(I_P) = \mathcal{R}(I_P)$ by 2.c.. Hence Lemma 2.1.3. implies that $((J\mathcal{R}(I))_*)_P = (J_P \mathcal{R}(I_P))_{*P} = \mathcal{R}(I_P)_{*P} = \mathcal{R}(I_P) = \mathcal{R}(I)_P$. Finally, we have $(J\mathcal{R}(I))_* = \bigcap_{P \in \text{spec}^1(R)} ((J\mathcal{R}(I))_*)_P = \bigcap_{P \in \text{spec}^1(R)} \mathcal{R}(I)_P = \mathcal{R}(I)$ by Lemma 2.1.3.. \square

Corollary 4.2. *Let R be an integral domain and $*$ a star operation on R .*

1. *If \widehat{R} is a Dedekind domain and $\mathcal{F}_{\widehat{R}/R} \in \mathbf{E}(\mathcal{F}_*(R))$, then $R = \widehat{R}$.*
2. *Suppose that one of the following conditions is satisfied:*
 - a. *$\mathcal{F}_{\widehat{R}/R} \neq \{0\}$ and \overline{R} is a semilocal principal ideal domain.*
 - b. *R is a $*$ -noetherian G -domain.*
 - c. *R is $*$ -noetherian and $*$ -max(R) = $\text{spec}^1(R)$.*

Then $\mathbf{E}(\mathcal{F}_(R)) = \{I \in \mathcal{F}_*(R) \mid I = \mathcal{R}(I)\}$.*

Proof. **1.** Let \widehat{R} be a Dedekind domain and $\mathcal{F}_{\widehat{R}/R} \in \mathbf{E}(\mathcal{F}_*(R))$. Then it follows by Theorem 4.1.1. that $\widehat{R} = \mathcal{F}_{\widehat{R}/R} \widehat{R} \subseteq R \subseteq \widehat{R}$, hence $R = \widehat{R}$.

2. This is an immediate consequence of Theorem 4.1.2. and Theorem 4.1.3.. \square

Theorem 4.3. *Let R be an integral domain, K a field of quotients of R and $*$ a star operation on R such that R is $*$ -noetherian. Suppose that one of the following conditions is fulfilled:*

- a. *\widehat{R} is a semilocal principal ideal domain.*
- b. *$*$ -max(R) = $\text{spec}^1(R)$ and \widehat{R}_P is a Krull domain for all $P \in \text{spec}^1(R)$.*

Then R is π -stable.*

Proof. **a.** Let \widehat{R} be a semilocal principal ideal domain and $I \in \mathcal{F}_*(R)$. Then $I\widehat{R} \in \mathcal{F}^\bullet(\widehat{R})$. Since \widehat{R} is a principal ideal domain, there exists some $a \in K^\bullet$ such that $I\widehat{R} = a\widehat{R}$, hence $a^{-1}I\widehat{R} = \widehat{R}$. Hence it follows that $a^{-1}I \not\subseteq M$ for all $M \in \text{max}(\widehat{R})$. Since \widehat{R} is semilocal, Lemma 3.2 implies that there exists some $u \in \mathbb{N}$ satisfying $a^{-u}I^u \not\subseteq \bigcup_{M \in \text{max}(\widehat{R})} M = \widehat{R} \setminus \widehat{R}^\times$. Consequently there exists some $b \in \widehat{R}^\times \cap a^{-u}I^u$, hence $R \subseteq b^{-1}a^{-u}I^u \subseteq \widehat{R}$. Let $J = b^{-1}a^{-u}(I^u)_*$. Then $J \in \mathcal{F}_*(R)$. Since R is $*$ -noetherian, there exists some finite subset $E \subseteq b^{-1}a^{-u}I^u$ satisfying $J = (E)_*$. Since E is finite and $E \subseteq \widehat{R}$, there exists some finitely generated R -submodule M of K such that $R[E] \subseteq M$, hence $R[E] \in \mathcal{F}^\bullet(R)$. Let $R' = (R[E])_*$. Then $R' \in \mathcal{F}_*(R)$ and $J \subseteq R'$. Since $R \subseteq J$, it follows for all $m \in \mathbb{N}$ that $(J^m)_* \subseteq (J^{m+1})_*$ and $(J^m)_* \subseteq ((R')^m)_* = (R[E]^m)_* = R'$. There exists some $x \in R^\bullet$ such that $xR' \subseteq R$, hence for all $m \in \mathbb{N}$ it follows that $x(J^m)_* \subseteq x(J^{m+1})_*$ and $x(J^m)_* \subseteq R$. Since R is $*$ -noetherian this implies that there exists some $r \in \mathbb{N}$ such that $x(J^r)_* = x(J^k)_*$ for all $k \in \mathbb{N}_{\geq r}$. This implies that $(J^r)_* = (J^{2r})_*$. Consequently,

$R \subseteq J \subseteq (J^r)_* \subseteq \mathcal{R}_*(J^r)$, hence $(J^r)_* = \mathcal{R}_*(J^r)$. This implies that $b^{-r}a^{-ru}(I^{ru})_* = \mathcal{R}_*(I^{ru})$. Let $n = ru \in \mathbb{N}$. Then $\mathcal{R}_*(I^n) = b^{-r}a^{-ru}(I^{ru})_* \subseteq (I^{ru}(\mathcal{R}_*(I^{ru}) : I^{ru}))_* \subseteq \mathcal{R}_*(I^{ru})$, hence $(I^n(\mathcal{R}_*(I^n) : I^n))_* = \mathcal{R}_*(I^n)$.

b. Let $*\text{-max}(R) = \text{spec}^1(R)$ and let \widehat{R}_P be a Krull domain for all $P \in \text{spec}^1(R)$. Let $I \in \mathcal{F}_*(R)$ and $\mathcal{P} = \{P \in \text{spec}^1(R) \mid I_P \neq R_P\}$. There is some $s \in R^\bullet$ such that $sI \subseteq R$ and some $s' \in I \setminus \{0\}$. Let $t = ss' \in R^\bullet$. Then $tR = ss'R \subseteq sIR = sI \subseteq I$. It follows that R is a Mori domain, hence Lemma 3.1.1. implies that $\{P \in \text{spec}^1(R) \mid st \in P\}$ is finite. Let us show that $\mathcal{P} \subseteq \{Q \in \text{spec}^1(R) \mid st \in Q\}$. Let $Q \in \mathcal{P}$ and assume that $st \in R \setminus Q$, then $st \in R_Q^\times$, hence $s, t \in R_Q^\times$. This implies that $I_Q = sI_Q = (sI)_Q \subseteq R_Q = tR_Q = (tR)_Q \subseteq I_Q$, hence $I_Q = R_Q$, a contradiction. It follows that \mathcal{P} is finite. If $P \in \text{spec}^1(R)$, then R_P is $*P$ -noetherian by Lemma 2.1.2., hence R_P is a Mori domain. Therefore \widehat{R}_P is a semilocal principal ideal domain by Lemma 3.1.5.. Since $I_P \in \mathcal{F}_{*P}^\bullet(R_P)$, it follows by a. and Lemma 2.2.5. that there exists a least $n_P \in \mathbb{N}$ such that $(I_P^k(\mathcal{R}_{*P}(I_P^k) : I_P^k))_{*P} = \mathcal{R}_{*P}(I_P^k)$ for all $k \in \mathbb{N}_{\geq n_P}$ (note that $n_P = 1$ if $P \notin \mathcal{P}$). If $n = \max(\{n_Q \mid Q \in \text{spec}^1(R)\} \cup \{1\})$, then it follows by Lemma 2.1.3. that $((I^n(\mathcal{R}_*(I^n) : I^n))_*)_Q = (I_Q^n(\mathcal{R}_{*Q}(I_Q^n) : I_Q^n))_{*Q} = \mathcal{R}_{*Q}(I_Q^n) = \mathcal{R}_*(I^n)_Q$ for all $Q \in \text{spec}^1(R)$. Consequently, we have $(I^n(\mathcal{R}_*(I^n) : I^n))_* = \bigcap_{Q \in \text{spec}^1(R)} ((I^n(\mathcal{R}_*(I^n) : I^n))_*)_Q = \bigcap_{Q \in \text{spec}^1(R)} \mathcal{R}_*(I^n)_Q = \mathcal{R}_*(I^n)$ by Lemma 2.1.3.. \square

Corollary 4.4. *Let R be an integral domain and $*$ a star operation on R . Assume that one of the following conditions is fulfilled:*

1. R is noetherian and $*\text{-max}(R) = \text{spec}^1(R)$.
2. R is $*\text{-noetherian}$, $*\text{-max}(R) = \text{spec}^1(R)$ and $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$.

Then R is $\pi*$ -stable.

Proof. **1.** Let R be noetherian, $*\text{-max}(R) = \text{spec}^1(R)$ and $P \in \text{spec}^1(R)$. Then R_P is noetherian with $\dim(R_P) = 1$. Therefore the theorem of Krull-Akizuki implies that \widehat{R}_P is a Dedekind domain, hence \widehat{R}_P is a Krull domain. Consequently, Theorem 4.3 implies that R is $\pi*$ -stable.

2. Let R be $*\text{-noetherian}$, $*\text{-max}(R) = \text{spec}^1(R)$, $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$ and $P \in \text{spec}^1(R)$. Then R_P is a Mori domain by Lemma 2.1.2.. Hence Lemma 3.1.4. implies that $\mathcal{F}_{\widehat{R}_P/R_P} = (\mathcal{F}_{\widehat{R}/R})_P \supseteq \mathcal{F}_{\widehat{R}/R} \neq \{0\}$, consequently \widehat{R}_P is a Krull domain by Lemma 3.1.4.. Therefore R is $\pi*$ -stable by Theorem 4.3. \square

Obviously, the conditions of Corollary 4.4. are satisfied for arbitrary noetherian 1-dimensional integral domains. We were not be able to decide whether $*\text{-noetherian}$ integral domains satisfying $*\text{-max}(R) = \text{spec}^1(R)$ are $\pi*$ -stable. By Theorem 2.3.3. and Corollary 4.2. it is sufficient to show that the $*\text{-ideal}$ semigroup of such integral domains is π -regular.

5. EXAMPLES

The examples and counterexamples of this section are based on subrings of the ring of formal power series $R[[X]]$ over an integral domain R . For a power series f we denote by $(f_i)_{i \in \mathbb{N}_0}$ the sequence of its coefficients, so that $f = \sum_{i \in \mathbb{N}_0} f_i X^i$. For $d \in R^\bullet$ we define $R_d = \{f \in R[[X]] \mid d \mid f_1\}$ and if $b, c \in R$ are such that $b \mid d$ and $c \mid d$, then we set $I_{b,c} = \{f \in R[[X]] \mid b \mid f_0, c \mid f_1\}$. Observe that R is completely integrally closed if and only if $R[[X]]$ is completely integrally closed (for a proof see [8, Theorem 16]). Recall that if $R[[X]]$ is a Mori domain, that R is a Mori domain (since $R[[X]] \cap K = R$ (where K is a field of quotients of R), this is a consequence of [1, Theorem 2.4]).

First we study some ring theoretical properties of R_d and investigate the elements of its v -ideal semigroup. Especially we investigate the completely integrally closed case, where the conductor turns out to be an idempotent of the v -ideal semigroup. In Lemma 5.3. and Lemma 5.4. we study most of the properties introduced in Section 2 with respect to the v -ideal semigroup of R_d . Moreover we explicitly calculate

the set of v -idempotents in some special cases. The most important counterexamples are consolidated in Example 5.5..

Lemma 5.1. *Let R be an integral domain, $d \in R^\bullet$ and K' a field of quotients of $R[[X]]$.*

1. R_d is an intermediate ring of R and $R[[X]]$, and K' is a quotient field of R_d .
2. For all $b, c \in R$ such that $b|d$ and $c|d$ it follows that $I_{b,c} = (b, cX, X^2, X^3)_{R_d} \in \mathcal{F}_v^\bullet(R_d)$, $(I_{b,c})^{-1} = I_{\frac{d}{c}, \frac{d}{b}}$, $\mathcal{R}(I_{b,c}) = \{f \in R[[X]] \mid c|bf_1\}$ and $(I_{b,c}^k)^{-1} = \{f \in R[[X]] \mid d|b^k f_1, d|b^{k-1} c f_0\}$ for all $k \in \mathbb{N}$.

Proof. 1. Of course, $R \subseteq R_d \subseteq R[[X]]$. Therefore it is sufficient to show that if $f, g \in R_d$, then $f+g, fg \in R_d$. Let $f, g \in R_d$. Then $f+g, fg \in R[[X]]$, $d|(f_1+g_1) = (f+g)_1$ and $d|(f_0g_1+f_1g_0) = (fg)_1$, hence $f+g, fg \in R_d$. Let $K'' \subseteq K'$ be the quotient field of R_d . Since $X = X^3X^{-2} \in K''$, it follows that $R[[X]] \subseteq K''$, hence $K'' = K'$.

2. Let $b, c \in R$ be such that $b|d$ and $c|d$. At first, let us show that $I_{b,c} = (b, cX, X^2, X^3)_{R_d}$. “ \subseteq ”: Let $f \in R[[X]]$ be such that $b|f_0$ and $c|f_1$. Then there are $v, w \in R$ such that $f_0 = vb$ and $f_1 = wc$, hence $f = vb + wcX + (\sum_{i \in \mathbb{N}_{\geq 2}, i \neq 3} f_i X^{i-2})X^2 + f_3 X^3 \in (b, cX, X^2, X^3)_{R_d}$. “ \supseteq ”: Since $b, cX, X^2, X^3 \in I_{b,c}$, it is sufficient to show that $I_{b,c}$ is an R_d -submodule of $R[[X]]$. Therefore let $g, h \in I_{b,c}$ and let $p, q \in R_d$. Then $gp + hq \in R[[X]]$, $b|(g_0p_0 + h_0q_0) = (gp + hq)_0$. It follows that $c|d|p_1$ and $c|d|q_1$. Consequently, $c|(g_1p_0 + g_0p_1 + h_1q_0 + h_0q_1) = (gp + hq)_1$, hence $gp + hq \in I_{b,c}$. Since $X^2 \in R_d^\bullet$ and $X^2(b, cX, X^2, X^3)_{R_d} = (bX^2, cX^3, X^4, X^5)_{R_d} \subseteq R_d$, it follows that $(b, cX, X^2, X^3)_{R_d}^k \in \mathcal{F}^\bullet(R_d)$.

Let $k \in \mathbb{N}$. If $g \in K'$ is such that $g(b, cX, X^2, X^3)_{R_d}^k \subseteq R_d$, then $gb^k \in R_d \subseteq R[[X]]$ and $gX^{2k} \in R_d \subseteq R[[X]]$. Hence $g \in R[[X]]$. This implies that $((b, cX, X^2, X^3)_{R_d}^k)^{-1} = \{f \in R[[X]] \mid f(b, cX, X^2, X^3)_{R_d}^k \subseteq R_d\}$. If $f \in R[[X]]$, then $X^2f \in R_d$. Hence $((b, cX, X^2, X^3)_{R_d}^k)^{-1} = \{f \in R[[X]] \mid fb^k, fb^{k-1}cX \in R_d\} = \{f \in R[[X]] \mid d|b^k f_1, d|b^{k-1} c f_0\}$.

Since $\frac{d}{b}|d$, $\frac{d}{c}|d$ and $b, c \in R$ are arbitrary it follows that $(b, cX, X^2, X^3)_{R_d}^{-1} = \{f \in R[[X]] \mid \frac{d}{c}|f_0, \frac{d}{b}|f_1\} = I_{\frac{d}{c}, \frac{d}{b}}$ and $(I_{b,c})_v = (I_{\frac{d}{c}, \frac{d}{b}})^{-1} = I_{b,c}$. Finally, it follows that $\mathcal{R}(I_{b,c}) = \{f \in R[[X]] \mid f(b, cX, X^2, X^3)_{R_d} \subseteq I_{b,c}\} = \{f \in R[[X]] \mid fb, fcX, fX^2, fX^3 \in I_{b,c}\} = \{f \in R[[X]] \mid c|bf_1\}$. \square

Lemma 5.2. *Let R be a completely integrally closed domain, $d \in R^\bullet$ and K' a field of quotients of $R[[X]]$.*

1. $\overline{R_d} = \widehat{R_d} = R[[X]] = (1, X)_{R_d}$.
2. $\mathcal{F}_{\widehat{R_d}/R_d} = I_{d,d} = (d, dX, X^2, X^3)_{R_d} \in \mathcal{E}(\mathcal{F}_v^\bullet(R_d))$.
3. Let $Q \in \text{spec}(R_d)$ be such that $\mathcal{F}_{\widehat{R_d}/R_d} \subseteq Q$. Then $ht(Q) \geq 2$.
4. $X \in \bigcap_{P \in \text{spec}^1(R_d)} (R_d)_P$.
5. Let R_d be a Mori domain and $P \in \text{spec}^1(R_d)$. Then $(R_d)_P$ is a discrete valuation domain.

Proof. 1. Let us show that $R[[X]] = (1, X)_{R_d}$. “ \subseteq ”: Let $f \in R[[X]]$. Then $f = f_1X + (\sum_{i \in \mathbb{N}_0, i \neq 1} f_i X^i)1 \in (1, X)_{R_d}$. “ \supseteq ”: Clear. Since $R[[X]]$ is completely integrally closed, this implies that $R[[X]] \subseteq \overline{R_d} \subseteq \widehat{R_d} \subseteq \widehat{R[[X]]} = R[[X]]$, hence $\overline{R_d} = \widehat{R_d} = R[[X]] = (1, X)_{R_d}$.

2. It follows by Lemma 5.1.2. that $I_{d,d} = (d, dX, X^2, X^3)_{R_d} \in \mathcal{F}_v^\bullet(R_d)$. Let us show that $\mathcal{F}_{\widehat{R_d}/R_d} = I_{d,d}$. “ \subseteq ”: Let $f \in \mathcal{F}_{\widehat{R_d}/R_d}$. Then $f \in R_d$, hence $d|f_1$. Since $fX \in R_d$, we have $d|(fX)_1 = f_0$. “ \supseteq ”: Since $\widehat{R_d} = R[[X]]$, it follows by 1. that $\{d, dX, X^2, X^3\} \subseteq \mathcal{F}_{\widehat{R_d}/R_d}$, hence $(d, dX, X^2, X^3)_{R_d} \subseteq \mathcal{F}_{\widehat{R_d}/R_d}$. Therefore Lemma 5.1.2. implies that $(I_{d,d}^2)^{-1} = \{f \in R[[X]] \mid d|d^2 f_0, d|d^2 f_1\} = R[[X]]$, hence $((\mathcal{F}_{\widehat{R_d}/R_d})^2)_v = (I_{d,d}^2)_v = (R[[X]])^{-1} = \mathcal{F}_{\widehat{R_d}/R_d}$.

3. Let $h : R[[X]] \rightarrow R$ be the canonical ringepimorphism and $g = h|_{R_d}$. Then g is a ringepimorphism. Let us show that $\text{Ker}(g) = (dX, X^2, X^3)_{R_d}$. “ \subseteq ”: Let $f \in \text{Ker}(g)$. Then $f_0 = 0$. Of course, $d|f_1$, hence $f = f_1X + X^2(\sum_{i \in \mathbb{N}_{\geq 2}, i \neq 3} f_i X^{i-2}) + f_3 X^3 \in (dX, X^2, X^3)_{R_d}$. “ \supseteq ”: Trivial. This implies that $R_d/(dX, X^2, X^3)_{R_d} \cong R$, consequently $(dX, X^2, X^3)_{R_d} \in \text{spec}(R_d)$. Since $\{0\} \subsetneq (dX, X^2, X^3)_{R_d} \subsetneq \mathcal{F}_{\widehat{R_d}/R_d} \subseteq Q$, we have $ht(Q) \geq 2$.

4. Let $P \in \text{spec}^1(R_d)$. Then it follows by 2. and 3. that $\{d, dX, X^2, X^3\} \not\subseteq P$.

Case1: $d \notin P$: Since $d, dX \in R_d$, we have $X = d^{-1}dX \in (R_d)_P$.

Case2: $dX \notin P$: Since $dX, dX^2 \in R_d$, it follows that $X = (dX)^{-1}dX^2 \in (R_d)_P$.

Case3: $X^2 \notin P$: Since $X^2, X^3 \in R_d$, we have $X = X^{-2}X^3 \in (R_d)_P$.

Case4: $X^3 \notin P$: Since $X^3, X^4 \in R_d$, it follows that $X = X^{-3}X^4 \in (R_d)_P$.

5. Since $X \in (R_d)_P$ (by 4.), it follows by Lemma 3.1.2. and 1. that $\widehat{(R_d)_P} = \widehat{(R_d)}_P = R[[X]]_P \subseteq ((R_d)_P)_P = (R_d)_P$, hence $(R_d)_P$ is completely integrally closed. Since $(R_d)_P$ is a Mori domain, this implies that $(R_d)_P$ is a Krull domain. Since $(R_d)_P$ is a Krull domain and $\dim((R_d)_P) = 1$, it follows that $(R_d)_P$ is a Dedekind domain and since $(R_d)_P$ is local this implies that $(R_d)_P$ is a discrete valuation domain. \square

Lemma 5.3. *Let R be a completely integrally closed domain, $d \in R^\bullet$ and K' a field of quotients of $R[[X]]$.*

1. $\{I_{b,c} \mid b, c \in R, b|d, c|d\} \subseteq \{I \in \mathcal{F}_v^\bullet(R_d) \mid \mathcal{F}_{\widehat{R_d}/R_d} \subseteq I \text{ and } I^2 \subseteq I\}$, and if R is a GCD-domain, then $\{I_{b,c} \mid b, c \in R, b|c|d, GCD(\frac{d}{b}, b) = R^\times\} \subseteq \mathbf{E}(\mathcal{F}_v^\bullet(R_d))$.
2. Let $\{I \in \mathcal{F}_v^\bullet(R_d) \mid \mathcal{F}_{\widehat{R_d}/R_d} \subseteq I \text{ and } I^2 \subseteq I\} \subseteq \{I_{b,c} \mid b, c \in R, b|d, c|d\}$. Then for all $a \in R$ such that $a^2|d$ it follows that $a \in R^\times$.
3. If R/dR is finite, then $\{I \mid I \text{ is an } R_d\text{-submodule of } \widehat{R_d} \text{ such that } \mathcal{F}_{\widehat{R_d}/R_d} \subseteq I\}$ is finite and $\mathcal{F}_v^\bullet(R_d)$ is almost complete.
4. If R/dR is finite, then for all $I \in \mathcal{F}_v^\bullet(R_d)$ such that $\mathcal{F}_{\widehat{R_d}/R_d} \subseteq I \subseteq \widehat{R_d}$ it follows that I is πv -regular.
5. Let $I \in \mathcal{F}_v^\bullet(R_d)$ be such that $\mathcal{F}_{\widehat{R_d}/R_d} \subseteq I \subseteq \widehat{R_d}$. Then there exists some $P \subseteq R \times R$ such that $I = \{f \in R[[X]] \mid \text{for all } (b, c) \in P \text{ it follows that } d|(bf_1 + cf_0)\}$.

Proof. 1. Let $b, c \in R$ be such that $b|d$ and $c|d$ and $I = I_{b,c}$. Then it follows by Lemma 5.1.2. that $I \in \mathcal{F}_v^\bullet(R_d)$. Since $b|d$ and $c|d$, it follows by Lemma 5.1.2. that $\mathcal{F}_{\widehat{R_d}/R_d} = \{f \in R[[X]] \mid d|f_0, d|f_1\} \subseteq \{f \in R[[X]] \mid b|f_0, c|f_1\} = I$. Since $I^2 = (b^2, bcX, bX^2, bX^3, c^2X^2, cX^3, cX^4, X^4, X^5, X^6)_{R_d}$ and all generators of I^2 are products of elements of I and R_d , it follows that $I^2 \subseteq I$.

Now let R be a GCD-domain, $b|c$ and $GCD(\frac{d}{b}, b) = R^\times$. It follows by Lemma 5.1.2. that $(I^2)^{-1} = \{f \in R[[X]] \mid \frac{d}{b}|bf_1, \frac{d}{c}|cf_0\}$. Since $b|c$, it follows that $\frac{d}{c}|\frac{d}{b}$, hence $GCD(\frac{d}{c}, b) = R^\times$. Since R is a GCD-domain, this implies that $(I^2)^{-1} = \{f \in R[[X]] \mid \frac{d}{b}|f_1, \frac{d}{c}|f_0\} = (\frac{d}{c}, \frac{d}{b}X, X^2, X^3)_{R_d} = I^{-1}$. Hence $I = (I^2)_v$, and thus $I \in \mathbf{E}(\mathcal{F}_v^\bullet(R_d))$.

2. Let $a \in R$ be such that $a^2|d$ and $I = (a^2, a^2X, aX + a, X^2, X^3)_{R_d}$. Of course, $I \in \mathcal{F}^\bullet(R_d)$. By Lemma 5.1.2. $I^{-1} = (I_{a^2, a^2} + (aX + a)R_d)^{-1} = (I_{a^2, a^2})^{-1} \cap (aX + a)^{-1}R_d = I_{\frac{d}{a^2}, \frac{d}{a^2}} \cap \{f \in K' \mid (aX + a)f \in R_d\} = \{f \in R[[X]] \mid \frac{d}{a^2}|f_0, \frac{d}{a^2}|f_1, \frac{d}{a}|(f_0 + f_1)\}$.

Let us show that $I^{-1} = (\frac{d}{a}, \frac{d}{a}X, \frac{d}{a^2}(1 - X), X^2, X^3)_{R_d}$. “ \subseteq ”: Let $f \in R[[X]]$ be such that $\frac{d}{a^2}|f_0, \frac{d}{a^2}|f_1$ and $\frac{d}{a}|(f_0 + f_1)$. Then there exist some $r, s \in R$ such that $f_0 + f_1 = \frac{d}{a}r$ and $f_0 = \frac{d}{a^2}s$, hence $f = f_0 + f_1X + \sum_{i \in \mathbb{N}_{\geq 2}} f_iX^i = s\frac{d}{a^2}(1 - X) + r\frac{d}{a}X + \sum_{i \in \mathbb{N}_{\geq 2}} f_iX^i \in (\frac{d}{a}, \frac{d}{a}X, \frac{d}{a^2}(1 - X), X^2, X^3)_{R_d}$. “ \supseteq ”: Trivial, since I^{-1} is an R_d -module.

Consequently, it follows by Lemma 5.1.2. that $I_v = (I^{-1})^{-1} = (I_{\frac{d}{a}, \frac{d}{a}} + \frac{d}{a^2}(1 - X)R_d)^{-1} = (I_{\frac{d}{a}, \frac{d}{a}})^{-1} \cap (\frac{d}{a^2}(1 - X))^{-1}R_d = I_{a, a} \cap \{f \in K' \mid \frac{d}{a^2}(1 - X)f \in R_d\} = \{f \in R[[X]] \mid a|f_0, a|f_1, a^2|(f_1 - f_0)\}$.

Let us show that $I_v = I$. “ \subseteq ”: Let $f \in R[[X]]$ be such that $a|f_0, a|f_1$ and $a^2|(f_1 - f_0)$. Then there exist some $r, s \in R$ such that $f_1 = f_0 + a^2r$ and $f_0 = as$. This implies that $f = f_0 + f_1X + \sum_{i \in \mathbb{N}_{\geq 2}} f_iX^i = s(aX + a) + ra^2X + \sum_{i \in \mathbb{N}_{\geq 2}} f_iX^i \in I$. “ \supseteq ”: Trivial.

Of course, $I \in \mathcal{F}_v^\bullet(R)$. Since $a^2|d$, it follows by Lemma 5.1.2. and Lemma 5.2.2. that $\mathcal{F}_{\widehat{R_d}/R_d} = I_{d, d} \subseteq I_{a^2, a^2} = (a^2, a^2X, X^2, X^3)_{R_d} \subseteq I$. Of course, $I^2 = (I_{a^2, a^2})^2 + (a^3X + a^3, a^4X + a^4, a^2X^2 + 2a^2X + a^2, aX^3 + aX^2, aX^4 + aX^3)_{R_d} \subseteq (a^2, a^2X, X^2, X^3)_{R_d} \subseteq I$. Therefore there are some $b_0, c_0 \in R$ such

that $b_0|d$, $c_0|d$ and $I = I_{b_0, c_0}$. It follows by Lemma 5.1.2. that $aX + a \in I_{b_0, c_0}$, hence $c_0|a$. Since $c_0X \in I = \{f \in R[[X]] \mid a|f_0, a|f_1, a^2|(f_1 - f_0)\}$, it follows that $a^2|c_0$. This implies that $a^2|a$, hence $a \in R^\times$.

3. Let R/dR be finite and $M = \{I \mid I \text{ is an } R_d\text{-submodule of } \widehat{R}_d \text{ such that } \mathcal{F}_{\widehat{R}_d/R_d} \subseteq I\}$. Let $f : R/dR \times R/dR \rightarrow \widehat{R}_d/\mathcal{F}_{\widehat{R}_d/R_d}$ be defined by $f((a + dR, b + dR)) = (a + bX) + \mathcal{F}_{\widehat{R}_d/R_d}$ for all $a, b \in R$. Let $a, a_1, b, b_1 \in R$ be such that $a + dR = a_1 + dR$ and $b + dR = b_1 + dR$. Then there exist some $c_1, d_1 \in R$ such that $a = a_1 + c_1d$ and $b = b_1 + d_1d$. Therefore Lemma 5.2.2. implies that $(a + bX) + \mathcal{F}_{\widehat{R}_d/R_d} = (a_1 + b_1X) + (c_1d + d_1dX) + \mathcal{F}_{\widehat{R}_d/R_d} = (a_1 + b_1X) + \mathcal{F}_{\widehat{R}_d/R_d}$, hence f is well-defined. Now let $g \in \widehat{R}_d = R[[X]]$. Then by Lemma 5.2.2. $\sum_{i \in \mathbb{N}_{\geq 2}} g_i X^i \in \mathcal{F}_{\widehat{R}_d/R_d}$, hence $f((g_0 + dR, g_1 + dR)) = (g_0 + g_1X) + \mathcal{F}_{\widehat{R}_d/R_d} = g + \mathcal{F}_{\widehat{R}_d/R_d}$, and thus f is surjective.

Since $R/dR \times R/dR$ is finite, this implies that $\widehat{R}_d/\mathcal{F}_{\widehat{R}_d/R_d}$ is finite. Since there exists a bijection between the set of all R_d -submodules of $\widehat{R}_d/\mathcal{F}_{\widehat{R}_d/R_d}$ and M , it follows that M is finite. It follows by Lemma 2.4.1. that $E(\mathcal{F}_v^\bullet(R_d)) \subseteq M$, hence $E(\mathcal{F}_v^\bullet(R_d))$ is finite. This implies that $\mathcal{F}_v^\bullet(R_d)$ is almost complete.

4. Let R/dR be finite and $I \in \mathcal{F}_v^\bullet(R_d)$ such that $\mathcal{F}_{\widehat{R}_d/R_d} \subseteq I \subseteq \widehat{R}_d$. It follows by Lemma 5.2.2. that $\mathcal{F}_{\widehat{R}_d/R_d} = ((\mathcal{F}_{\widehat{R}_d/R_d})^k)_v \subseteq (I^k)_v \subseteq ((\widehat{R}_d)^k)_v = \widehat{R}_d$ for all $k \in \mathbb{N}$. Since $\{I \in \mathcal{F}_v^\bullet(R_d) \mid \mathcal{F}_{\widehat{R}_d/R_d} \subseteq I \subseteq \widehat{R}_d\}$ is finite, there are some $r, s \in \mathbb{N}$ such that $r \neq s$ and $(I^r)_v = (I^s)_v$. Hence I is πv -regular by Lemma 1.1.2..

5. Let $P = \{(b, c) \in R \times R \mid b + cX \in I^{-1}\}$. Let us show that $I = \{f \in R[[X]] \mid \text{for all } (b, c) \in P \text{ we have } d|(bf_1 + cf_0)\}$. “ \subseteq ”: Let $f \in I$ and $(b, c) \in P$. Since $b + cX \in I^{-1}$, it follows that $(b + cX)f \in R_d$ and hence $d|(b + cX)f_1 = bf_1 + cf_0$. “ \supseteq ”: Let $f \in R[[X]]$ be such that $d|(bf_1 + cf_0)$ for all $(b, c) \in P$. It is sufficient to show that $fg \in R_d$ for all $g \in I^{-1}$. Let $g \in I^{-1}$. Then $g \in I^{-1} \subseteq (R_d :_{K'} \mathcal{F}_{\widehat{R}_d/R_d}) = \widehat{R}_d = R[[X]]$. It follows by Lemma 5.2.2. that $\sum_{i \in \mathbb{N}_{\geq 2}} g_i X^i \in (X^2, X^3)_{R_d} \subseteq \mathcal{F}_{\widehat{R}_d/R_d} \subseteq I^{-1}$, hence $g_0 + g_1X \in I^{-1}$. Therefore $(g_0, g_1) \in P$, and thus $d|(g_0f_1 + g_1f_0) = (gf)_1$. This implies that $gf \in R_d$. \square

Lemma 5.4. *Let R be a completely integrally closed domain, $d \in R^\bullet$ and K' a field of quotients of $R[[X]]$.*

1. *The following assertions are equivalent:*

- a.** $d \in R^\times$.
- b.** R_d is completely integrally closed.
- c.** $\mathcal{F}_v^\bullet(R_d)$ is a group.
- d.** $\mathcal{F}_v^\bullet(R_d)$ is a Clifford semigroup.
- e.** $E(\mathcal{F}_v^\bullet(R_d)) = \{F \in \mathcal{F}_v^\bullet(R_d) \mid F = \mathcal{R}(F)\}$.
- f.** $\mathcal{F}_{\widehat{R}_d/R_d} \in \{F \in \mathcal{F}_v^\bullet(R_d) \mid F = \mathcal{R}(F)\}$.

2. *If d is a product of pairwise non-associated prime elements of R , then $\{I \in \mathcal{F}_v^\bullet(R_d) \mid \mathcal{F}_{\widehat{R}_d/R_d} \subseteq I \text{ and } I^2 \subseteq I\} = \{I_{b,c} \mid b, c \in R, b|d, c|d\}$.*

3. *If d is a product of prime elements of R , then $E(\mathcal{F}_v^\bullet(R_d)) = \{I_{b,c} \mid b, c \in R, b|c|d, \text{GCD}(b, \frac{d}{b}) = R^\times\}$ and $\mathcal{F}_v^\bullet(R_d)$ is almost complete.*

4. *If $R[[X]]$ is factorial and if d is a prime element of R or R/dR is finite, then $\mathcal{F}_v^\bullet(R_d)$ is complete.*

Proof. **1. a. \Rightarrow b.** Let $d \in R^\times$. Then $R_d = \{f \in R[[X]] \mid d|f_1\} = R[[X]] = \widehat{R}_d$ by Lemma 5.2.1., hence R_d is completely integrally closed.

b. \Rightarrow c. Clear. **c. \Rightarrow d.** Trivial.

d. \Rightarrow a. Let $\mathcal{F}_v^\bullet(R_d)$ be a Clifford semigroup and $I = \mathcal{F}_{\widehat{R}_d/R_d} + XR_d$. It follows by Lemma 5.2.2. that $I = (d, X, X^2, X^3)_{R_d}$. We show that $(I^2)_v = (I\mathcal{F}_{\widehat{R}_d/R_d})_v = \mathcal{F}_{\widehat{R}_d/R_d}$. It follows by Lemma 5.1.2. and 5.2.2. that $\mathcal{F}_{\widehat{R}_d/R_d} = ((\mathcal{F}_{\widehat{R}_d/R_d})^2)_v \subseteq (I\mathcal{F}_{\widehat{R}_d/R_d})_v \subseteq (I^2)_v = (\{f \in R[[X]] \mid d|d^2f_1, d|df_0\})^{-1} = (R[[X]])^{-1} = \mathcal{F}_{\widehat{R}_d/R_d}$, hence $(I^2)_v = (I\mathcal{F}_{\widehat{R}_d/R_d})_v = \mathcal{F}_{\widehat{R}_d/R_d}$. Since I is v -regular, there exist some $E \in E(I)$ and

$J \in \mathcal{F}_v^\bullet(R_d)$ such that $(IJ)_v = E$, hence $(\mathcal{F}_{\widehat{R}_d/R_d} J^2)_v = (I^2 J^2)_v = (E^2)_v = E$. This implies that $(\mathcal{F}_{\widehat{R}_d/R_d} E)_v = ((\mathcal{F}_{\widehat{R}_d/R_d})^2 J^2)_v = (\mathcal{F}_{\widehat{R}_d/R_d} J^2)_v = E$. It follows by Lemma 2.4.1. that $\mathcal{F}_{\widehat{R}_d/R_d} \subseteq E = (E\mathcal{F}_{\widehat{R}_d/R_d})_v \subseteq (\widehat{R}_d \mathcal{F}_{\widehat{R}_d/R_d})_v = \mathcal{F}_{\widehat{R}_d/R_d}$, hence $E = \mathcal{F}_{\widehat{R}_d/R_d}$. This implies that $\mathcal{F}_{\widehat{R}_d/R_d} \in \mathbf{E}(I)$, hence $\mathcal{F}_{\widehat{R}_d/R_d} = (I\mathcal{F}_{\widehat{R}_d/R_d})_v = I$. Finally, it follows by Lemma 5.2.2. that $X \in I = \mathcal{F}_{\widehat{R}_d/R_d} = I_{d,d}$. Therefore $d|1$ and thus $d \in R^\times$.

c. \Rightarrow **e.** Clear. **e.** \Rightarrow **f.** This is an immediate consequence of Lemma 5.2.2..

f. \Rightarrow **a.** Let $\mathcal{F}_{\widehat{R}_d/R_d} \in \{F \in \mathcal{F}_v^\bullet(R_d) \mid F = \mathcal{R}(F)\}$. Then by Lemma 5.2.2., $1 \in \mathcal{F}_{\widehat{R}_d/R_d} = I_{d,d}$. It follows that $d|1$, hence $d \in R^\times$.

2. Let $n \in \mathbb{N}$ and let $(p_i)_{i=1}^n$ be a finite sequence of pairwise non-associated prime elements of R such that $d = \prod_{i=1}^n p_i$. Now we show the equality. “ \supseteq ”: This follows by Lemma 5.3.1.. “ \subseteq ”: Let $I \in \mathcal{F}_v^\bullet(R_d)$ be such that $\mathcal{F}_{\widehat{R}_d/R_d} \subseteq I$ and $I^2 \subseteq I$. Then $I \subseteq \mathcal{R}(I) \subseteq \widehat{R}_d = R[[X]]$. By Lemma 5.3.5. there exists some $P \subseteq R \times R$ such that $I = \{f \in R[[X]] \mid d|(b_0 f_1 + c_0 f_0) \text{ for all } (b_0, c_0) \in P\}$. Let $M = \{i \in [1, n] \mid \text{there is some } (b_0, c_0) \in P \text{ such that } p_i \nmid c_0\}$, $N = \{i \in [1, n] \mid \text{there is some } (b_0, c_0) \in P \text{ such that } p_i \nmid b_0\}$, $b = \prod_{i \in M} p_i$ and $c = \prod_{i \in N} p_i$. Next we show that $I = I_{b,c}$.

“ \subseteq ”: Let $f \in I$. Then $f^2 \in I^2 \subseteq I$. If $(b_1, c_1) \in P$, then $d|(b_1(f^2)_1 + c_1(f^2)_0)$. Therefore $d|(b_1(2f_0 f_1 + c_1 f_0^2))$, and thus $d|(b_1 f_0 f_1 + f_0(b_1 f_1 + c_1 f_0))$. Since $d|(b_1 f_1 + c_1 f_0)$, it follows that $d|b_1 f_0 f_1$. Now we prove that $b|f_0$. It is sufficient to show that $p_i|f_0$ for all $i \in M$. Let $i \in M$. Then there exists some $(b_0, c_0) \in P$ such that $p_i \nmid c_0$.

Case1: $p_i \nmid b_0$: Since $p_i|d|b_0 f_0 f_1$, we have $p_i|f_0$ or $p_i|f_1$. If $p_i|f_1$, then since $p_i|d|(b_0 f_1 + c_0 f_0)$, we have $p_i|f_0$.

Case2: $p_i|b_0$: Since $p_i|d|(b_0 f_1 + c_0 f_0)$, we have $p_i|f_0$.

Now we prove that $c|f_1$. It is sufficient to show that for all $i \in N$ it follows that $p_i|f_1$. Let $i \in N$. Then there exists some $(b_0, c_0) \in P$ such that $p_i \nmid b_0$. Since $p_i|d|b_0 f_0 f_1$, we have $p_i|f_0$ or $p_i|f_1$. If $p_i|f_0$, then $p_i|f_1$, since $p_i|d|(b_0 f_1 + c_0 f_0)$.

“ \supseteq ”: Since $X^2, X^3 \in \mathcal{F}_{\widehat{R}_d/R_d} \subseteq I$, we have to show that $b, cX \in I$. Let us show that $b \in I$. It is sufficient to show that for all $i \in [1, n]$ and all $(b_1, c_1) \in P$ it follows that $p_i|c_1 b$. Let $i \in [1, n]$ and $(b_1, c_1) \in P$.

Case1: For all $(b_0, c_0) \in P$ it follows that $p_i|c_0$: Of course, $p_i|c_1|c_1 b$.

Case2: There exists some $(b_0, c_0) \in P$ such that $p_i \nmid c_0$: Since $i \in M$, we have $p_i|b|c_1 b$.

Let us show that $cX \in I$. It is sufficient to show that for all $i \in [1, n]$ and all $(b_1, c_1) \in P$ it follows that $p_i|b_1 c$. Let $i \in [1, n]$ and $(b_1, c_1) \in P$.

Case1: For all $(b_0, c_0) \in P$ it follows that $p_i|b_0$: Of course, $p_i|b_1|b_1 c$.

Case2: There is some $(b_0, c_0) \in P$ such that $p_i \nmid b_0$: Since $i \in N$, it follows that $p_i|c|b_1 c$.

3. Without restriction let $n \in \mathbb{N}$, $(p_i)_{i=1}^n$ a finite sequence of pairwise non-associated primes of R and $(k_i)_{i=1}^n \in \mathbb{N}^{[1, n]}$ be such that $d = \prod_{i=1}^n p_i^{k_i}$. Claim: $(I_{p_i, 1}^{k_i+1})_v = I_{p_i^{k_i}, p_i^{k_i}}$ for all $i \in [1, n]$. Proof of the claim:

Let $i \in [1, n]$. Then it follows by Lemma 5.1.2. that $(I_{p_i, 1}^{k_i+1})^{-1} = \{f \in R[[X]] \mid d|p_i^{k_i+1} f_1, d|p_i^{k_i} f_0\} = \{f \in R[[X]] \mid \frac{d}{p_i^{k_i}}|f_1, \frac{d}{p_i^{k_i}}|f_0\} = I_{\frac{d}{p_i^{k_i}}, \frac{d}{p_i^{k_i}}}$, hence $(I_{p_i, 1}^{k_i+1})_v = (I_{\frac{d}{p_i^{k_i}}, \frac{d}{p_i^{k_i}}})^{-1} = I_{p_i^{k_i}, p_i^{k_i}}$ by Lemma 5.1.2..

Next we show the equality. “ \subseteq ”: Let $I \in \mathbf{E}(\mathcal{F}_v^\bullet(R_d))$. Then we have $\mathcal{F}_{\widehat{R}_d/R_d} \subseteq I \subseteq \widehat{R}_d$ by Lemma 2.4.1., hence there exists some $P \subseteq R \times R$ such that $I = \{f \in R[[X]] \mid d|(b f_1 + c f_0) \text{ for all } (b, c) \in P\}$ by Lemma 5.3.5.. Let $M = \{i \in [1, n] \mid p_i|f_0 \text{ for all } f \in I\}$, $l_i = \max\{r \in [0, k_i] \mid p_i^r|f_1 \text{ for all } f \in I\}$ for all $i \in [1, n]$, $b = \prod_{j \in M} p_j^{k_j}$ and $c = \prod_{j=1}^n p_j^{l_j}$. Next we show that $b|c$. Let $i \in M$. Then $I \subseteq I_{p_i, 1}$ by Lemma 5.1.2., and hence $I = (I^{k_i+1})_v \subseteq (I_{p_i, 1}^{k_i+1})_v = I_{p_i^{k_i}, p_i^{k_i}}$ by the claim. It follows that $p_i^{k_i}|f_1$ for all $f \in I$. Therefore $l_i = k_i$, hence $b|c$. Of course, $c|d$ and since $(p_i)_{i=1}^n$ is a sequence of pairwise non-associated prime elements of R , we have $\text{GCD}(b, \frac{d}{b}) = R^\times$.

It remains to prove that $I = I_{b,c}$. “ \subseteq ”: Let $f \in I$. If $j \in M$, then $I \subseteq I_{p_j^{k_j}, p_j^{k_j}}$. Therefore $p_j^{k_j}|f_0$, hence $b|f_0$. Of course, $p_j^{l_j}|f_1$ for all $j \in [1, n]$, hence $c|f_1$. “ \supseteq ”: It follows by Lemma 5.2.2. and Lemma 2.4.1.

that $X^2, X^3 \in \mathcal{F}_{\widehat{R}_d/R_d} \subseteq I$. Due to Lemma 5.1.2., it is sufficient to show that $b \in I$ and $cX \in I$. We have to prove that for all $j \in [1, n]$ and all $(b_1, c_1) \in P$ it follows that $p_j^{k_j} | c_1 b$ and $p_j^{k_j} | b_1 c$. Let $j \in [1, n]$ and $(b_1, c_1) \in P$.

Case1: $j \in M$: Since $p_j^{k_j} | b | c$, we have $p_j^{k_j} | c_1 b$ and $p_j^{k_j} | c b_1$.

Case2: $j \notin M$: There is some $f \in I$ such that $p_j \nmid f_0$. Since $f^2 \in I^2 \subseteq I$, we have $d | (b_1(f^2)_1 + c_1(f^2)_0)$, hence $d | (b_1 f_0 f_1 + f_0(b_1 f_1 + c_1 f_0))$. Since $f \in I$, it follows that $p_j^{k_j} | d | b_1 f_0 f_1$ and since $p_j \nmid f_0$, we have $p_j^{k_j} | b_1 f_1$. Since $p_j^{k_j} | d | (b_1 f_1 + c_1 f_0)$, it follows that $p_j^{k_j} | c_1 | c_1 \prod_{i \in M} p_i^{k_i}$. If $l_j = k_j$, then $p_j^{k_j} | b_1 c$, hence we may assume that $l_j < k_j$. There exists some $g \in I$ such that $p_j^{l_j} | g_1$ and $p_j^{l_j+1} \nmid g_1$. It follows that $p_j^{k_j} | d | (b_1 g_1 + c_1 g_0)$, hence $p_j^{k_j} | b_1 g_1$ and hence $p_j^{k_j-l_j} | b_1$. This implies that $p_j^{k_j} = p_j^{k_j-l_j} p_j^{l_j} | b_1 c$.

“ \supseteq “: Let $b, c \in R$ be such that $b | c | d$ and $GCD(b, \frac{d}{b}) = R^\times$. We set $I = (b, cX, X^2, X^3)_{R_d}$. It follows by Lemma 5.1.2. that $I \in \mathcal{F}_v^\bullet(R_d)$ and $(I^2)^{-1} = \{f \in R[[X]] \mid \frac{d}{b} | b f_1, \frac{d}{c} | b f_0\}$.

Let us show that $(I^2)^{-1} = I_{\frac{d}{c}, \frac{d}{b}}$. “ \subseteq “: Let $f \in (I^2)^{-1}$, then $\frac{d}{c} | b f_0$ and $\frac{d}{b} | b f_1$. Since $\frac{d}{c} | \frac{d}{b}$, we have $GCD(b, \frac{d}{c}) = R^\times$. Of course, $\frac{d}{b}$ and $\frac{d}{c}$ are associate to products of prime elements of R . Therefore it is straightforward to show that $\frac{d}{c} | f_0$ and $\frac{d}{b} | f_1$. Consequently, $f \in I_{\frac{d}{c}, \frac{d}{b}}$. “ \supseteq “: Trivial. Therefore Lemma 5.1.2. implies that $(I^2)^{-1} = I^{-1}$, hence $(I^2)_v = I$. Now it is straightforward to prove that $E(\mathcal{F}_v^\bullet(R_d))$ is finite. Hence it follows that $\mathcal{F}_v^\bullet(R_d)$ is almost complete.

4. Let $R[[X]]$ be factorial. At first let d be a prime element of R . It follows by Lemma 5.2.2., 2. and 3. that $\{J \in \mathcal{F}_v^\bullet(R_d) \mid \mathcal{F}_{\widehat{R}_d/R_d} \subseteq J \text{ and } J^2 \subseteq J\} = \{\mathcal{F}_{\widehat{R}_d/R_d}, \mathcal{F}_{\widehat{R}_d/R_d} + XR_d, R_d, \widehat{R}_d\}$, $E(\mathcal{F}_v^\bullet(R_d)) = \{\mathcal{F}_{\widehat{R}_d/R_d}, R_d, \widehat{R}_d\}$ and $\mathcal{F}_v^\bullet(R_d)$ is almost complete. We have to show that $\mathcal{F}_v^\bullet(R_d)$ is πv -regular. Let $I \in \mathcal{F}_v^\bullet(R_d)$. Then $\mathcal{R}(I) \in \{R_d, \widehat{R}_d\}$.

Case1: $\mathcal{R}(I) = R_d$: Since $\mathcal{F}_{\mathcal{R}(I)/R_d} = R_d \in E(I)$, we have I is v -regular by Lemma 2.4.2..

Case2: $\mathcal{R}(I) = \widehat{R}_d$: Since $\widehat{R}_d \in E(I)$ and $\widehat{R}_d \notin E(R_d)$, it follows by Lemma 2.2.4. and Lemma 2.4.1. that $(I(\widehat{R}_d :_{K'} I))_v \in \{J \in \mathcal{F}_v^\bullet(R_d) \mid \mathcal{F}_{\widehat{R}_d/R_d} \subseteq J \subseteq \widehat{R}_d, \widehat{R}_d \in E(J) \text{ and } J^2 \subseteq J\} = \{\mathcal{F}_{\widehat{R}_d/R_d}, \mathcal{F}_{\widehat{R}_d/R_d} + XR_d, \widehat{R}_d\}$. Since $\widehat{R}_d = R[[X]]$ is factorial and $(\widehat{R}_d :_{K'} I) \in \mathcal{F}_{v_{\widehat{R}_d}}^\bullet(\widehat{R}_d)$, it follows that there is some $c \in K'^\bullet$ such that $(\widehat{R}_d :_{K'} I) = c \widehat{R}_d$. Hence $(I(\widehat{R}_d :_{K'} I))_v = (c I \widehat{R}_d)_v = c I$, and thus $I \in \{c^{-1} \mathcal{F}_{\widehat{R}_d/R_d}, c^{-1}(\mathcal{F}_{\widehat{R}_d/R_d} + XR_d), c^{-1} \widehat{R}_d\}$.

Case2.1: $I = c^{-1} \mathcal{F}_{\widehat{R}_d/R_d}$: Since $\mathcal{F}_{\mathcal{R}(I)/R_d} \in E(I)$, we have I is v -regular by Lemma 2.4.2..

Case2.2: $I = c^{-1}(\mathcal{F}_{\widehat{R}_d/R_d} + XR_d)$: Since $(I^2)_v = c^{-2} \mathcal{F}_{\widehat{R}_d/R_d}$ and $\mathcal{F}_{\mathcal{R}_v(I^2)/R_d} \in E((I^2)_v)$, it follows by Lemma 2.4.2. that $(I^2)_v$ is v -regular, hence I is πv -regular.

Case2.3: $I = c^{-1} \widehat{R}_d$: Since $c \widehat{R}_d \in \mathcal{F}_v^\bullet(R_d)$ and $(I c \widehat{R}_d)_v = \widehat{R}_d$, we have I is v -regular.

Now let R/dR be finite. It follows by Lemma 5.3.3. that $\mathcal{F}_v^\bullet(R_d)$ is almost complete. Let $J \in \mathcal{F}_v^\bullet(R_d)$. Then since $\widehat{R}_d = R[[X]]$ is factorial, it follows by Lemma 2.4.3. and Lemma 5.2.2. that there exists some $c \in K'^\bullet$ such that $\mathcal{F}_{\widehat{R}_d/R_d} \subseteq cJ \subseteq \widehat{R}_d$. Therefore Lemma 5.3.4. implies that cJ is πv -regular, hence J is πv -regular. \square

Example 5.5. *Each of the following properties is satisfied by some integral domain R :*

1. R is noetherian, $\dim(R) = \dim(\overline{R}) = 2$, \overline{R} is local, noetherian and factorial, $\mathcal{F}_{\widehat{R}/R} \in E(\mathcal{F}_v^\bullet(R)) \setminus \{I \in \mathcal{F}_v^\bullet(R) \mid I = \mathcal{R}(I)\}$ and R_P is a discrete valuation domain for all $P \in \text{spec}^1(R)$.
2. R is neither a Mori domain nor completely integrally closed and yet $\mathcal{F}_v^\bullet(R)$ is almost complete.
3. $\mathcal{F}_{\widehat{R}/R} \neq \{0\}$, $\dim(\widehat{R}) = 2$, \widehat{R} is local, noetherian and factorial and there exists some $I \in \mathcal{F}_v^\bullet(R)$ such that $\mathcal{R}_v(I^n) \subsetneq \mathcal{R}_v(I^{n+1})$ for all $n \in \mathbb{N}$. Especially $\mathcal{F}_v^\bullet(R)$ is not π -regular.

Proof. 1. Let S be a discrete valuation domain, L a field of quotients of S , X an indeterminate over L , $d \in S^\bullet \setminus S^\times$, and $R = \{f \in S[[X]] \mid d | f_1\}$. It follows by Lemma 5.2.1. that $\overline{R} = \widehat{R} = S[[X]] = (1, X)_R$,

hence \bar{R} is a local, noetherian and factorial domain and $\dim(R) = \dim(\bar{R}) = 2$. It follows by the theorem of Eakin-Nagata that R is noetherian, hence Lemma 5.2.5. implies that R_P is a discrete valuation domain for all $P \in \text{spec}^1(R)$. Since $d \notin S^\times$ it follows by Lemma 5.2.2. and Lemma 5.4.1. that $\mathcal{F}_{\hat{R}/R} \in \mathbb{E}(\mathcal{F}_v^\bullet(R)) \setminus \{I \in \mathcal{F}_v^\bullet(R) \mid I = \mathcal{R}(I)\}$.

2. Let S be a completely integrally closed domain, that is not a Mori domain, for example the ring of algebraic integers. Let Y be an indeterminate over S . It follows that $S[[Y]]$ is a completely integrally closed domain that is not a Mori domain. Of course, Y is a prime element of $S[[Y]]$. Let L be a field of quotients of $S[[Y]]$, X an indeterminate over L and $R = \{f \in (S[[Y]])[[X]] \mid Y \mid_{S[[Y]]} f_1\}$. Then Lemma 5.4.3. implies that $\mathcal{F}_v^\bullet(R)$ is almost complete. Since $X \notin R$ it follows that R is not completely integrally closed. Assume that R is a Mori domain. Then Lemma 3.1.4. and Lemma 5.2.1. imply that $\hat{R} = (S[[Y]])[[X]]$ is a Mori domain, hence $S[[Y]]$ is a Mori domain, a contradiction.

3. Let S be an integral domain that is not a field. Let K' be a quotient field of S , Y an indeterminate over K' and $T = S + YK'[[Y]]$. It follows that T and $K'[[Y]]$ have the same field of quotients, we denote it by L . Since $K'[[Y]]$ is a principal ideal domain, it follows that $\hat{T} \subseteq \widehat{K'[[Y]]} = K'[[Y]]$. On the other hand $YK'[[Y]] \subseteq T$. Since $Y \in T^\bullet$, we have $K'[[Y]] \subseteq \hat{T}$, hence $\hat{T} = K'[[Y]]$. It follows that \hat{T} is a discrete valuation domain. There exists some $b \in S^\bullet \setminus S^\times$. Of course, $b \in T \setminus T^\times$ and $\bigcap_{n \in \mathbb{N}} b^n T \supseteq YK'[[Y]] \neq \{0\}$. Let $d \in T^\bullet$ be such that $d \in \bigcap_{n \in \mathbb{N}} b^n T$.

Let X be an indeterminate over L , $R = \{f \in T[[X]] \mid d \mid f_1\}$ and $I = (b, dX, X^2, X^3)_R$. It follows that $\hat{T}[[X]]$ is local, noetherian, factorial and $\dim(\hat{T}[[X]]) = 2$. This implies that $\hat{R} \subseteq \widehat{\hat{T}[[X]]} = \hat{T}[[X]]$. Since $X^2 Y \hat{T}[[X]] \subseteq X^2 T[[X]] \subseteq R$ and $X^2 Y \in R^\bullet$, we have $\hat{T}[[X]] \subseteq \hat{R}$. Therefore $\hat{R} = \hat{T}[[X]]$ and $\mathcal{F}_{\hat{R}/R} \neq \{0\}$. By Lemma 5.1.2. it follows that $(I^n)^{-1} = \{f \in T[[X]] \mid d \mid b^n f_1, d \mid b^{n-1} d f_0\} = (1, \frac{d}{b^n} X, X^2, X^3)_R$ for all $n \in \mathbb{N}$. Hence it follows by Lemma 5.1.2. that $\mathcal{R}_v(I^n) = ((I^n)_v : I^n) = (R : (I^n)^{-1} I^n) = \mathcal{R}((I^n)^{-1}) = \mathcal{R}((1, \frac{d}{b^n} X, X^2, X^3)_R) = \{f \in T[[X]] \mid \frac{d}{b^n} \mid f_1\} = (1, \frac{d}{b^n} X, X^2, X^3)_R$ for all $n \in \mathbb{N}$. Assume that there exists some $m \in \mathbb{N}$ such that $\mathcal{R}_v(I^m) = \mathcal{R}_v(I^{m+1})$, then $\frac{d}{b^{m+1}} X \in \mathcal{R}_v(I^{m+1}) = \mathcal{R}_v(I^m) = \{f \in T[[X]] \mid \frac{d}{b^m} \mid f_1\}$, hence $\frac{d}{b^m} \mid \frac{d}{b^{m+1}}$. This implies that $b \in T^\times$, a contradiction. Hence Lemma 1.1.3. and Lemma 2.2.4. imply that $\mathcal{F}_v^\bullet(R)$ is not π -regular. \square

Note that Example 5.5.1. shows that the idempotents of the v -ideal semigroup of a Mori domain need not be trivial. Moreover by Example 5.5.2. it follows that the converse of Theorem 2.6.1. does not hold. By Example 5.5.3. we get that the π -regularity of the v -ideal semigroup does not descend from the complete integral closure.

Acknowledgement. This work was supported by the Austrian Science Fund FWF (Project Number P21576-N18) and is part of my doctoral thesis. I want to thank my advisor Professor Franz Halter-Koch for the many helpful suggestions. I am indebted to the referee for pointing out a few inaccuracies in a previous version of the paper.

REFERENCES

- [1] V. Barucci, *Mori domains*, Non-noetherian commutative ring theory, Math. Appl., vol. 520, Kluwer Academic Publishers, 2000, pp. 57-73.
- [2] S. Bazzoni, *Class semigroups of Prüfer domains*, J. Algebra **184** (1996), 613-631.
- [3] S. Bazzoni, *Idempotents of the class semigroup of a Prüfer domain of finite character*, Abelian groups, modules theory and topology, Lect. Notes Pure Appl. Math. 201, Marcel Dekker, 1998, pp. 79-89.
- [4] S. Bazzoni, *Groups in the class semigroup of a Prüfer domain of finite character*, Comm. Algebra **28** (2000), 5157-5167.
- [5] S. Bazzoni, *Clifford regular domains*, J. Algebra **238** (2007), 703-722.
- [6] S. Bazzoni, S. Kabbaj, *Class semigroups and t -class semigroups of integral domains*, Commutative Algebra, Noetherian and Non-Noetherian Perspectives, 2010, pp. 47-76.
- [7] S. Bazzoni, L. Salce, *Groups in the class semigroup of valuation domains*, Israel J. Math. **95** (1996), 135-155.
- [8] J.W. Brewer, *Power series over commutative rings*, Marcel Dekker, 1981.

- [9] E.C. Dade, O. Taussky, H. Zassenhaus, *On the theory of orders, in particular on the semigroup of ideal classes and genera on an order in an algebraic number field*, Math. Ann. **148** (1962), 31-64.
- [10] L. Fuchs, L. Salce, *Modules over non-Noetherian domains*, Mathematical Surveys and Monographs, American Mathematical Society, 2001.
- [11] S. Gabelli, G. Picozza, *Star stable domains*, J. Pure Appl. Algebra **208** (2007), 853-866.
- [12] A. Geroldinger, F. Halter-Koch, *Non-unique factorizations: Algebraic, combinatorial and analytic theory*, Monogr. Textb. Pure Appl. Math., Chapman and Hall/CRC, 2006.
- [13] R.W. Gilmer, *Multiplicative ideal theory*, Marcel Dekker, 1972.
- [14] P.A. Grillet, *Semigroups: An introduction to the structure theory*, Marcel Dekker, 1995.
- [15] F. Halter-Koch, *Ideal systems*, Monogr. Textb. Pure Appl. Math., Marcel Dekker, 1998.
- [16] F. Halter-Koch, *Localizing systems, module systems and semistar operations*, J. Algebra **238** (2001), 723-761.
- [17] F. Halter-Koch, *Ideal semigroups of noetherian domains and Ponizovksi decompositions*, J. Pure Appl. Algebra **209** (2007), 763-770.
- [18] F. Halter-Koch, *Clifford semigroups of ideals in monoids and domains*, Forum Math. **21** (2009), 1001-1020.
- [19] F. Halter-Koch, *Multiplicative ideal theory in the context of commutative monoids*, Commutative Algebra, Noetherian and Non-Noetherian Perspectives, 2010, pp. 203-231.
- [20] S. Kabbaj, A. Mimouni, *Class semigroups of integral domains*, J. Algebra **264** (2003), 620-640.
- [21] S. Kabbaj, A. Mimouni, *t-class semigroups of integral domains*, J. Reine Angew. Math. **612** (2007), 213-229.
- [22] S. Kabbaj, A. Mimouni, *Constituent groups of Clifford semigroups arising from t-closure*, J. Algebra **321** (2009), 1443-1452.
- [23] P. Zanardo, U. Zannier, *The class semigroup of orders in number fields*, Math. Proc. Cambridge Phil. Soc. **115** (1994), 379-391.

INSTITUT FÜR MATHEMATIK UND WISSENSCHAFTLICHES RECHNEN, KARL-FRANZENS-UNIVERSITÄT, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA

E-mail address: andreas.reinhart@uni-graz.at