

# NUMERICAL ANALYSIS OF A NONSMOOTH QUASILINEAR ELLIPTIC CONTROL PROBLEM: II. FINITE ELEMENT DISCRETIZATION AND ERROR ESTIMATES

Christian Clason\*

Vu Huu Nhu<sup>†</sup>

Arnd Rösch<sup>‡</sup>

**Abstract** In this paper, we carry out the numerical analysis of a nonsmooth quasilinear elliptic optimal control problem, where the coefficient in the divergence term of the corresponding state equation is not differentiable with respect to the state variable. Despite the lack of differentiability of the nonlinearity in the quasilinear elliptic equation, the corresponding control-to-state operator is of class  $C^1$  but not of class  $C^2$ . Analogously, the discrete control-to-state operators associated with the approximated control problems are proven to be of class  $C^1$  only. By using an explicit second-order sufficient optimality condition, we prove a priori error estimates for a variational approximation, a piecewise constant approximation, and a continuous piecewise linear approximation of the continuous optimal control problem. The numerical tests confirm these error estimates.

**Key words** Optimal control, nonsmooth optimization, quasilinear elliptic equation, piecewise differentiable function, sufficient optimality condition, error estimate, finite element approximation

## 1 INTRODUCTION

We investigate the nonsmooth quasilinear elliptic optimal control problem

$$(P) \quad \begin{cases} \min_{u \in L^\infty(\Omega)} j(u) := \int_{\Omega} L(x, y_u(x)) \, dx + \frac{\nu}{2} \|u\|_{L^2(\Omega)}^2 \\ \text{s.t.} & -\operatorname{div}[(b + a(y_u)) \nabla y_u] = u \quad \text{in } \Omega, \quad y_u = 0 \text{ on } \partial\Omega, \\ & \alpha \leq u(x) \leq \beta \quad \text{a.e. } x \in \Omega. \end{cases}$$

Here  $\Omega$  is a two-dimensional bounded, convex and polygonal domain;  $L : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function that is of class  $C^2$  with respect to (w.r.t.) the second variable;  $b : \bar{\Omega} \rightarrow \mathbb{R}$  is a Lipschitz continuous function;  $a : \mathbb{R} \rightarrow \mathbb{R}$  is Lipschitz continuous but not differentiable; and constants  $\alpha, \beta, \nu \in \mathbb{R}$  satisfying  $\beta > \alpha$  and  $\nu > 0$ . We refer to Section 2 for the precise assumptions on the data of (P).

The state equation in the optimal control problem (P) occurs, for instance, in models of heat conduction in which the coefficient in the divergence term acts as the heat conductivity and is a function of two variables: the temperature  $y$  variable and the spatial coordinate  $x$  variable; see, e.g. [5, 45]. When the data belong to class  $C^2$ , the numerical analysis of the discrete approximation of optimal control problems governed by such state equations were studied by Casas et al. in [16, 17] for distributed control and in [10] for Neumann control.

---

\*Faculty of Mathematics, Universität Duisburg-Essen, 45117 Essen, Germany; *current address*: Institute of Mathematics and Scientific Computing, University of Graz, Heinrichstrasse 36, 8010 Graz, Austria (c.clason@uni-graz.at, ORCID: 0000-0002-9948-8426)

<sup>†</sup>Faculty of Fundamental Sciences, PHENIKAA University, Yen Nghia, Ha Dong, Hanoi 12116, Vietnam (nhu.vuhuu@phenikaa-uni.edu.vn, ORCID: 0000-0003-4279-3937)

<sup>‡</sup>Faculty of Mathematics, University of Duisburg-Essen, Thea-Leymann-Strasse 9, 45127 Essen, Germany (arnd.roesch@uni-due.de)

Let us briefly comment on other works concerning the error analysis of optimal control problems governed by partial differential equations (PDEs), in particular by elliptic PDEs. For control-constrained elliptic problems, we refer to the early papers [31, 32] for linear elliptic control problems; to [2, 14] for semilinear elliptic problems. For state-constrained control problems, we mention only the recent contributions [28, 42] and refer to the survey paper [36] for further references. Although the error analysis for smooth PDE-constrained problems has been intensively investigated, there are very few contributions on this topic for nonsmooth PDE-constrained optimal control. Here we want to mention the work [37] concerning error analysis for optimal control of a coupled PDE–ODE system, where the nonsmooth nonlinearity acts on a semilinear ODE. For works related to optimal control of obstacle problems, we refer to [41, 19] and the references therein. Based on a quadratic growth condition, a priori error estimates were established in [37, 41]. To the authors' best knowledge, this is the first work that exploits a second-order sufficient optimality condition to show a priori error estimates for the discretization of optimal control problems governed by nonsmooth PDEs.

In this paper, our main aim is to derive the convergence analysis and error estimates of the discretization of (P) under the explicit second-order sufficient optimality conditions established in [24] (first derived in general form in [22]). As the coefficient  $a$  in the state equation is Lipschitz continuous but not differentiable, there are two major difficulties in deriving the error analysis. The first issue arises in studying error estimates of the discretization of the adjoint state equation; the other is the lack of the second-order differentiability of the cost functional. We therefore cannot apply an abstract theorem on error estimates shown in [17] and in [13]. To deal with the first issue, we introduce the function  $Z_{y,\hat{y}}$  defined in (3.6) that measures pointwise the difference between the gradients of the superposition mappings of  $a$  associated with two distinct states  $y$  and  $\hat{y}$ . This allows us to derive thereafter both  $L^2$ - and an  $H_0^1$ - error estimates for the approximation of the adjoint state equation. Let us emphasize that the solutions to the linearized state equation can only be shown to be in a  $W^{1,p}$ -space (see Remark 2.4) and thus we cannot directly employ the standard duality argument based on the Aubin–Nitsche trick; see, e.g. [11, 26, 7]. In order to derive  $L^2$ -error estimates, we will instead introduce an *adjusted* linearized state equation in (4.19) below. Under the assumption that the jump functional  $\Sigma$  defined in (2.13) to be finite at the optimal state  $\bar{y}$ , we show that the solutions to this adjusted equation belong to a fractional Sobolev space of order greater than 1. We can then apply standard interpolation error estimates to obtain the desired result. For handling the second issue, we will exploit an structural assumption on the optimal state and employ an explicit formula of a second-order generalized derivative of the objective functional, formulated in [24]. Based on the second-order sufficient optimality conditions for (P) from [22, 24], we then prove general error estimates for variational, piecewise constant, and continuous piecewise linear approximations of the optimal control, which generalize those of Theorem 2.14 in [17] and of Lemma 5.2 in [13], and can be applied for the case where the cost functional  $j$  is of class  $C^1$  but not necessarily  $C^2$ ; see Theorem 5.10.

The plan of the paper is as follows. This section ends with our notation. In the next section, we make the assumptions for (P) and provide some preliminary results from [22] and [24]. Section 3 is devoted to the numerical approximation of the state equation by finite elements and the local well-posedness and differentiability of the discrete counterpart of the control-to-state operator. In Section 4, the error analysis of the adjoint state equation is investigated. Finally, the main results of the paper are presented in Section 5. There, the convergence and error estimates of local minima of discrete optimal control problems are, respectively, stated in Section 5.1 and Section 5.2. The numerical tests illustrating the obtained results are given in Section 5.3. Finally, the verification of a structural assumption and a computation of the jump functional for a specific situation are carried out in Appendix A, while the regularity of solutions in a fractional Sobolev space to the adjusted linearized state equation is shown in Appendix B.

**Notation.** We denote by  $B_X(u, \rho)$  and  $\overline{B}_X(u, \rho)$  the open and closed balls in a Banach space  $X$  of radius  $\rho > 0$  centered at  $u \in X$ , respectively. For Banach spaces  $X$  and  $Y$ , the notation  $X \hookrightarrow (\Subset) Y$  is understood that  $X$  is continuously (compactly) embedded in  $Y$ . Let  $X$  be a Banach space with its dual  $X^*$ , the symbol  $\langle \cdot, \cdot \rangle_{X^*, X}$  stands for the dual product of  $X$  and  $X^*$ . For a given function  $g : \overline{\Omega} \rightarrow \mathbb{R}$  and a subset  $A \subset \mathbb{R}$ ,  $\{g \in A\}$  denotes the set of all points  $x \in \overline{\Omega}$  for which  $g(x) \in A$ . For functions  $g_1, g_2$  and subsets  $A_1, A_2 \subset \mathbb{R}$ , we set  $\{g_1 \in A_1, g_2 \in A_2\} := \{g_1 \in A_1\} \cap \{g_2 \in A_2\}$ . For any set  $\omega \subset \overline{\Omega}$ , we denote by  $\mathbb{1}_\omega$  the characteristic function of  $\omega$ , i.e.,  $\mathbb{1}_\omega(x) = 1$  if  $x \in \omega$  and  $\mathbb{1}_\omega(x) = 0$  otherwise. We write the symbol  $C$  for a generic positive constant, which may be different at different places of occurrence and the notation, e.g.  $C_\xi$  for a constant depending only on the parameter  $\xi$ . For a measurable two-dimensional subset  $M$ , by  $\text{meas}_{\mathbb{R}^2}(M)$ , we denote the two-dimensional Lebesgue measure of  $M$ . Finally, the symbol  $\mathcal{H}^1$  denotes the one-dimensional Hausdorff measure on  $\mathbb{R}^2$  that is scaled as in [30], Def. 2.1.

## 2 MAIN ASSUMPTIONS AND PRELIMINARY RESULTS

In this section we first present assumptions, which will be used in the whole paper, and then state some preliminary results on the state equation, the adjoint state equation, as well as the first-order and explicit second-order sufficient optimality conditions from [22] and [24].

We first address the salient point, which is the structure of the nondifferentiable nonlinearity  $a$ . In this work, we assume that  $a$  is defined by

$$(2.1) \quad a(t) := \mathbb{1}_{(-\infty, \bar{t}]}(t)a_0(t) + \mathbb{1}_{(\bar{t}, \infty)}(t)a_1(t) \quad \text{for all } t \in \mathbb{R},$$

for a given number  $\bar{t} \in \mathbb{R}$  and given functions  $a_0 \in C^2((-\infty, \bar{t}])$  and  $a_1 \in C^2([\bar{t}, \infty))$  with  $a_0(\bar{t}) = a_1(\bar{t})$ . Obviously,  $a$  is Lipschitz continuous on  $\mathbb{R}$  and twice continuously differentiable on  $(-\infty, \bar{t}) \cup (\bar{t}, \infty)$ , but not even of class  $C^1$  in general. However,  $a$  is directionally differentiable and its directional derivative for all  $t \in \mathbb{R}$  in direction  $s \in \mathbb{R}$  is given by

$$(2.2) \quad a'(t; s) = \mathbb{1}_{(-\infty, \bar{t})}(t)a'_0(t)s + \mathbb{1}_{(\bar{t}, \infty)}(t)a'_1(t)s + \mathbb{1}_{\{\bar{t}\}}(t)[\mathbb{1}_{(0, \infty)}(s)a'_1(\bar{t})s + \mathbb{1}_{(-\infty, 0)}(s)a'_0(\bar{t})s].$$

**Remark 2.1.** Let us emphasize that the results and the underlying analysis in this paper can be applied to the situation in which the function  $a$  is continuous and is twice continuously differentiable on finitely many intervals (i.e., a finitely  $PC^2$  function; see [22] for a precise definition). However, in order to keep the presentation concise and to be able to focus on the main arguments, we restrict the presentation to the simplest such situation given by (2.1).

The following assumptions shall hold throughout the following.

- (A1)  $\Omega \subset \mathbb{R}^2$  is an open bounded convex polygonal.
- (A2) The Lipschitz continuous function  $b : \overline{\Omega} \rightarrow \mathbb{R}$  satisfies  $b(x) \geq \underline{b} > 0$  for all  $x \in \overline{\Omega}$ .
- (A3)  $a : \mathbb{R} \rightarrow \mathbb{R}$  is nonnegative and given by (2.1).
- (A4)  $L : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function that is of class  $C^2$  w.r.t. the second variable with  $L(\cdot, 0) \in L^1(\Omega)$ . Besides, for any  $M > 0$ , there exist  $C_M > 0$  and  $\psi_M \in L^{\bar{p}}(\Omega)$  ( $\bar{p} > 2$ ) such that  $|\frac{\partial L}{\partial y}(x, y)| \leq \psi_M(x)$  and  $|\frac{\partial^2 L}{\partial y^2}(x, y)| \leq C_M$  for all  $y \in \mathbb{R}$  with  $|y| \leq M$ , and a.e.  $x \in \Omega$ .

In the remainder of this subsection, we state some known results for the state equation, the adjoint state equation, and the optimality conditions for (P); see, e.g. [22] and [24]. Let us first study the state equation

$$(2.3) \quad -\text{div}[(b + a(y))\nabla y] = u \text{ in } \Omega, \quad y = 0 \text{ on } \partial\Omega.$$

**Theorem 2.2** ([24], Thm. 3.2 and cf. [22], Thms. 3.1 and 3.5). *Assume that Assumptions (A1) to (A3) are verified. Then, the control-to-state mapping  $S : W^{-1,p}(\Omega) \ni u \mapsto y_u \in W_0^{1,p}(\Omega)$  is of class  $C^1$ , where  $y_u$  is the unique solution to (2.3). Moreover, for any  $u, v \in W^{-1,p}(\Omega)$  with  $p > 2$  and  $y_u := S(u)$ ,  $z_v := S'(u)v$  uniquely satisfies*

$$(2.4) \quad -\operatorname{div}[(b + a(y_u))\nabla z_v + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) z_v \nabla y_u] = v \text{ in } \Omega, \quad z_v = 0 \text{ on } \partial\Omega.$$

Moreover, a number  $p_* > 2$  exists and satisfies that for any  $p \in [2, p_*)$  and for any bounded set  $U \subset L^p(\Omega)$ , there hold  $S(u) \in W_0^{1,p}(\Omega) \cap W^{2,p}(\Omega)$  and  $\|S(u)\|_{W^{2,p}(\Omega)} \leq C_U$ .

We now investigate the adjoint state equation

$$(2.5) \quad -\operatorname{div}[(b + a(y_u))\nabla \varphi] + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \nabla y_u \cdot \nabla \varphi = v \text{ in } \Omega, \quad \varphi = 0 \text{ on } \partial\Omega$$

for  $u \in W^{-1,p}(\Omega)$ ,  $p > 2$ ,  $v \in H^{-1}(\Omega)$ , and  $y_u := S(u)$ .

**Theorem 2.3** ([24], Thm. 3.3 and cf. [22], Lem. 4.1). *Assume that Assumptions (A1) to (A3) are satisfied. Let  $p, q > 2$  be arbitrary. Then, for any  $u \in W^{-1,p}(\Omega)$ ,  $v \in H^{-1}(\Omega)$ , there exists a unique  $\varphi \in H_0^1(\Omega)$  which solves (2.5). Moreover, if  $U$  is a bounded subset in  $L^p(\Omega)$ , then for any  $u \in U$  and any  $v \in L^q(\Omega)$ , the solution  $\varphi$  to (2.5) is an element of  $H^2(\Omega) \cap W^{1,\infty}(\Omega)$  and there holds  $\|\varphi\|_{H^2(\Omega)} + \|\varphi\|_{W^{1,\infty}(\Omega)} \leq C_U \|v\|_{L^q(\Omega)}$ . Furthermore, if  $u \in L^p(\Omega)$  and  $v \in L^r(\Omega)$  with  $r \in (2, p_*)$ , then  $\varphi \in W^{2,r}(\Omega)$ , where  $p_*$  is given as in Theorem 2.2.*

**Remark 2.4.** In spite of the  $W^{2,p}$ -regularity of the state and adjoint state, the function  $z_v := S'(u)v$  determined in (2.4) belongs to  $W^{1,p}(\Omega)$  only. This fact is due to the nondifferentiability of the function  $a$ .

The optimal control problem (P) can be transferred in the following form

$$(P) \quad \min_{u \in \mathcal{U}_{ad}} j(u) = \int_{\Omega} L(x, S(u)(x)) \, dx + \frac{\nu}{2} \|u\|_{L^2(\Omega)}^2,$$

where the admissible set is defined as

$$\mathcal{U}_{ad} := \{u \in L^\infty(\Omega) \mid \alpha \leq u(x) \leq \beta \quad \text{for a.e. } x \in \Omega\}.$$

Thanks to Assumptions (A1) to (A4), the cost functional  $j : L^2(\Omega) \rightarrow \mathbb{R}$  is first-order continuously differentiable and satisfies

$$(2.6) \quad j'(u)v = \int_{\Omega} (\varphi_u + \nu u)v \, dx \quad \text{for } u, v \in L^2(\Omega)$$

with  $\varphi_u \in H_0^1(\Omega)$  solving (2.5) corresponding to the right-hand side term  $v$  substituted by  $\frac{\partial L}{\partial y}(\cdot, S(u))$ ; see [22], Thm. 4.2. We have the following first-order necessary optimality conditions from Theorem 4.3 in [22].

**Theorem 2.5** ([22], Thm. 4.3). *Let Assumptions (A1) to (A4) hold. Then (P) admits at least one local minimizer  $\bar{u}$ . Furthermore, an adjoint state  $\bar{\varphi} \in H_0^1(\Omega)$  exists and fulfills the following first-order optimality conditions*

$$(2.7a) \quad -\operatorname{div}[(b + a(\bar{y}))\nabla \bar{y}] = \bar{u} \quad \text{in } \Omega, \quad \bar{y} = 0 \text{ on } \partial\Omega,$$

$$(2.7b) \quad -\operatorname{div}[(b + a(\bar{y}))\nabla \bar{\varphi}] + \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'(\bar{y}) \nabla \bar{y} \cdot \nabla \bar{\varphi} = \frac{\partial L}{\partial y}(x, \bar{y}) \text{ in } \Omega, \quad \bar{\varphi} = 0 \text{ on } \partial\Omega,$$

$$(2.7c) \quad \int_{\Omega} (\bar{\varphi} + \nu \bar{u})(u - \bar{u}) \, dx \geq 0 \quad \text{for all } u \in \mathcal{U}_{ad},$$

with  $\bar{y} := S(\bar{u})$ . Moreover,  $\bar{y} \in W^{2,p}(\Omega)$  and  $\bar{\varphi} \in W^{2,r}(\Omega)$  for any  $p, r \in (2, p_*)$  and  $r \leq \bar{p}$  with  $\bar{p}$  and  $p_*$ , respectively, defined in Assumption (A4) and Theorem 2.2. Consequently,  $\bar{y}, \bar{\varphi} \in C^1(\bar{\Omega})$  and  $\bar{u} \in C^{0,1}(\bar{\Omega})$ .

Assume now that  $\bar{\varphi} \in H_0^1(\Omega)$  fulfills (2.7). The *critical cone* of the problem (P) at  $\bar{u}$  is defined by

$$(2.8) \quad C(\mathcal{U}_{ad}; \bar{u}) := \{v \in L^2(\Omega) \mid v \geq 0 \text{ if } \bar{u} = \alpha, v \leq 0 \text{ if } \bar{u} = \beta, v = 0 \text{ if } \bar{\varphi} + v\bar{u} \neq 0 \text{ a.e. in } \Omega\}.$$

In the rest of this subsection, we shall provide second-order sufficient optimality conditions for (P). On that account, we need introduce the curvature functional of  $j$ , which can be separated into three contributions. For any  $(u, y, \varphi) \in L^2(\Omega) \times H^1(\Omega) \times W^{1,\infty}(\Omega)$ , the smooth part and the first-order nonsmooth part of the curvature in direction  $(v_1, v_2) \in L^2(\Omega)^2$  are defined by

$$\begin{aligned} Q_s(u, y, \varphi; v_1, v_2) &:= \frac{1}{2} \int_{\Omega} \frac{\partial^2 L}{\partial y^2}(\cdot, y) z_{v_1} z_{v_2} \, dx + \frac{\nu}{2} \int_{\Omega} v_1 v_2 \, dx - \frac{1}{2} \int_{\Omega} \mathbb{1}_{\{y \neq \bar{t}\}} a''(y) z_{v_1} z_{v_2} \nabla y \cdot \nabla \varphi \, dx, \\ Q_1(u, y, \varphi; v_1, v_2) &:= -\frac{1}{2} \int_{\Omega} [a'(y; z_{v_1}) \nabla z_{v_2} + a'(y; z_{v_2}) \nabla z_{v_1}] \cdot \nabla \varphi \, dx \end{aligned}$$

with  $z_{v_i} := S'(u)v_i$ ,  $i = 1, 2$ . The critical part of the curvature is of course the second-order nonsmooth part involving some additional notation. Let  $\delta > 0$  be arbitrary but fixed and set

$$(2.9) \quad \begin{cases} \Omega_{y, \hat{y}}^2 := \{\hat{y} \in (\bar{t}, \bar{t} + \delta), y \in (\bar{t} - \delta, \bar{t})\}, \\ \Omega_{y, \hat{y}}^3 := \{\hat{y} \in (\bar{t} - \delta, \bar{t}), y \in [\bar{t}, \bar{t} + \delta)\} \end{cases}$$

for given functions  $y, \hat{y} \in C(\bar{\Omega})$ . For any  $s \in \mathbb{R}$ ,  $u, v \in L^2(\Omega)$ ,  $y \in C(\bar{\Omega}) \cap H^1(\Omega)$ , and  $\varphi \in W^{1,\infty}(\Omega)$ , we define

$$(2.10) \quad \begin{cases} \zeta_0(u, y; s, v) := -\{a'\}_{\bar{t}+0}^{\bar{t}-0}(\bar{t} - S(u + sv)) \mathbb{1}_{\Omega_{S(u+sv), y}^3}, \\ \zeta_1(u, y; s, v) := \{a'\}_{\bar{t}+0}^{\bar{t}-0}(\bar{t} - S(u + sv)) \mathbb{1}_{\Omega_{S(u+sv), y}^2}, \\ \zeta(u, y; s, v) := \zeta_0(u, y; s, v) + \zeta_1(u, y; s, v) = \{a'\}_{\bar{t}+0}^{\bar{t}-0}(\bar{t} - S(u + sv)) \left[ \mathbb{1}_{\Omega_{S(u+sv), y}^2} - \mathbb{1}_{\Omega_{S(u+sv), y}^3} \right]. \end{cases}$$

Here  $\{a'\}_{\bar{t}+0}^{\bar{t}-0}$  denotes the difference between the one-sided derivatives of  $a$  at  $\bar{t}$  from left and right, i.e.,

$$\{a'\}_{\bar{t}+0}^{\bar{t}-0} := \lim_{t \rightarrow \bar{t}^-} a'(t) - \lim_{t \rightarrow \bar{t}^+} a'(t) = a'_0(\bar{t}) - a'_1(\bar{t}).$$

We then determine for any  $\{s_n\} \in c_0^+ := \{\{s_n\} \subset (0, \infty) \mid s_n \rightarrow 0\}$  and  $v \in L^2(\Omega)$  the term

$$(2.11) \quad \begin{aligned} \tilde{Q}(u, y, \varphi; \{s_n\}, v) &:= \liminf_{n \rightarrow \infty} \frac{1}{s_n^2} \int_{\Omega} \sum_{i=0}^1 \zeta_i(u, y; s_n, v) \nabla y \cdot \nabla \varphi \, dx \\ &= \{a'\}_{\bar{t}+0}^{\bar{t}-0} \liminf_{n \rightarrow \infty} \frac{1}{s_n^2} \int_{\Omega} (\bar{t} - S(u + s_n v)) \left[ \mathbb{1}_{\Omega_{S(u+s_n v), y}^2} - \mathbb{1}_{\Omega_{S(u+s_n v), y}^3} \right] \nabla y \cdot \nabla \varphi \, dx. \end{aligned}$$

The second-order nonsmooth part of the curvature in direction  $v \in L^2(\Omega)$  is thus defined as

$$Q_2(u, y, \varphi; v) := \inf \{ \tilde{Q}(u, y, \varphi; \{s_n\}, v) \mid \{s_n\} \in c_0^+ \}.$$

We finally identify the total curvature in direction  $v$  as

$$(2.12) \quad Q(u, y, \varphi; v) := Q_s(u, y, \varphi; v, v) + Q_1(u, y, \varphi; v, v) + Q_2(u, y, \varphi; v).$$

**Remark 2.6.** The definitions of the sets  $\Omega_{y, \hat{y}}^2$  and  $\Omega_{y, \hat{y}}^3$  in (2.9) are derived from the ones for  $\Omega_{y, \hat{y}}^{1,2}$  and  $\Omega_{y, \hat{y}}^{0,3}$ , respectively, in Lemma 3.3 in [22] for the situation in which  $K := 1$ ,  $t_0 := -\infty$ ,  $t_1 := \bar{t}$ , and  $t_2 := \infty$ . Analogously, the definitions of  $Q_s$ ,  $Q_1$ , and  $Q_2$  in this section can be obtained from those in § 5.1 in [22].

Thanks to Proposition 5.6 and Lemma 5.7 in [22], we have the weak lower semicontinuity of  $Q_2$  in the last variable and there holds

$$|Q_2(u, S(u), \varphi; v)| \leq \Sigma(S(u)) \|\nabla \varphi\|_{L^\infty(\Omega)} \|S'(u)v\|_{L^\infty(\Omega)}^2 \quad \text{for all } u, v \in L^2(\Omega) \text{ and } \varphi \in W^{1,\infty}(\Omega).$$

Here  $\Sigma : C(\bar{\Omega}) \cap W^{1,1}(\Omega) \rightarrow \mathbb{R} \cup \{\infty\}$  is the *jump functional* and defined by

$$(2.13) \quad \Sigma(y) := \sigma_0 \limsup_{r \rightarrow 0^+} \frac{1}{r} \sum_{m=1}^2 \int_{\Omega} [\mathbb{1}_{\{0 < |y - \bar{t}| \leq r\}} |\partial_{x_m} y|] dx, \quad y \in W^{1,1}(\Omega) \cap C(\bar{\Omega})$$

with

$$(2.14) \quad \sigma_0 := |\{a'\}_{\bar{t}+0}^{\bar{t}-0}| = |a'_0(\bar{t}) - a'_1(\bar{t})|.$$

Moreover, it follows from Corollary 5.5 in [22] that, for any  $u \in L^2(\Omega)$ ,  $\{s_n\} \in c_0^+$  and  $v_n \rightharpoonup v$  in  $L^2(\Omega)$ , there holds

$$(2.15) \quad \liminf_{n \rightarrow \infty} \frac{1}{s_n^2} \int_{\Omega} \sum_{i=0}^1 \zeta_i(u, S(u); s_n, v_n) \nabla S(u) \cdot \nabla \varphi dx = \tilde{Q}(u, S(u), \varphi; \{s_n\}, v) \geq Q_2(u, S(u), \varphi; v),$$

provided that  $\Sigma(S(u)) < \infty$ .

**Theorem 2.7 (explicit second-order sufficient optimality conditions, [24], Thm. 3.23).** *Assume that Assumptions (A1) to (A4) are fulfilled. Let  $\bar{u}$  be an admissible control of (P) such that  $\{\bar{y} = \bar{t}\}$  decomposes into finitely many connected components and that on each such connected component  $C$ , either*

$$(2.16) \quad \nabla \bar{y}(x) \neq 0 \quad \text{for all } x \in C$$

or

$$(2.17) \quad \begin{cases} \nabla \bar{y} = 0 \text{ on } C, \\ \text{meas}_{\mathbb{R}^2}(\{0 < |\bar{y} - \bar{t}| < r\} \cap C^{\varepsilon_0}) \leq c_s r, \end{cases} \quad \text{for all } r \in (0, r_0), \text{ for some constants } \varepsilon_0, r_0 > 0$$

holds with  $\bar{y} := S(\bar{u})$  and

$$C^{\varepsilon_0} := \{x \in \bar{\Omega} \mid \text{dist}(x, C) < \varepsilon_0\}.$$

Assume further that there exists a  $\bar{\varphi} \in W_0^{1,\bar{p}}(\Omega) \cap W^{1,\infty}(\Omega)$ , with  $\bar{p}$  defined in Assumption (A4), that together with  $\bar{u}, \bar{y}$  fulfills (2.7) and

$$(2.18) \quad \begin{aligned} Q(\bar{u}, \bar{y}, \bar{\varphi}; v) &= \frac{1}{2} \int_{\Omega} \frac{\partial^2 L}{\partial y^2}(\cdot, \bar{y}) z_v^2 dx + \frac{\nu}{2} \int_{\Omega} v^2 dx - \frac{1}{2} \int_{\Omega} \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a''(\bar{y}) z_v^2 \nabla \bar{y} \cdot \nabla \bar{\varphi} dx \\ &\quad - \int_{\Omega} a'(\bar{y}; z_v) \nabla z_v \cdot \nabla \bar{\varphi} dx \\ &\quad + \frac{1}{2} [a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y} = \bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}| > 0\}} z_v^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) \\ &> 0 \quad \text{for all } v \in C(\mathcal{U}_{ad}; \bar{u}) \setminus \{0\} \end{aligned}$$

with  $z_v := S'(\bar{u})v$ . Then constants  $c_0, \rho_0 > 0$  exist and fulfill

$$j(\bar{u}) + c_0 \|u - \bar{u}\|_{L^2(\Omega)}^2 \leq j(u) \quad \text{for all } u \in \mathcal{U}_{ad} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \rho_0).$$

The following will be used later to show the error estimates for the approximation of (P).

**Proposition 2.8** ([24], Thm. 3.19). *Let  $\bar{u} \in L^2(\Omega)$  be arbitrary and let  $\bar{y} := S(\bar{u})$ . Assume that  $\{\bar{y} = \bar{t}\}$  decomposes into finitely many connected components and that on each such connected component  $C$ , either (2.16) or (2.17) is fulfilled. Let  $v \in L^2(\Omega)$  and  $\bar{\varphi} \in C^1(\bar{\Omega}) \cap W^{2,1}(\Omega)$ . Then, for any  $\{s_n\} \in c_0^+$  and  $\{v_n\} \subset L^2(\Omega)$  such that  $v_n \rightarrow v$  in  $L^2(\Omega)$ ,  $y_n := S(\bar{u} + s_n v_n) \rightarrow \bar{y}$  in  $C^1(\bar{\Omega})$ , and  $(y_n - \bar{y})/s_n \rightarrow w$  in  $W_0^{1,p}(\Omega)$  for some  $p > 2$ , there hold*

$$(2.19) \quad \tilde{Q}(\bar{u}, \bar{y}, \bar{\varphi}; \{s_n\}, v) = \frac{1}{2} [a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y}=\bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}|>0\}} w^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x)$$

and

$$(2.20) \quad \lim_{n \rightarrow \infty} \frac{1}{s_n^2} \int_{\Omega} (2\bar{t} - \bar{y} - y_n) [\mathbb{1}_{\Omega_{y_n, \bar{y}}^2} - \mathbb{1}_{\Omega_{y_n, \bar{y}}^3}] \nabla \bar{y} \cdot \nabla \bar{\varphi} dx = 0.$$

Finally, the wellposedness of  $\Sigma$ , defined in (2.13), is guaranteed by either (2.16) or (2.17) as stated below.

**Proposition 2.9** ([24], Prop. 3.20). *Let  $\bar{y} \in C^1(\bar{\Omega})$  be such that  $\{\bar{y} = \bar{t}\}$  decomposes into finitely many connected components and that on each such connected component  $C$ , either (2.16) or (2.17) is fulfilled. Then  $\Sigma(\bar{y}) < \infty$ .*

### 3 ANALYSIS OF THE DISCRETE STATE EQUATION

In this section, we study the discrete version of the state equation (2.3) and show error estimates of solutions to the discrete state equation (3.1), local uniqueness of these solutions, and local differentiability of the solution operators of (3.1). To this end, we introduce a family of regular triangulations  $\{\mathcal{T}_h\}_{h>0} : \bar{\Omega} = \bigcup_{T \in \mathcal{T}_h} T$  for all  $h > 0$ . For each element  $T \in \mathcal{T}_h$ , we denote by  $\varrho(T)$  and  $\delta(T)$  the diameter of  $T$  and the diameter of the largest ball contained in  $T$ , respectively. The mesh size of  $\mathcal{T}_h$  will be denoted by  $h := \max_{T \in \mathcal{T}_h} \varrho(T)$ . This triangulation is assumed to be *regular* in the sense that there exist  $\bar{\varrho}, \bar{\delta} > 0$  such that  $\frac{\varrho(T)}{\delta(T)} \leq \bar{\delta}$  and  $\frac{h}{\varrho(T)} \leq \bar{\varrho}$  for all  $T \in \mathcal{T}_h$  and  $h > 0$ ; see, e.g. [20].

We will employ the standard continuous piecewise linear finite elements for the state  $y$  and set

$$V_h := \{v_h \in C(\bar{\Omega}) \mid v_h|_T \in \mathcal{P}_1 \text{ for all } T \in \mathcal{T}_h, v_h = 0 \text{ on } \partial\Omega\},$$

where  $\mathcal{P}_1$  stands for the space of polynomials of degree equal at most 1. The discrete approximation of the state equation (2.3) for  $y_h \in V_h$  is then

$$(3.1) \quad \int_{\Omega} (b + a(y_h)) \nabla y_h \cdot \nabla v_h dx = \int_{\Omega} u v_h dx \quad \text{for all } v_h \in V_h.$$

While the existence of solutions to (3.1) follows from Theorem 3.1 in [9], the uniqueness of solutions is still an open problem. However, if  $a : \mathbb{R} \rightarrow \mathbb{R}$  is assumed to be bounded, then we have uniqueness provided that  $h$  is small enough; see Theorem 4.1 in [9]. Below, we provide some error estimates for solutions to (3.1) that are sufficiently close to the solutions of (2.3).

In what follows, we fix  $\bar{u} \in L^2(\Omega)$  and set  $\bar{y} := S(\bar{u})$ . From **Theorem 2.2** and the continuous embedding  $L^2(\Omega) \hookrightarrow W^{-1,p}(\Omega)$  for any  $p > 1$ , we then have  $\bar{y} \in W_0^{1,p}(\Omega) \cap H^2(\Omega)$ .

**Theorem 3.1** ([16], Thm. 3.1). *Let  $\rho_0 > 0$  be arbitrary but fixed,  $U := \bar{B}_{L^2(\Omega)}(\bar{u}, \rho_0)$ , and let  $p \geq 2$ . Assume that **Assumptions (A1) to (A3)** are fulfilled. Then there exists a constant  $h_0 > 0$  such that for any  $u \in U$  and  $h < h_0$ , there exists at least one solution  $y_h(u)$  to (3.1) satisfying for  $y_u := S(u)$*

$$(3.2) \quad \|y_u - y_h(u)\|_{L^2(\Omega)} + h \|y_u - y_h(u)\|_{H_0^1(\Omega)} + h \|y_u - y_h(u)\|_{L^\infty(\Omega)} \leq C_U h^2,$$

$$(3.3) \quad \|y_u - y_h(u)\|_{W_0^{1,p}(\Omega)} \leq C_{U,p} h^{2/p}.$$

*Proof.* The estimates for the norms in  $L^2$ ,  $H_0^1$ , and  $W^{1,p}$  are shown in Theorem 3.1 in [16], while the estimate for the  $L^\infty$  norm can be obtained similar to estimate (3.11) in [16].  $\square$

The following theorem guarantees the local uniqueness of solutions to (3.1). Its proof is similar to that of Theorem 4.2 in [10] with slight modifications and is thus omitted here.

**Theorem 3.2.** *Let  $p > 2$  be arbitrary and let  $h_0$  be defined in Theorem 3.1. Under Assumptions (A1) to (A3), there exist  $h_1 \in (0, h_0)$ ,  $\rho > 0$ , and  $\kappa_\rho > 0$  such that for any  $h < h_1$  and any  $u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho)$ , (3.1) admits a unique solution in  $\overline{B}_{W_0^{1,p}(\Omega)}(\bar{y}, \kappa_\rho) \cap V_h$ .*

From now on, let us fix  $\tilde{p} \geq 4$  and let  $h_0, h_1, \rho$ , and  $\kappa_\rho$  be the constants defined in Theorems 3.1 and 3.2 for  $p = \tilde{p}$ . In the rest of this section, we shall investigate the differentiability of the discrete solution operator

$$(3.4) \quad S_h : B_{L^2(\Omega)}(\bar{u}, \rho) \ni u \mapsto y_h(u) \in \overline{B}_{W_0^{1,\tilde{p}}(\Omega)}(\bar{y}, \kappa_\rho) \cap V_h,$$

where  $y_h(u)$  is the unique solution to (3.1) in  $\overline{B}_{W_0^{1,\tilde{p}}(\Omega)}(\bar{y}, \kappa_\rho)$  from Theorem 3.2.

For any  $y, \hat{y} \in C(\overline{\Omega}) \cap W^{1,1}(\Omega)$ , we define functions  $T_{y,\hat{y}}$  and  $Z_{y,\hat{y}}$  on  $\Omega$  via

$$(3.5) \quad T_{y,\hat{y}} := \mathbb{1}_{\{\hat{y} \neq \bar{t}\}} [a(y) - a(\hat{y}) - a'(\hat{y})(y - \hat{y})]$$

and

$$(3.6) \quad Z_{y,\hat{y}} := \mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y - \mathbb{1}_{\{\hat{y} \neq \bar{t}\}} a'(\hat{y}) \nabla \hat{y}.$$

In order to prove the differentiability of  $S_h$ , we need the following lemmas.

**Lemma 3.3** ([22], Lem. 3.3). *Let Assumption (A3) be fulfilled. Assume that  $y_n \rightarrow y$  in  $W_0^{1,p}(\Omega)$  as  $n \rightarrow \infty$  with  $p > 2$ . Then*

$$\frac{1}{\|y_n - y\|_{W_0^{1,p}(\Omega)}} \|T_{y_n, y} \nabla y\|_{L^p(\Omega)} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

**Lemma 3.4.** *Let Assumption (A3) be fulfilled and let  $y, \hat{y} \in C(\overline{\Omega}) \cap W^{1,1}(\Omega)$  and  $M > 0$  be arbitrary such that  $\|y - \hat{y}\|_{C(\overline{\Omega})} < \delta$  with  $\delta$  defined in (2.9) and  $\|y\|_{C(\overline{\Omega})}, \|\hat{y}\|_{C(\overline{\Omega})} \leq M$ . Then*

$$(3.7) \quad Z_{y,\hat{y}} = Z_{y,\hat{y}}^{(1)} + Z_{y,\hat{y}}^{(2)} + Z_{y,\hat{y}}^{(3)} + Z_{y,\hat{y}}^{(4)}$$

for

$$\begin{aligned} Z_{y,\hat{y}}^{(1)} &:= \mathbb{1}_{\{\hat{y} \in (-\infty, \bar{t}), y \in (-\infty, \bar{t})\}} [a'_0(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] + \mathbb{1}_{\{\hat{y} \in (\bar{t}, \infty), y \in (\bar{t}, \infty)\}} [a'_1(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}], \\ Z_{y,\hat{y}}^{(2)} &:= \mathbb{1}_{\{\hat{y} = \bar{t}\}} \mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y, \quad Z_{y,\hat{y}}^{(3)} := [a'_0(\bar{t}) - a'_1(\bar{t})] [\mathbb{1}_{\Omega_{y,\hat{y}}^2} - \mathbb{1}_{\Omega_{y,\hat{y}}^3}] \nabla \hat{y}, \end{aligned}$$

and

$$\begin{aligned} Z_{y,\hat{y}}^{(4)} &:= \mathbb{1}_{\Omega_{y,\hat{y}}^2} [a'_0(y) \nabla(y - \hat{y}) + (a'_0(y) - a'_0(\bar{t})) \nabla \hat{y} + (a'_1(\bar{t}) - a'_1(\hat{y})) \nabla \hat{y}] \\ &\quad + \mathbb{1}_{\Omega_{y,\hat{y}}^3} [a'_1(y) \nabla(y - \hat{y}) + (a'_1(y) - a'_1(\bar{t})) \nabla \hat{y} + (a'_0(\bar{t}) - a'_0(\hat{y})) \nabla \hat{y}], \end{aligned}$$

with the sets  $\Omega_{y,\hat{y}}^2$  and  $\Omega_{y,\hat{y}}^3$  defined as in (2.9). Moreover, there exists a constant  $C_M > 0$  such that a.e. in  $\Omega$ ,

$$(3.8) \quad \begin{cases} |Z_{y,\hat{y}}^{(1)}| + |Z_{y,\hat{y}}^{(2)}| \leq C_M [ \|y - \hat{y}\| |\nabla \hat{y}| + |\nabla(y - \hat{y})| ], \\ |Z_{y,\hat{y}}^{(4)}| \leq C_M [ \|y - \hat{y}\| |\nabla \hat{y}| + |\nabla(y - \hat{y})| ] (\mathbb{1}_{\Omega_{y,\hat{y}}^2} + \mathbb{1}_{\Omega_{y,\hat{y}}^3}). \end{cases}$$

Consequently,  $Z_{y,\hat{y}} \rightarrow 0$  in  $L^p(\Omega)$  as  $y \rightarrow \hat{y}$  in  $W^{1,p}(\Omega) \cap C(\overline{\Omega})$  for any  $p \geq 1$ .

*Proof.* Clearly, we deduce from the fact  $\nabla \hat{y} = 0$  a.e. in  $\{\hat{y} = \bar{t}\}$  (see [18]) that

$$(3.9) \quad Z_{y,\hat{y}} = \mathbb{1}_{\{\hat{y} \in (-\infty, \bar{t})\}} [\mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] + \mathbb{1}_{\{\hat{y} \in (\bar{t}, \infty)\}} [\mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}] \\ + \mathbb{1}_{\{\hat{y} = \bar{t}\}} \mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y = Z_0^1 + Z_1^1 + Z_{y,\hat{y}}^{(2)}$$

with

$$Z_0^1 := \mathbb{1}_{\{\hat{y} \in (-\infty, \bar{t})\}} [\mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] \quad \text{and} \quad Z_1^1 := \mathbb{1}_{\{\hat{y} \in (\bar{t}, \infty)\}} [\mathbb{1}_{\{y \neq \bar{t}\}} a'(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}].$$

Since  $\|y - \hat{y}\|_{C(\bar{\Omega})} < \delta$  and  $\nabla y = 0$  a.e. in  $\{y = \bar{t}\}$ , we can write

$$Z_0^1 = \mathbb{1}_{\{\hat{y} \in (-\infty, \bar{t}), y \in (-\infty, \bar{t})\}} [a'_0(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] + \mathbb{1}_{\{\hat{y} \in (\bar{t}-\delta, \bar{t}), y \in [\bar{t}, \bar{t}+\delta)\}} [a'_1(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] \\ =: Z_0^{1,2} + Z_0^{1,1}$$

and

$$Z_1^1 = \mathbb{1}_{\{\hat{y} \in (\bar{t}, \bar{t}+\delta), y \in (\bar{t}-\delta, \bar{t}]\}} [a'_0(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}] + \mathbb{1}_{\{\hat{y} \in (\bar{t}, \infty), y \in (\bar{t}, \infty)\}} [a'_1(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}] \\ =: Z_1^{1,1} + Z_1^{1,2}.$$

Thus, we have from the definition of  $Z_{y,\hat{y}}^{(1)}$  that

$$(3.10) \quad Z_0^{1,2} + Z_1^{1,2} = \mathbb{1}_{\{\hat{y} \in (-\infty, \bar{t}), y \in (-\infty, \bar{t})\}} [a'_0(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] + \mathbb{1}_{\{\hat{y} \in (\bar{t}, \infty), y \in (\bar{t}, \infty)\}} [a'_1(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}] \\ = Z_{y,\hat{y}}^{(1)}.$$

By using the definition of  $\Omega_{y,\hat{y}}^3$  in (2.9), we now write

$$Z_0^{1,1} = \mathbb{1}_{\{\hat{y} \in (\bar{t}-\delta, \bar{t}), y \in [\bar{t}, \bar{t}+\delta)\}} [a'_1(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] = \mathbb{1}_{\Omega_{y,\hat{y}}^3} [a'_1(y) \nabla y - a'_0(\hat{y}) \nabla \hat{y}] \\ = \mathbb{1}_{\Omega_{y,\hat{y}}^3} [a'_1(\bar{t}) - a'_0(\bar{t})] \nabla \hat{y} + \tilde{Z}_0^{1,1}$$

with

$$\tilde{Z}_0^{1,1} := \mathbb{1}_{\Omega_{y,\hat{y}}^3} [a'_1(y) \nabla(y - \hat{y}) + (a'_1(y) - a'_1(\bar{t})) \nabla \hat{y} + (a'_0(\bar{t}) - a'_0(\hat{y})) \nabla \hat{y}].$$

Similarly, the definition of  $\Omega_{y,\hat{y}}^2$  in (2.9) implies that

$$Z_1^{1,1} = \mathbb{1}_{\{\hat{y} \in (\bar{t}, \bar{t}+\delta), y \in (\bar{t}-\delta, \bar{t}]\}} [a'_0(y) \nabla y - a'_1(\hat{y}) \nabla \hat{y}] \\ = \mathbb{1}_{\Omega_{y,\hat{y}}^2} [a'_0(\bar{t}) - a'_1(\bar{t})] \nabla \hat{y} + \tilde{Z}_1^{1,1}$$

with

$$\tilde{Z}_1^{1,1} := \mathbb{1}_{\Omega_{y,\hat{y}}^2} [a'_0(y) \nabla(y - \hat{y}) + (a'_0(y) - a'_0(\bar{t})) \nabla \hat{y} + (a'_1(\bar{t}) - a'_1(\hat{y})) \nabla \hat{y}].$$

Obviously, we have

$$\tilde{Z}_0^{1,1} + \tilde{Z}_1^{1,1} = Z_{y,\hat{y}}^{(4)}$$

and there then holds

$$Z_{y,\hat{y}}^{(3)} + Z_{y,\hat{y}}^{(4)} = Z_0^{1,1} + Z_1^{1,1}.$$

From this and (3.9)–(3.10), we derive (3.7). Moreover, (3.8) is derived by combining the definition of  $Z_{y,\hat{y}}^{(k)}$ , Assumption (A3), the estimates

$$|y(x) - \bar{t}|, |\hat{y}(x) - \bar{t}| \leq |y(x) - \hat{y}(x)| \quad \text{for a.e. } x \in \Omega_{y,\hat{y}}^2 \cup \Omega_{y,\hat{y}}^3$$

due to the definition of  $\Omega_{y,\hat{y}}^2$  and  $\Omega_{y,\hat{y}}^3$ , and the fact that  $\nabla \hat{y} = 0$  a.e. in  $\{\hat{y} = \bar{t}\}$  (see; e.g. Remark 2.6 in [18]). Finally, the claimed convergence follows from (3.7), (3.8), the fact that  $\mathbb{1}_{\Omega_{y,\hat{y}}^2}, \mathbb{1}_{\Omega_{y,\hat{y}}^3} \rightarrow 0$  a.e. in  $\Omega$  as  $y \rightarrow \hat{y}$  in  $C(\bar{\Omega})$ , and Lebesgue's dominated convergence theorem.  $\square$

For any  $h \in (0, h_1)$  and  $y_h \in V_h$ , we now define the operator  $D_{h, y_h} : V_h \rightarrow V_h^*$  via

$$(3.11) \quad \langle D_{h, y_h} w_h, z_h \rangle := \int_{\Omega} [(b + a(y_h)) \nabla w_h + \mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h) \nabla y_h w_h] \cdot \nabla z_h \, dx, \quad w_h, z_h \in V_h.$$

**Lemma 3.5.** *Let all assumptions of Theorem 3.1 hold. Then for any  $h \in (0, h_1)$  and any  $\{y_h^k\} \subset V_h$  converging to  $y_h \in V_h$  in  $H_0^1(\Omega)$  as  $k \rightarrow \infty$ , there holds  $\|D_{h, y_h^k} - D_{h, y_h}\|_{\mathbb{L}(V_h, V_h^*)} \rightarrow 0$ .*

*Proof.* Let  $w_h, v_h \in V_h$  be arbitrary such that  $\|w_h\|_{H_0^1(\Omega)}, \|v_h\|_{H_0^1(\Omega)} \leq 1$  and  $h \in (0, h_1)$  be arbitrary but fixed. Assume that  $\{y_h^k\} \subset V_h$  converges to  $y_h \in V_h$  in  $H_0^1(\Omega)$  as  $k \rightarrow \infty$ . By virtue of the inverse inequality [20], Thm. 3.2.6, we deduce that  $y_h^k \rightarrow y_h$  in  $W_0^{1, \tilde{p}}(\Omega)$  and hence in  $C(\bar{\Omega})$  as  $k \rightarrow \infty$ . We can therefore assume that  $\|y_h^k - y_h\|_{C(\bar{\Omega})} < \delta$  for all  $k \in \mathbb{N}$  large enough. On the other hand, we have

$$\langle (D_{h, y_h^k} - D_{h, y_h}) w_h, v_h \rangle = \int_{\Omega} [(a(y_h^k) - a(y_h)) \nabla w_h + Z_{y_h^k, y_h} w_h] \cdot \nabla v_h \, dx.$$

Together with the Hölder inequality, this yields that

$$\begin{aligned} \|D_{h, y_h^k} - D_{h, y_h}\|_{\mathbb{L}(V_h, V_h^*)} &\leq \|a(y_h^k) - a(y_h)\|_{L^\infty(\Omega)} + \|Z_{y_h^k, y_h}\|_{L^{\tilde{p}}(\Omega)} \|w_h\|_{L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)} \\ &\leq \|a(y_h^k) - a(y_h)\|_{L^\infty(\Omega)} + C \|Z_{y_h^k, y_h}\|_{L^{\tilde{p}}(\Omega)}, \end{aligned}$$

where we have employed the continuous embedding  $H_0^1(\Omega) \hookrightarrow L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)$  and the fact that  $\|w_h\|_{H_0^1(\Omega)} \leq 1$  to obtain the last inequality. The first term on the right-hand side of the last estimate tends to zero as  $k \rightarrow \infty$  since  $y_h^k \rightarrow y_h$  in  $C(\bar{\Omega})$  as  $k \rightarrow \infty$ . Moreover, the second term tends to zero as a result of Lemma 3.4.  $\square$

**Lemma 3.6.** *Let all assumptions of Theorem 3.1 hold. Then there exists a constant  $h_2 \in (0, h_1)$  such that for any  $h \in (0, h_2)$  and any  $y_h \in \bar{B}_{W_0^{1, \tilde{p}}(\Omega)}(\bar{y}, \kappa_\rho) \cap V_h$ , the operator  $D_{h, y_h} : V_h \rightarrow V_h^*$  is an isomorphism.*

*Proof.* Since  $V_h$  is finite-dimensional and  $D_{h, y_h}$  is linear, it suffices to prove that there exists an  $h_2 \in (0, h_1)$  such that for any  $h \in (0, h_2)$  and  $y_h \in \bar{B}_{W_0^{1, \tilde{p}}(\Omega)}(\bar{y}, \kappa_\rho) \cap V_h$ , the equation

$$(3.12) \quad D_{h, y_h} w_h = 0$$

admits the unique solution  $w_h = 0$ . We argue by contradiction. Assume for any  $k \geq 1$  that there exist  $h_k \in (0, h_1)$ ,  $y_{h_k} \in \bar{B}_{W_0^{1, \tilde{p}}(\Omega)}(\bar{y}, \kappa_\rho) \cap V_{h_k}$ , and  $w_{h_k} \in V_{h_k} \setminus \{0\}$  such that  $h_k \rightarrow 0^+$  and  $w_{h_k}$  solves (3.12) for  $h = h_k$  and  $y_h = y_{h_k}$ . By setting  $\hat{w}_{h_k} := \frac{w_{h_k}}{\|w_{h_k}\|_{L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)}}$ , we deduce that

$$(3.13) \quad \|\hat{w}_{h_k}\|_{L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)} = 1 \quad \text{and} \quad D_{h_k, y_{h_k}} \hat{w}_{h_k} = 0.$$

Furthermore, as a result of the embedding  $W_0^{1, \tilde{p}}(\Omega) \Subset C(\bar{\Omega})$ , there hold that  $\|y_{h_k}\|_{C(\bar{\Omega})} \leq M$  for all  $k \geq 1$  and some constant  $M > 0$  independent of  $k$  and that

$$(3.14) \quad y_{h_k} \rightarrow y \text{ in } C(\bar{\Omega}) \quad \text{for some } y \in W_0^{1, \tilde{p}}(\Omega).$$

Testing the second equation in (3.13) by  $\hat{w}_{h_k}$ , Hölder's inequality thus gives

$$\underline{b} \|\nabla \hat{w}_{h_k}\|_{L^2(\Omega)} \leq \|\nabla y_{h_k}\|_{L^{\tilde{p}}(\Omega)} \|\mathbb{1}_{\{y_{h_k} \neq \bar{t}\}} a'(y_{h_k})\|_{L^\infty(\Omega)} \|\hat{w}_{h_k}\|_{L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)} \leq C_{M, \rho}$$

for some constant  $C_{M,\rho} > 0$ . From this and the compact embedding  $H_0^1(\Omega) \Subset L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)$ , a subsequence argument shows that we can assume that

$$(3.15) \quad \hat{w}_{h_k} \rightharpoonup \hat{w} \text{ in } H_0^1(\Omega) \quad \text{and} \quad \hat{w}_{h_k} \rightarrow \hat{w} \text{ in } L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)$$

for some  $\hat{w} \in H_0^1(\Omega)$ . Moreover, there exist an element  $\mathbb{b} \in L^{\tilde{p}}(\Omega)^2$  and a subsequence of  $\{\mathbb{b}_k\}$  with  $\mathbb{b}_k := \mathbb{1}_{\{y_{h_k} \neq \bar{i}\}} a'(y_{h_k}) \nabla y_{h_k}$ , denoted in the same way, such that  $\mathbb{b}_k \rightharpoonup \mathbb{b}$  weakly in  $L^{\tilde{p}}(\Omega)^2$ . By fixing any  $v \in H^2(\Omega) \cap H_0^1(\Omega)$  and testing the last equation in (3.13) with  $v_{h_k} := \Pi_{h_k} v \in V_{h_k}$ , where  $\Pi_{h_k}$  is the interpolation operator, we have

$$\int_{\Omega} [(b + a(y_{h_k})) \nabla \hat{w}_{h_k} + \hat{w}_{h_k} \mathbb{b}_k] \cdot \nabla v_{h_k} \, dx = 0 \quad \text{for all } k \geq 1.$$

Letting  $k \rightarrow \infty$  and exploiting the limits (3.14), (3.15),  $\mathbb{b}_k \rightharpoonup \mathbb{b}$  in  $L^{\tilde{p}}(\Omega)^N$ , and  $v_{h_k} \rightarrow v$  in  $H_0^1(\Omega)$ , we can conclude that  $\int_{\Omega} [(b + a(y)) \nabla \hat{w} + \hat{w} \mathbb{b}] \cdot \nabla v \, dx = 0$ . From this, the density of  $H^2(\Omega) \cap H_0^1(\Omega)$  in  $H_0^1(\Omega)$ , and Theorem 2.6 in [9], we conclude that  $\hat{w} = 0$ , contradicting the fact that  $\|\hat{w}\|_{L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)} = \lim_{k \rightarrow \infty} \|\hat{w}_{h_k}\|_{L^{2\tilde{p}/(\tilde{p}-2)}(\Omega)} = 1$ .  $\square$

As a consequence of Lemmas 3.5 and 3.6 and the implicit function theorem, we obtain the differentiability of  $S_h$ .

**Theorem 3.7.** *Let all assumptions of Theorem 3.1 hold. Then, for any  $h \in (0, h_2)$ , the operator  $S_h$  defined in (3.4) is of class  $C^1$ . Moreover, for any  $u \in B_{L^2(\Omega)}(\bar{u}, \rho)$ , let  $y_h(u) := S_h(u)$ . Then for any  $v \in L^2(\Omega)$ , the Fréchet derivative  $S'_h(u)v =: z_h$  is the unique solution to*

$$(3.16) \quad \int_{\Omega} [(b + a(y_h(u))) \nabla z_h + \mathbb{1}_{\{y_h(u) \neq \bar{i}\}} a'(y_h(u)) z_h \nabla y_h(u)] \cdot \nabla w_h \, dx = \int_{\Omega} v w_h \, dx \quad \text{for all } w_h \in V_h.$$

*Proof.* We first consider for any  $h \in (0, h_2)$  the mapping  $F_h : B_{L^2(\Omega)}(\bar{u}, \rho) \times V_h \rightarrow V_h^*$  defined via

$$(3.17) \quad \langle F_h(u, y_h), v_h \rangle = \int_{\Omega} (b + a(y_h)) \nabla y_h \cdot \nabla v_h - u v_h \, dx, \quad u \in B_{L^2(\Omega)}(\bar{u}, \rho), y_h, v_h \in V_h.$$

Clearly,  $F_h(u, y_h(u)) = 0$  and  $F_h$  is continuously partially differentiable in  $u$ . We now prove that  $F_h$  is partially differentiable in  $y_h$  with  $\frac{\partial F_h}{\partial y_h}(u, y_h) = D_{h,y_h}$ , where  $D_{h,y_h}$  is defined in (3.11). We thus derive the differentiability of  $S_h$  according to Lemmas 3.5 and 3.6 as well as a simple computation. To this end, by taking any  $v_h \in V_h$  and  $\{w_h^k\} \subset V_h$  with  $\|w_h^k\|_{H_0^1(\Omega)} \rightarrow 0$  as  $k \rightarrow \infty$  and  $\|v_h\|_{H_0^1(\Omega)} \leq 1$ , we deduce from a straightforward computation that

$$\langle F_h(u, y_h + w_h^k) - F_h(u, y_h) - D_{h,y_h} w_h^k, v_h \rangle = \int_{\Omega} [T_{y_h^k, y_h} \nabla y_h + (a(y_h^k) - a(y_h)) \nabla w_h^k] \cdot \nabla v_h \, dx,$$

where  $y_h^k := y_h + w_h^k$  and  $T_{y,y}$  is defined in (3.5). This gives

$$\|F_h(u, y_h + w_h^k) - F_h(u, y_h) - D_{h,y_h} w_h^k\|_{\mathbb{L}(V_h, V_h^*)} \leq \|T_{y_h^k, y_h} \nabla y_h\|_{L^2(\Omega)} + \|a(y_h^k) - a(y_h)\|_{L^\infty(\Omega)} \|w_h^k\|_{H_0^1(\Omega)}.$$

Moreover, in view of inverse estimates [20], Thm. 3.2.6, we have  $y_h^k \rightarrow y_h$  in  $W_0^{1,\tilde{p}}(\Omega)$  and hence in  $C(\bar{\Omega})$  as  $k \rightarrow \infty$ . Then Lemma 3.3 and the embedding  $W_0^{1,\tilde{p}}(\Omega) \hookrightarrow H_0^1(\Omega)$  imply that

$$\frac{1}{\|w_h^k\|_{H_0^1(\Omega)}} \|F_h(u, y_h + w_h^k) - F_h(u, y_h) - D_{h,y_h} w_h^k\|_{\mathbb{L}(V_h, V_h^*)} \rightarrow 0,$$

which gives that  $\frac{\partial F_h}{\partial y_h}(u, y_h) = D_{h,y_h}$ . We have shown that  $F_h(u, S_h(u)) = 0$  and  $\frac{\partial F_h}{\partial y_h}(u, y_h) = D_{h,y_h}$ . We then deduce from the Implicit Function Theorem and Lemmas 3.5 and 3.6 that  $S_h$  is of class  $C^1$ . Finally, (3.16) follows from (3.11) and (3.17).  $\square$

#### 4 NUMERICAL ANALYSIS OF THE ADJOINT STATE EQUATION

In this section, we will carry out the numerical analysis of the adjoint equation (2.5). For any  $h \in (0, h_2)$ ,  $u \in B_{L^2(\Omega)}(\bar{u}, \rho)$ ,  $v \in L^2(\Omega)$  and  $y_h := S_h(u)$ , we approximate (2.5) using the triangulation  $\mathcal{T}_h$  by

$$(4.1) \quad \int_{\Omega} (b + a(y_h)) \nabla \varphi_h \cdot \nabla w_h + \mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h) w_h \nabla y_h \cdot \nabla \varphi_h \, dx = \int_{\Omega} v w_h \, dx \quad \text{for all } w_h \in V_h.$$

From the bijectivity of  $D_{h, y_h}$  shown in Lemma 3.6, we deduce the existence and uniqueness of solutions to (4.1).

**Theorem 4.1.** *Let all assumptions of Theorem 3.7 hold. Then for all  $h \in (0, h_2)$ ,  $u \in B_{L^2(\Omega)}(\bar{u}, \rho)$ , and  $v \in L^2(\Omega)$ , there exists a unique solution  $\varphi_h \in V_h$  to (4.1).*

In order to derive error estimates for the full approximation (4.1) of (2.5), we first consider the continuous problem (2.5) with  $y_h(u)$  in place of  $y_u$ .

**Lemma 4.2.** *Let all assumptions of Theorem 3.7 hold. Then for any  $h \in (0, h_2)$ ,  $u \in B_{L^2(\Omega)}(\bar{u}, \rho)$ ,  $y_h := S_h(u)$ , and  $v \in L^2(\Omega)$ , the equation*

$$(4.2) \quad -\operatorname{div}[(b + a(y_h)) \nabla \tilde{\varphi}] + \mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h) \nabla y_h \cdot \nabla \tilde{\varphi} = v \quad \text{in } \Omega, \quad \tilde{\varphi} = 0 \quad \text{on } \partial\Omega,$$

has a unique solution  $\tilde{\varphi}$  in  $H^2(\Omega) \cap H_0^1(\Omega)$ . Moreover,

$$(4.3) \quad \|\varphi - \tilde{\varphi}\|_{H^2(\Omega)} \leq C_\rho h \|v\|_{L^2(\Omega)} \quad \text{and} \quad \|\varphi - \tilde{\varphi}\|_{L^2(\Omega)} \leq C_\rho h^2 \|v\|_{L^2(\Omega)}$$

for some constant  $C_\rho$  independent of  $u, v$ , and  $h$ , where  $\varphi$  is the unique solution to (2.5).

*Proof.* From Theorem 2.2, the continuous embedding  $W_0^{1, \tilde{p}}(\Omega) \hookrightarrow C(\bar{\Omega})$ , and (3.3) for  $p := \tilde{p} \geq 4$ , there holds

$$\|\mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h) \nabla y_h\|_{L^{\tilde{p}}(\Omega)} + \|b + a(y_h(u))\|_{W^{1, \tilde{p}}(\Omega)} \leq C_\rho \quad \text{for all } h \in (0, h_2), u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho).$$

A standard argument then proves the existence of solutions  $\tilde{\varphi}$  to (4.2) in  $H^2(\Omega) \cap H_0^1(\Omega)$ ; see. e.g. Theorem 2.6 in [9] and the proof of Lemma 4.1 in [22]. Moreover, we have

$$(4.4) \quad \|\tilde{\varphi}\|_{H^2(\Omega)} \leq C_\rho \|v\|_{L^2(\Omega)}.$$

Setting  $\psi := \varphi - \tilde{\varphi}$  and subtracting the equations corresponding to  $\varphi$  and  $\tilde{\varphi}$  yields

$$(4.5) \quad -\operatorname{div}[(b + a(y_u)) \nabla \psi] + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \nabla y_u \cdot \nabla \psi = g_{u, h} \quad \text{in } \Omega, \quad \psi = 0 \quad \text{on } \partial\Omega,$$

with

$$g_{u, h} := \operatorname{div}[(a(y_u) - a(y_h)) \nabla \tilde{\varphi}] + Z_{y_h, y_u} \cdot \nabla \tilde{\varphi}.$$

By the chain rule [33], Thm. 7.8 and the fact that  $y_u, y_h \in W_0^{1, \tilde{p}}(\Omega)$  and that  $\tilde{\varphi} \in H^2(\Omega)$ , we can write

$$(4.6) \quad g_{u, h} = (a(y_u) - a(y_h)) \Delta \tilde{\varphi}.$$

Similar to (4.4), there holds

$$\|\psi\|_{H^2(\Omega)} \leq C_\rho \|g_{u, h}\|_{L^2(\Omega)} \leq C_\rho \|a(y_u) - a(y_h)\|_{L^\infty(\Omega)} \|\Delta \tilde{\varphi}\|_{L^2(\Omega)}.$$

Combining this with the Lipschitz continuity of  $a$  on bounded sets, the  $L^\infty$ -estimate in (3.2), and (4.4) yields the first estimate in (4.3). To show the second estimate, set  $z_{u, \psi} := S'(u)\psi$  and note that  $\psi = S'(u)^* g_{u, h}$ . We then deduce from (4.6) that

$$(4.7) \quad \|\psi\|_{L^2(\Omega)}^2 = \int_{\Omega} g_{u, h} z_{u, \psi} \, dx \leq C_\rho \|z_{u, \psi}\|_{L^\infty(\Omega)} \|\Delta \tilde{\varphi}\|_{L^2(\Omega)} \|y_u - y_h\|_{L^2(\Omega)}.$$

By [Theorem 2.2](#) and the compact embedding  $L^2(\Omega) \Subset W^{-1,\tilde{p}}(\Omega)$ , we have

$$(4.8) \quad \sup\{\|S'(u)\|_{\mathbb{L}(W^{-1,\tilde{p}}(\Omega), W_0^{1,\tilde{p}}(\Omega))} \mid u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho)\} \leq C_\rho.$$

The continuous embeddings  $W_0^{1,\tilde{p}}(\Omega) \hookrightarrow L^\infty(\Omega)$  and  $L^2(\Omega) \hookrightarrow W^{-1,\tilde{p}}(\Omega)$  therefore yield

$$\|z_{u,\psi}\|_{L^\infty(\Omega)} \leq C\|z_{u,\psi}\|_{W_0^{1,\tilde{p}}(\Omega)} \leq C_\rho\|\psi\|_{W^{-1,\tilde{p}}(\Omega)} \leq C_\rho\|\psi\|_{L^2(\Omega)}.$$

The inequality (4.7) thus yields

$$\|\psi\|_{L^2(\Omega)} \leq C_\rho\|\Delta\tilde{\varphi}\|_{L^2(\Omega)}\|y_u - y_h\|_{L^2(\Omega)}.$$

This, (3.2) and (4.4) yield the last estimate in (4.3).  $\square$

Below, we shall estimate the term  $Z_{y_h,y}$  defined in (3.6). We first observe from the  $L^\infty$ -error estimate in (3.2) that

$$(4.9) \quad \|S(u) - S_h(u)\|_{L^\infty(\Omega)} \leq C_\infty h \quad \text{for all } u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}, h \in (0, h_2)$$

for some positive constant  $C_\infty$ . For any  $y \in W^{1,1}(\Omega) \cap C(\overline{\Omega})$  and  $r > 0$ , let

$$(4.10) \quad V(y, r) := \sigma_0 \sum_{m=1}^2 \mathbb{1}_{\{0 < |y - \bar{t}| \leq r\}} |\partial_{x_m} y| \quad \text{and} \quad \Sigma_r(y) := \frac{1}{r} V(y, r)$$

with  $\sigma_0$  determined as in (2.14).

**Proposition 4.3.** *Let  $r > 0$ ,  $y \in W^{1,1}(\Omega) \cap C(\overline{\Omega})$ , and  $\hat{y} \in W^{1,\infty}(\Omega)$  be arbitrary and let  $\kappa := r + \|y - \hat{y}\|_{C(\overline{\Omega})}$ . Then*

$$(i) \quad V(y, r) \leq V(\hat{y}, \kappa) + \sigma_0 \sum_{m=1}^2 |\partial_{x_m} y - \partial_{x_m} \hat{y}| \text{ for a.e. in } \Omega;$$

$$(ii) \quad \|V(\hat{y}, r)\|_{L^2(\Omega)}^2 \leq 2r\sigma_0 \|\nabla \hat{y}\|_{L^\infty(\Omega)} \|\Sigma_r(\hat{y})\|_{L^1(\Omega)}.$$

*Proof.* The proof of the second claim is straightforward. It remains to prove the first assertion. To this end, we now observe that  $\{0 < |y - \bar{t}| \leq r\} \subset \{|\hat{y} - \bar{t}| \leq \kappa\}$  and  $|\partial_{x_m} y| \leq |\partial_{x_m} \hat{y}| + |\partial_{x_m} y - \partial_{x_m} \hat{y}|$ . There thus holds

$$\begin{aligned} V(y, r) &\leq \sum_{m=1}^2 \sigma_0 \mathbb{1}_{\{|\hat{y} - \bar{t}| \leq \kappa\}} [|\partial_{x_m} \hat{y}| + |\partial_{x_m} y - \partial_{x_m} \hat{y}|] \\ &= \sum_{m=1}^2 \sigma_0 \mathbb{1}_{\{|\hat{y} - \bar{t}| \leq \kappa\}} |\partial_{x_m} \hat{y}| + \sum_{m=1}^2 \sigma_0 \mathbb{1}_{\{|\hat{y} - \bar{t}| \leq \kappa\}} |\partial_{x_m} y - \partial_{x_m} \hat{y}| \\ &= V(\hat{y}, \kappa) + \sigma_0 \sum_{m=1}^2 \mathbb{1}_{\{|\hat{y} - \bar{t}| \leq \kappa\}} |\partial_{x_m} y - \partial_{x_m} \hat{y}|, \end{aligned}$$

where we have employed the fact that  $\nabla \hat{y}$  vanishes a.e. in  $\{\hat{y} = \bar{t}\}$  in order to obtain the last identity. This yields the first claim.  $\square$

**Lemma 4.4.** *There exist an  $h_3 \in (0, h_2]$  and a constant  $L_\rho > 0$  such that for all  $u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$  and  $h \in (0, h_3)$ , there hold*

$$(4.11) \quad \|Z_{y_h,y}\|_{L^2(\Omega)} \leq L_\rho h + \|V(y, \|y_h - y\|_{L^\infty(\Omega)})\|_{L^2(\Omega)}$$

and

$$(4.12) \quad \|Z_{y,\bar{y}}\|_{L^2(\Omega)} \leq L_\rho \|u - \bar{u}\|_{L^2(\Omega)} + \|V(\bar{y}, \|y - \bar{y}\|_{L^\infty(\Omega)})\|_{L^2(\Omega)}$$

with  $y := S(u)$  and  $y_h := S_h(u)$ .

*Proof.* By [Theorem 2.2](#) (also, see, [Theorems 3.1](#) and [3.5](#) in [\[22\]](#)), there exists a constant  $M_{1,\rho}$  such that

$$(4.13) \quad \|S(u)\|_{W^{1,\infty}(\Omega)} \leq M_{1,\rho} \quad \text{for all } u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}.$$

Setting  $h_3 := \min\{h_2, \delta 2^{-1} C_\infty^{-1}\}$  and exploiting [\(4.9\)](#) shows that  $\|y - y_h\|_{C(\bar{\Omega})} < \delta$  for any  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$  and  $h \in (0, h_3)$ . From the definition of  $Z_{y_h, y}$  in [\(3.6\)](#) and [Lemma 3.4](#), we arrive at

$$(4.14) \quad Z_{y_h, y} = Z_{y_h, y}^{(1)} + Z_{y_h, y}^{(2)} + Z_{y_h, y}^{(3)} + Z_{y_h, y}^{(4)}.$$

By [\(3.8\)](#), [\(4.13\)](#), and [Theorem 3.1](#), we have

$$(4.15) \quad \|Z_{y_h, y}^{(k)}\|_{L^2(\Omega)} \leq L_\rho h \quad \text{for } k = 1, 2, 4.$$

On the other hand, we have

$$(4.16) \quad \begin{cases} \Omega_{y_h, y}^2 = \{y \in (\bar{t}, \bar{t} + \delta), y_h \in (\bar{t} - \delta, \bar{t})\} \subset \{0 < y - \bar{t} \leq \|y_h - y\|_{L^\infty(\Omega)}\}, \\ \Omega_{y_h, y}^3 = \{y \in (\bar{t} - \delta, \bar{t}), y_h \in [\bar{t}, \bar{t} + \delta)\} \subset \{0 < \bar{t} - y \leq \|y_h - y\|_{L^\infty(\Omega)}\}, \end{cases}$$

which together with the definitions of  $Z_{y_h, y}^{(3)}$  in [Lemma 3.4](#) and of  $V$  in [\(4.10\)](#), and [\(2.14\)](#), show that

$$(4.17) \quad |Z_{y_h, y}^{(3)}| \leq V(y, \|y_h - y\|_{L^\infty(\Omega)})$$

a.e. in  $\Omega$ . Combining this with [\(4.14\)](#) and [\(4.15\)](#), we obtain [\(4.11\)](#).

For [\(4.12\)](#), we first see that

$$\|Z_{y, \bar{y}}^{(k)}\|_{L^2(\Omega)} \leq C \|y - \bar{y}\|_{H_0^1(\Omega)} \leq C \|u - \bar{u}\|_{L^2(\Omega)} \quad \text{for } k = 1, 2, 4,$$

which is similar to [\(4.15\)](#) and is derived by using [\(3.8\)](#), [\(4.13\)](#), [Theorem 2.2](#), and the continuity of the mapping  $S : L^2(\Omega) \hookrightarrow W^{-1, \tilde{p}}(\Omega) \rightarrow W_0^{1, \tilde{p}}(\Omega) \hookrightarrow H_0^1(\Omega)$ . Finally, similar to [\(4.17\)](#), one has

$$|Z_{y, \bar{y}}^{(3)}| \leq V(y, \|y - \bar{y}\|_{L^\infty(\Omega)})$$

a.e. in  $\Omega$ . We thus obtain [\(4.12\)](#).  $\square$

From now on, let us fix constants  $\gamma$ ,  $p_0$ , and  $p_1$  satisfying

$$(4.18) \quad 1 - \frac{2}{p_0} \leq \gamma < \min\left\{\frac{1}{p_0}, 1 - \frac{2}{p_1}\right\} < \frac{1}{2}, \quad p_0 > 2 = N, \quad \text{and} \quad p_1 \in (2, p_*),$$

where  $p_*$  is given in [Theorem 2.2](#).

In order to derive an  $L^2$ -error estimate of  $\tilde{\varphi} - \varphi_h$ , we cannot directly employ a duality argument based on the Aubin–Nitsche trick since the linearized state equation [\(2.4\)](#) admits solutions belonging to  $W^{1, p}(\Omega)$  only due to the nondifferentiability of the function  $a$ ; see [Remark 2.4](#). To overcome this difficulty, we now consider the following *adjusted* linearized state equation:

$$(4.19) \quad -\operatorname{div}[(b + a(y_u))\nabla \tilde{z}_{u, v} + \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'(\bar{y}) \tilde{z}_{u, v} \nabla \bar{y}] = v \text{ in } \Omega, \quad \tilde{z}_{u, v} = 0 \text{ on } \partial\Omega$$

for all  $u \in \mathcal{U}_{ad}$  and  $v$  belonging to  $W^{-1+\gamma, 2}(\Omega)$ , the dual space of a Sobolev space of fractional order. Compared to [\(2.4\)](#), we have here partly linearized the state equation [\(2.3\)](#) at the optimal state  $\bar{y}$  by replacing the vector-valued function  $\mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \nabla y_u$  in [\(2.4\)](#) by  $\mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'(\bar{y}) \nabla \bar{y}$ . We can therefore exploit assumptions imposed on  $\bar{y}$  to derive the necessary regularity of solutions  $\tilde{z}_{u, v}$  to [\(4.19\)](#). Indeed, as we will see later in [Proposition B.1](#) in [Appendix B](#), the finiteness of  $\Sigma(\bar{y})$  implies that

$$\mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'(\bar{y}) \nabla \bar{y} \in (W^{\gamma, p_0}(\Omega))^2 = (W_0^{\gamma, p_0}(\Omega))^2,$$

where the constants  $\gamma$  and  $p_0$  are fixed and satisfy [\(4.18\)](#). From this and the  $W^{1+\gamma, 2}$ -regularity of solutions to [\(4.19\)](#), we can then show that  $\tilde{z}_{u, v} \in W^{1+\gamma, 2}(\Omega)$  whenever  $v \in W^{-1+\gamma, 2}(\Omega)$ . This regularity of solutions to [\(4.19\)](#) will play an important role in establishing a priori error estimates for the discretization of the adjoint state equation [\(2.5\)](#).

**Theorem 4.5.** *If  $\Sigma(\bar{y}) < \infty$ , then there exist constants  $\bar{h} := \bar{h}(\bar{u}) \in (0, h_3)$ ,  $\bar{\rho} := \bar{\rho}(\bar{u}) \leq \min\{\rho, \hat{\rho}, \rho_{\bar{p}}\}$  (with  $\hat{\rho}$  and  $\rho_{\bar{p}}$  being the constants in [Proposition B.3](#) associated with  $p := \bar{p}$ ), and  $C_{\bar{u}} > 0$  such that*

$$\|\varphi - \varphi_h\|_{L^2(\Omega)} \leq C_{\bar{u}} \varepsilon_h^u \|v\|_{L^2(\Omega)} \quad \text{and} \quad \|\varphi - \varphi_h\|_{H_0^1(\Omega)} \leq C_{\bar{u}} h \|v\|_{L^2(\Omega)}$$

for all  $h \in (0, \bar{h})$ ,  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\rho}) \cap \mathcal{U}_{ad}$ , and  $v \in L^2(\Omega)$ , where

$$(4.20) \quad \varepsilon_h^u := h^{1+\gamma} + h \|Z_{S(u), \bar{y}}\|_{L^2(\Omega)} + h \|V(S(u), \|S_h(u) - S(u)\|_{L^\infty(\Omega)})\|_{L^2(\Omega)},$$

and  $\varphi$  and  $\varphi_h$  are the unique solutions to (2.5) and (4.1), respectively.

*Proof.* Let  $\tilde{\varphi}$  be the solution of (4.2). To simplify the notation, set  $y_u := S(u)$  and  $y_h := S_h(u)$  for any  $h \in (0, h_3)$  and  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \min\{\rho, \hat{\rho}, \rho_{\bar{p}}\}) \cap \mathcal{U}_{ad}$ . We divide the proof into three steps.

*Step 1: Existence of a constant  $C_{1, h_3, \rho}$  such that*

$$(4.21) \quad \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)} \leq C_{1, h_3, \rho} (h^\gamma + \|Z_{y_u, \bar{y}}\|_{L^2(\Omega)} + \|V(y_u, \|y_h - y_u\|_{L^\infty(\Omega)})\|_{L^2(\Omega)}) \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)}$$

for all  $h \in (0, h_3)$ ,  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \min\{\rho, \hat{\rho}, \rho_{\bar{p}}\}) \cap \mathcal{U}_{ad}$ , and  $v \in L^2(\Omega)$ .

To prove (4.21), first let  $\hat{z}$  be the unique solution to the adjusted linearized state equation (4.19) corresponding to  $v := \tilde{\varphi} - \varphi_h$ . Since  $\tilde{\varphi} - \varphi_h \in H_0^1(\Omega)$ , one has  $\tilde{\varphi} - \varphi_h \in W^{-1+\gamma, 2}(\Omega)$  due to the embeddings  $H_0^1(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow W^{-1+\gamma, 2}(\Omega)$ . From [Proposition B.3](#) and the finiteness of  $\Sigma(\bar{y})$ , we have that  $\hat{z} \in W^{1+\gamma, 2}(\Omega)$  and that

$$(4.22) \quad \|\hat{z}\|_{W^{1+\gamma, 2}(\Omega)} \leq C \|\tilde{\varphi} - \varphi_h\|_{W^{-1+\gamma, 2}(\Omega)} \leq C \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}.$$

Moreover, we also have that

$$(4.23) \quad \hat{z} = S'(u)(\tilde{\varphi} - \varphi_h - \operatorname{div}(Z_{y_u, \bar{y}} \hat{z}));$$

see the identity (B.11) in the proof of [Proposition B.3](#) for  $v := \tilde{\varphi} - \varphi_h$ . Testing (2.4) for  $v := \tilde{\varphi} - \varphi_h - \operatorname{div}(Z_{y_u, \bar{y}} \hat{z})$  by  $\tilde{\varphi} - \varphi_h$  and using (4.23) thus yields

$$(4.24) \quad \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^2 + \int_{\Omega} Z_{y_u, \bar{y}} \cdot \nabla(\tilde{\varphi} - \varphi_h) \hat{z} \, dx \\ = \int_{\Omega} (b + a(y_u)) \nabla \hat{z} \cdot \nabla(\tilde{\varphi} - \varphi_h) + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \hat{z} \nabla y_u \cdot \nabla(\tilde{\varphi} - \varphi_h) \, dx.$$

Furthermore, applying assertion (i) in [Proposition B.3](#) for  $p := \bar{p}$  and using the embedding  $L^2(\Omega) \hookrightarrow W^{-1, \bar{p}}(\Omega)$  shows that

$$(4.25) \quad \|\hat{z}\|_{W_0^{1, \bar{p}}(\Omega)} \leq C_{\rho} \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)} \quad \text{for all } h \in (0, h_3), u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\rho}) \cap \mathcal{U}_{ad}, v \in L^2(\Omega).$$

Consider for any  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \min\{\rho, \hat{\rho}, \rho_{\bar{p}}\}) \cap \mathcal{U}_{ad}$  and  $h \in (0, h_3)$  the bilinear operators  $B_u : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$  and  $B_{u, h} : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$  defined via

$$B_u(z, w) := \int_{\Omega} [(b + a(y_u)) \nabla z + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) z \nabla y_u] \cdot \nabla w \, dx, \\ B_{u, h}(z, w) := \int_{\Omega} [(b + a(y_h)) \nabla z + \mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h) z \nabla y_h] \cdot \nabla w \, dx.$$

From this and (4.24), we obtain for any  $w_h \in V_h$  that

$$\begin{aligned} \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^2 + \int_{\Omega} Z_{y_u, \tilde{y}} \cdot \nabla(\tilde{\varphi} - \varphi_h) \hat{z} \, dx &= B_u(\hat{z}, \tilde{\varphi} - \varphi_h) = B_u(\hat{z} - w_h, \tilde{\varphi} - \varphi_h) + B_u(w_h, \tilde{\varphi} - \varphi_h) \\ &= B_u(\hat{z} - w_h, \tilde{\varphi} - \varphi_h) + [B_u(w_h, \tilde{\varphi} - \varphi_h) - B_{u,h}(w_h, \tilde{\varphi} - \varphi_h)], \end{aligned}$$

where we have used the fact that  $B_{u,h}(w_h, \tilde{\varphi} - \varphi_h) = 0$  which follows from combining (4.1) with (4.2). We now estimate the first integral in the left-hand side and two summands in the right-hand side of the above identity. From Hölder's inequality, there holds

$$\begin{aligned} (4.26) \quad \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^2 - \|Z_{y_u, \tilde{y}}\|_{L^2(\Omega)} \|\hat{z}\|_{L^\infty(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \\ \leq \|b + a(y_u)\|_{L^\infty(\Omega)} \|\hat{z} - w_h\|_{H_0^1(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \\ + \|\mathbb{1}_{\{y_u \neq \tilde{t}\}} a'(y_u) \nabla y_u\|_{L^\infty(\Omega)} \|\hat{z} - w_h\|_{L^2(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \\ + \|a(y_u) - a(y_h)\|_{L^4(\Omega)} \|w_h\|_{W_0^{1,4}(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \\ + \|Z_{y_h, y_u}\|_{L^2(\Omega)} \|w_h\|_{L^\infty(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \end{aligned}$$

for all  $w_h \in V_h$ . Moreover, we have from Assumption (A3), the continuous embedding  $H^{N/4}(\Omega) \hookrightarrow L^4(\Omega)$  with  $N = 2$  (see, e.g. [34], Thm. 1.4.4.1), interpolation theory [7], Thm. 14.2.7, and (3.2) that

$$\begin{aligned} (4.27) \quad \|a(y_u) - a(y_h)\|_{L^4(\Omega)} &\leq C \|y_u - y_h\|_{L^4(\Omega)} \leq C \|y_u - y_h\|_{H^{N/4}(\Omega)} \\ &\leq C \|y_u - y_h\|_{L^2(\Omega)}^{1-\frac{2}{4}} \|y_u - y_h\|_{H_0^1(\Omega)}^{\frac{2}{4}} \leq Ch^{\frac{3}{2}}. \end{aligned}$$

We then deduce from this, (4.26), (4.13), the assumptions on  $b$  and  $a$ , and (4.11) that

$$\begin{aligned} (4.28) \quad \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^2 &\leq C_\rho \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \left[ \|\hat{z} - w_h\|_{H_0^1(\Omega)} + h^{\frac{3}{2}} \|w_h\|_{W_0^{1,4}(\Omega)} \right] \\ &\quad + (h + \|V(y_u, \|y_h - y_u\|_{L^\infty(\Omega)})\|_{L^2(\Omega)}) \|w_h\|_{L^\infty(\Omega)} + \|Z_{y_u, \tilde{y}}\|_{L^2(\Omega)} \|\hat{z}\|_{L^\infty(\Omega)} \end{aligned}$$

for all  $w_h \in V_h$ . Moreover, from standard interpolation error estimates (see, e.g. Theorem 4.4.20 in [7]) and applying the estimates (4.4.21) for  $(p, s, m) := (4, 1, 1)$  and (4.4.22) for  $(p, s, m, n) := (\tilde{p}, 0, 1, 2)$ , we have that

$$(4.29) \quad \begin{cases} \|\hat{z} - \Pi_h \hat{z}\|_{W_0^{1,4}(\Omega)} \leq C_1 \|\hat{z}\|_{W_0^{1,4}(\Omega)} \leq C_1 \|\hat{z}\|_{W_0^{1,\tilde{p}}(\Omega)}, \\ \|\hat{z} - \Pi_h \hat{z}\|_{L^\infty(\Omega)} \leq C_1 h^{1-\frac{2}{\tilde{p}}} \|\hat{z}\|_{W_0^{1,\tilde{p}}(\Omega)}, \end{cases}$$

where we have used the fact that  $\tilde{p} \geq 4$  to derive the first line in (4.29). On the other hand, from Theorem 6.1 in [29] and arguments similar to the ones in Example 3 in [29], we also obtain that

$$(4.30) \quad \|\hat{z} - \Pi_h \hat{z}\|_{H_0^1(\Omega)} \leq C_2 h^\gamma \|\hat{z}\|_{W^{1+\gamma,2}(\Omega)};$$

see also the error estimates for the interpolation operator  $\Pi_h : W^{1+\gamma,2}(\Omega) \rightarrow V_h$  in the proof of Lemma 3.1 in [26]. Combining (4.29) with (4.30), the triangle inequality, and the embedding  $W_0^{1,\tilde{p}}(\Omega) \hookrightarrow L^\infty(\Omega) \cap W_0^{1,4}(\Omega)$  (due to  $\tilde{p} \geq 4$ ) as well as estimates (4.22) and (4.25) gives

$$\|\hat{z} - \Pi_h \hat{z}\|_{H_0^1(\Omega)} \leq C_3 h^\gamma \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}, \quad \|\hat{z}\|_{L^\infty(\Omega)} + \|\Pi_h \hat{z}\|_{L^\infty(\Omega)} + \|\Pi_h \hat{z}\|_{W_0^{1,4}(\Omega)} \leq C_4 \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}.$$

By choosing  $w_h := \Pi_h \hat{z}$  in (4.28) and using these above estimates and the fact that  $\gamma < \frac{1}{2}$ , we obtain (4.21).

Step 2: Existence of a constant  $C_{2,h_3,\rho}$  such that

$$(4.31) \quad \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \leq C_{2,h_3,\rho} (h\|v\|_{L^2(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)})$$

for all  $h \in (0, h_3)$ ,  $u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$ , and  $v \in L^2(\Omega)$ .

To show this, we first consider for any  $u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$  and  $h \in (0, h_3)$ , the bilinear mapping  $S_{u,h} : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$  defined by

$$S_{u,h}(\omega, \psi) := \int_{\Omega} (b + a(y_h)) \nabla \omega \cdot \nabla \psi \, dx.$$

From Assumptions (A2) and (A3), we obtain

$$(4.32) \quad \underline{b} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)}^2 \leq S_{u,h}(\tilde{\varphi} - \varphi_h, \tilde{\varphi} - \varphi_h) = S_{u,h}(\tilde{\varphi} - \varphi_h, \tilde{\varphi} - \Pi_h \tilde{\varphi}) + S_{u,h}(\tilde{\varphi} - \varphi_h, \Pi_h \tilde{\varphi} - \varphi_h).$$

Moreover, the Cauchy–Schwarz inequality, the uniform boundedness of  $\{y_h\}$  on  $C(\overline{\Omega})$ , and Assumptions (A2) and (A3) yield that

$$(4.33) \quad \begin{aligned} S_{u,h}(\tilde{\varphi} - \varphi_h, \tilde{\varphi} - \Pi_h \tilde{\varphi}) &\leq C_{\rho} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \|\tilde{\varphi} - \Pi_h \tilde{\varphi}\|_{H_0^1(\Omega)} \\ &\leq C_{\rho} h \|\tilde{\varphi}\|_{H^2(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \leq C_{\rho} h \|v\|_{L^2(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)}, \end{aligned}$$

where we have exploited the interpolation error [20] and (4.4) in order to obtain the last two estimates. Now using (4.1) and (4.2), we deduce from Hölder's inequality that

$$\begin{aligned} S_{u,h}(\tilde{\varphi} - \varphi_h, \Pi_h \tilde{\varphi} - \varphi_h) &= - \int_{\Omega} \mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h) \nabla y_h \cdot \nabla (\tilde{\varphi} - \varphi_h) (\Pi_h \tilde{\varphi} - \varphi_h) \, dx \\ &\leq \|\mathbb{1}_{\{y_h \neq \bar{t}\}} a'(y_h)\|_{L^\infty(\Omega)} \|\nabla y_h\|_{L^4(\Omega)} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \|\Pi_h \tilde{\varphi} - \varphi_h\|_{L^4(\Omega)}. \end{aligned}$$

Combing this with the uniform boundedness in  $C(\overline{\Omega})$  of  $\{y_h\}$  and the embedding  $W_0^{1,\tilde{p}}(\Omega) \hookrightarrow W_0^{1,4}(\Omega)$ , we obtain that

$$S_{u,h}(\tilde{\varphi} - \varphi_h, \Pi_h \tilde{\varphi} - \varphi_h) \leq C_{\rho} (\|\bar{y}\|_{W_0^{1,\tilde{p}}(\Omega)} + \kappa_{\rho}) \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \|\Pi_h \tilde{\varphi} - \varphi_h\|_{L^4(\Omega)}.$$

The combination of a triangle inequality and the embedding  $H_0^1(\Omega) \hookrightarrow L^4(\Omega)$  with Theorem 3.1.6 in [20] further implies that

$$\begin{aligned} \|\Pi_h \tilde{\varphi} - \varphi_h\|_{L^4(\Omega)} &\leq \|\Pi_h \tilde{\varphi} - \tilde{\varphi}\|_{L^4(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^4(\Omega)} \leq C \|\Pi_h \tilde{\varphi} - \tilde{\varphi}\|_{H_0^1(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^4(\Omega)} \\ &\leq Ch \|\tilde{\varphi}\|_{H^2(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^4(\Omega)} \\ &\leq Ch \|v\|_{L^2(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^4(\Omega)}, \end{aligned}$$

where we have used (4.4) to obtain the last inequality. Similar to (4.27), we find that

$$\|\tilde{\varphi} - \varphi_h\|_{L^4(\Omega)} \leq C \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^{1-\frac{2}{4}} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)}^{\frac{2}{4}}.$$

We then have

$$S_{u,h}(\tilde{\varphi} - \varphi_h, \Pi_h \tilde{\varphi} - \varphi_h) \leq C_{\rho} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \left( h \|v\|_{L^2(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^{\frac{1}{2}} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)}^{\frac{1}{2}} \right),$$

which, together with (4.32) and (4.33), yields

$$\|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \leq C_{\rho} \left( h \|v\|_{L^2(\Omega)} + \|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)}^{\frac{1}{2}} \|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)}^{\frac{1}{2}} \right).$$

Applying the Cauchy–Schwarz inequality then gives (4.31).

*Step 3: Existence of constants  $\bar{h} \in (0, h_3)$  and  $\bar{\rho} := \bar{\rho}(\bar{u}) \leq \min\{\rho, \hat{\rho}, \rho_{\bar{\rho}}\}$ .*

To show this, we first obtain from the definition of  $\Sigma(\bar{y})$  in (2.13) and of  $\Sigma_r$  in (4.10) the existence of a  $r_* > 0$  such that  $\|\Sigma_r(\bar{y})\|_{L^1(\Omega)} \leq \Sigma(\bar{y}) + 1$  for all  $r \in (0, r_*)$ . This together with (ii) in Proposition 4.3 yields

$$(4.34) \quad \|V(\bar{y}, r)\|_{L^2(\Omega)} \leq C_{\bar{y}} r^{1/2} (\Sigma(\bar{y}) + 1)^{1/2} \quad \text{for all } r \in (0, r_*).$$

Moreover, thanks to Theorem 2.2 and the embeddings  $L^2(\Omega) \hookrightarrow W^{-1,p}(\Omega)$  and  $W_0^{1,p}(\Omega) \hookrightarrow C(\bar{\Omega}) \cap H_0^1(\Omega)$  for some  $p > 2$ , one has

$$\|y_u - \bar{y}\|_{L^\infty(\Omega)} + \|y_u - \bar{y}\|_{H_0^1(\Omega)} \leq C_\rho \|u - \bar{u}\|_{L^2(\Omega)}$$

for all  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$  and some constant  $C_\rho$ . Now Proposition 4.3 (i), (4.9), and the monotonic growth of  $V(y, \cdot)$  imply that

$$(4.35) \quad \begin{aligned} \|V(y_u, \|y_h - y_u\|_{L^\infty(\Omega)})\|_{L^2(\Omega)} &\leq \|V(\bar{y}, \|y_h - y_u\|_{L^\infty(\Omega)} + \|y_u - \bar{y}\|_{L^\infty(\Omega)})\|_{L^2(\Omega)} + C_{\bar{y},\rho} \|y_u - \bar{y}\|_{H_0^1(\Omega)} \\ &\leq \|V(\bar{y}, C_\infty h + C_\rho \|u - \bar{u}\|_{L^2(\Omega)})\|_{L^2(\Omega)} + C_{\bar{y},\rho} C_\rho \|u - \bar{u}\|_{L^2(\Omega)} \end{aligned}$$

for some constant  $C_{\bar{y},\rho}$  and for all  $h \in (0, h_3)$  and  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$ . Besides, from (4.12) and the monotonic growth of  $V(\bar{y}, \cdot)$ , there holds

$$\begin{aligned} \|Z_{y_u, \bar{y}}\|_{L^2(\Omega)} &\leq L_\rho \|u - \bar{u}\|_{L^2(\Omega)} + \|V(\bar{y}, \|y_u - \bar{y}\|_{L^\infty(\Omega)})\|_{L^2(\Omega)} \\ &\leq L_\rho \|u - \bar{u}\|_{L^2(\Omega)} + \|V(\bar{y}, C_\rho \|u - \bar{u}\|_{L^2(\Omega)})\|_{L^2(\Omega)} \end{aligned}$$

for all  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$ . We then have

$$\|Z_{y_u, \bar{y}}\|_{L^2(\Omega)} + \|V(y_u, \|y_h - y_u\|_{L^\infty(\Omega)})\|_{L^2(\Omega)} \leq C_{1,\rho} \|u - \bar{u}\|_{L^2(\Omega)} + 2\|V(\bar{y}, C_\infty h + C_\rho \|u - \bar{u}\|_{L^2(\Omega)})\|_{L^2(\Omega)}$$

for all  $h \in (0, h_3)$  and  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \cap \mathcal{U}_{ad}$  with  $C_{1,\rho} := (C_{\bar{y},\rho} C_\rho + L_\rho)$ .

Next, fix  $\bar{h} \in (0, h_3)$  and  $\bar{\rho} \leq \min\{\rho, \hat{\rho}, \rho_{\bar{\rho}}\}$  such that

$$\bar{h}^\gamma + 2C_{\bar{y}} [C_\infty \bar{h} + C_\rho \bar{\rho}]^{\frac{1}{2}} (\Sigma(\bar{y}) + 1)^{\frac{1}{2}} + C_{1,\rho} \bar{\rho} \leq \frac{1}{2C_{1,h_3,\rho} C_{2,h_3,\rho}} \quad \text{and} \quad C_\infty \bar{h} + C_\rho \bar{\rho} < r_*,$$

where  $C_{1,h_3,\rho}$  and  $C_{2,h_3,\rho}$  are defined in (4.21) and (4.31), respectively. From this, (4.34), and (4.35), we conclude that

$$\bar{h}^\gamma + \|Z_{y_u, \bar{y}}\|_{L^2(\Omega)} + \|V(y_u, \|y_h - y_u\|_{L^\infty(\Omega)})\|_{L^2(\Omega)} \leq \frac{1}{2C_{1,h_3,\rho} C_{2,h_3,\rho}}$$

for all  $h \in (0, \bar{h})$ ,  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\rho}) \cap \mathcal{U}_{ad}$ .

The combination of this with (4.21) and (4.31) yields  $\|\tilde{\varphi} - \varphi_h\|_{H_0^1(\Omega)} \leq 2C_{2,h_3,\rho} h \|v\|_{L^2(\Omega)}$ , and together with (4.21), we obtain  $\|\tilde{\varphi} - \varphi_h\|_{L^2(\Omega)} \leq C\varepsilon_h^u \|v\|_{L^2(\Omega)}$ . Combining the last two estimates with Lemma 4.2 and the triangle inequality, we arrive at the desired conclusion.  $\square$

## 5 DISCRETIZATION OF THE CONTROL PROBLEM

In this section, we discretize the control problem (P), show convergence of the discretizations, and derive error estimates of the discrete optimal solutions. In the following, we will consider three different discretizations of the control:

(i) *variational discretization*:  $\mathcal{U}_h = L^\infty(\Omega)$  (see, e.g., [35]);

(ii) *piecewise constant discretization*:

$$\mathcal{U}_h = \mathcal{U}_h^0 := \{u \in L^\infty(\Omega) \mid u|_T \in \mathbb{R} \text{ for all } T \in \mathcal{T}_h\};$$

(iii) *continuous piecewise linear discretization*:

$$\mathcal{U}_h = \mathcal{U}_h^1 := \{u \in L^\infty(\Omega) \mid u|_T \in \mathcal{P}_1 \text{ for all } T \in \mathcal{T}_h\}.$$

Unless specified, any claim for  $\mathcal{U}_h$  should be understood to hold for all three cases. For any  $h > 0$ , we now set  $\mathcal{U}_{ad,h} := \mathcal{U}_{ad} \cap \mathcal{U}_h$ . If  $\mathcal{U}_h = \mathcal{U}_h^0$ , then by  $\mathcal{I}_h$  we denote the linear projection from  $L^2(\Omega)$  onto  $\mathcal{U}_h^0$ . If  $\mathcal{U}_h = \mathcal{U}_h^1$ , then  $\mathcal{I}_h : L^2(\Omega) \rightarrow \mathcal{U}_h^1$  denotes the Carstensen quasi-interpolation operator [8]. In both situations, we have  $\mathcal{I}_h u \rightarrow u$  strongly in  $L^2(\Omega)$  as  $h \rightarrow 0$  for all  $u \in L^2(\Omega)$  and  $\mathcal{I}_h u \in \mathcal{U}_{ad,h}$  for all  $u \in \mathcal{U}_{ad}$ ; see, e.g. [27].

For any  $h > 0$ , we define the discretized optimal control problem

$$(P_h) \quad \min\{J(y_h, u_h) : u_h \in \mathcal{U}_{ad,h} \text{ and } y_h \in V_h \text{ satisfies (3.1) for } u := u_h\}$$

with

$$J(y, u) := \int_{\Omega} L(x, y(x)) \, dx + \frac{\nu}{2} \|u\|_{L^2(\Omega)}^2.$$

Note that the discrete operator  $S_h$  defined in (3.4) does not appear in  $(P_h)$ , since this operator is well-defined only locally by [Theorem 3.2](#).

## 5.1 CONVERGENCE OF DISCRETE MINIMIZERS

We first have the convergence of minimizers of  $(P_h)$ .

**Theorem 5.1** (cf. [Theorem 4.1](#) in [13]). *Assume that  $a$  satisfies the growth condition*

$$(5.1) \quad |a(t)| \leq C_0 + C_1 |t|^m \quad \text{for all } t \in \mathbb{R}$$

for some positive constants  $C_0, C_1$  and  $m \geq 1$ . Then there exists an  $\tilde{h}_0 > 0$  such that  $(P_h)$  admits at least one global minimizer  $(\bar{y}_h, \bar{u}_h)$  for all  $0 < h < \tilde{h}_0$ . Moreover, if  $\{(\bar{y}_h, \bar{u}_h)\}_{0 < h < \tilde{h}_0}$  is a sequence of solutions to  $(P_h)$ , then there exists a subsequence that converges strongly in  $H_0^1(\Omega) \times L^2(\Omega)$  to some  $(\bar{y}, \bar{u})$  as  $h \rightarrow 0$ , where  $\bar{u}$  is a global solution to (P).

*Proof.* The existence of discrete solutions to  $(P_h)$  is proven similarly to Claim 1 in the proof of [Theorem 4.1](#) in [13]. Moreover, there exists a constant  $\tilde{h}_0 > 0$  such that  $(P_h)$  admits at least one minimizer  $(\bar{y}_h, \bar{u}_h)$  for all  $0 < h < \tilde{h}_0$ . The remainder of this proof is now divided into three steps as follows:

- *Claim 1: Weak convergence of  $\{(\bar{y}_h, \bar{u}_h)\}$  to  $(\bar{y}, \bar{u})$  satisfying (2.3).* Indeed, the boundedness of  $\{\bar{u}_h\}$  in  $L^\infty(\Omega)$  and thus in  $L^2(\Omega)$  is due to the  $L^\infty$ -boundedness of  $\mathcal{U}_{ad}$ . Since  $(\bar{y}_h, \bar{u}_h)$  satisfies (3.1), there holds

$$(5.2) \quad \int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla v_h \, dx = \int_{\Omega} \bar{u}_h v_h \, dx \quad \text{for all } v_h \in V_h.$$

In particular, one has

$$\int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx = \int_{\Omega} \bar{y}_h \bar{u}_h \, dx.$$

Combining this with [Assumption \(A3\)](#) and [Assumption \(A2\)](#), we deduce from the Cauchy–Schwarz and Poincaré inequalities the boundedness of  $\{\bar{y}_h\}$  in  $H_0^1(\Omega)$ . From this and the compact embedding

$H_0^1(\Omega) \Subset L^p(\Omega)$  for any  $p \geq 1$ , we can take a subsequence, denoted in the same way, of  $\{(\bar{y}_h, \bar{u}_h)\}_{0 < h < \tilde{h}_0}$  that satisfies

$$(5.3) \quad (\bar{y}_h, \bar{u}_h) \rightharpoonup (\bar{y}, \bar{u}) \text{ in } H_0^1(\Omega) \times L^2(\Omega), \quad \bar{y}_h(x) \rightarrow \bar{y}(x) \text{ for a.e. } x \in \Omega, \text{ and } \bar{y}_h \rightarrow \bar{y} \text{ in } L^p(\Omega)$$

for some  $(\bar{y}, \bar{u}) \in H_0^1(\Omega) \times \mathcal{U}_{ad}$  and for any  $p \geq 1$ , for instance,  $p := 4m$  with constant  $m$  defined in (5.1). Thanks to the growth condition (5.1), we conclude from the last limit and the generalized Lebesgue Dominated Convergence Theorem that

$$(5.4) \quad a(\bar{y}_h) \rightarrow a(\bar{y}) \quad \text{strongly in } L^4(\Omega).$$

Now take any  $v \in H_0^1(\Omega) \cap W^{2,4}(\Omega)$  and choose  $v_h := \Pi_h v \in V_h$ . Then  $v_h \rightarrow v$  in  $W^{1,4}(\Omega)$  as  $h \rightarrow 0$ ; see, e.g. [20]. Letting  $h \rightarrow 0$  in the equation (5.2) and exploiting the first limit in (5.3) as well as the limit in (5.4) then yields

$$\int_{\Omega} (b + a(\bar{y})) \nabla \bar{y} \cdot \nabla v \, dx = \int_{\Omega} \bar{u} v \, dx \quad \text{for all } v \in H_0^1(\Omega) \cap H^2(\Omega),$$

which, together with the density of  $H_0^1(\Omega) \cap W^{2,4}(\Omega)$  in  $H_0^1(\Omega)$ , implies that  $\bar{y} = S(\bar{u})$ .

• *Claim 2: Optimality of  $\bar{u}$ .* Let us show that  $\bar{u}$  is a global solution of (P). For that purpose, we first observe from the inclusion  $\mathcal{U}_{ad,h} \subset \mathcal{U}_{ad}$  and from the first limit in (5.3) that  $\bar{u} \in \mathcal{U}_{ad}$ . Take  $u \in \mathcal{U}_{ad}$  arbitrarily and choose  $u_h := u$  if  $\mathcal{U}_h = L^\infty(\Omega)$  and  $u_h := \mathcal{I}_h u$  if  $\mathcal{U}_h = \mathcal{U}_h^i$  with  $i = 0, 1$ . One has  $u_h \in \mathcal{U}_{ad,h}$  and  $u_h \rightarrow u$  strongly in  $L^2(\Omega)$  as  $h \rightarrow 0$ . In view of Theorem 3.1, for  $h$  small enough there exists at least one solution  $y_h(u_h)$  of (3.1) such that  $y_h(u_h) \rightarrow y_u := S(u)$  strongly in  $H_0^1(\Omega) \cap C(\bar{\Omega})$ . From this and the optimality of  $(\bar{y}_h, \bar{u}_h)$ , we have

$$(5.5) \quad \liminf_{h \rightarrow 0} J(\bar{y}_h, \bar{u}_h) \leq \liminf_{h \rightarrow 0} J(y_h(u_h), u_h) \leq \limsup_{h \rightarrow 0} J(y_h(u_h), u_h) = J(y_u, u) = j(u).$$

On the other hand, it follows from the limits in (5.3) and the weak lower semicontinuity of the  $L^2$ -norm that

$$(5.6) \quad \liminf_{h \rightarrow 0} J(\bar{y}_h, \bar{u}_h) \geq J(\bar{y}, \bar{u}) = j(\bar{u}),$$

which together with (5.5) gives  $j(\bar{u}) \leq j(u)$ . Since  $u \in \mathcal{U}_{ad}$  was arbitrary,  $\bar{u}$  is a global optimal solution to (P).

• *Claim 3: Strong convergence in  $H_0^1(\Omega) \times L^2(\Omega)$ .* In fact, by plugging  $u := \bar{u}$  into (5.5) and (5.6) and using the limits for  $\bar{y}_h$  in (5.3), we can conclude that  $\|\bar{u}_h\|_{L^2(\Omega)}^2 \rightarrow \|\bar{u}\|_{L^2(\Omega)}^2$ . Combining this with the limit for  $\bar{u}_h$  in (5.3) yields

$$(5.7) \quad \bar{u}_h \rightarrow \bar{u} \quad \text{strongly in } L^2(\Omega).$$

It remains to prove the strong convergence of  $\{\bar{y}_h\}$  in  $H_0^1(\Omega)$ . To this end, by the weak lower semicontinuity of the functional  $H_0^1(\Omega) \ni y \mapsto \int_{\Omega} (b(x) + a(y(x))) \nabla y(x) \cdot \nabla y(x) \, dx \in \mathbb{R} \cup \{\infty\}$ ; see, e.g. Theorem 1.3 in [25], we deduce from the first limit in (5.3) that

$$(5.8) \quad \int_{\Omega} (b + a(\bar{y})) \nabla \bar{y} \cdot \nabla \bar{y} \, dx \leq \liminf_{h \rightarrow 0} \int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx.$$

Moreover, (5.2) and (5.7) imply that

$$\begin{aligned} \liminf_{h \rightarrow 0} \int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx &\leq \limsup_{h \rightarrow 0} \int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx \\ &= \limsup_{h \rightarrow 0} \int_{\Omega} \bar{u}_h \bar{y}_h \, dx = \int_{\Omega} \bar{y} \bar{u} \, dx \\ &= \int_{\Omega} (b + a(\bar{y})) \nabla \bar{y} \cdot \nabla \bar{y} \, dx. \end{aligned}$$

Combining this with (5.8), one has

$$(5.9) \quad \lim_{h \rightarrow 0} \int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx = \int_{\Omega} (b + a(\bar{y})) \nabla \bar{y} \cdot \nabla \bar{y} \, dx.$$

Moreover, thanks to Assumption (A2) and by the nonnegativity of  $a$ , there hold  $b(x) - \frac{b}{2} \geq \frac{b}{2} > 0$  and  $\frac{b}{2} + a(\bar{y}_h(x)) \geq \frac{b}{2} > 0$  for all  $x \in \Omega$ . We then deduce from the limit in (5.7), the weak limit of  $\{\bar{y}_h\}$  in  $H_0^1(\Omega)$ , and Theorem 1.3 in [25] that

$$\begin{aligned} \int_{\Omega} (b(x) - \frac{b}{2}) |\nabla \bar{y}|^2 \, dx &\leq \liminf_{h \rightarrow 0} \int_{\Omega} (b(x) - \frac{b}{2}) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx \\ &\leq \limsup_{h \rightarrow 0} \int_{\Omega} (b(x) - \frac{b}{2}) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx \\ &= \limsup_{h \rightarrow 0} \left[ \int_{\Omega} \bar{u}_h \bar{y}_h \, dx - \int_{\Omega} (\frac{b}{2} + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx \right] \\ &= \int_{\Omega} \bar{u} \bar{y} \, dx - \liminf_{h \rightarrow 0} \int_{\Omega} (\frac{b}{2} + a(\bar{y}_h)) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx \\ &\leq \int_{\Omega} \bar{u} \bar{y} \, dx - \int_{\Omega} (\frac{b}{2} + a(\bar{y})) \nabla \bar{y} \cdot \nabla \bar{y} \, dx \\ &= \int_{\Omega} (b(x) - \frac{b}{2}) |\nabla \bar{y}|^2 \, dx, \end{aligned}$$

where we have employed the equations for  $\bar{y}_h$  and  $\bar{y}$  to derive the first and the last identities. We therefore have

$$\lim_{h \rightarrow 0} \int_{\Omega} \left( b(x) - \frac{b}{2} \right) \nabla \bar{y}_h \cdot \nabla \bar{y}_h \, dx = \int_{\Omega} \left( b(x) - \frac{b}{2} \right) |\nabla \bar{y}|^2 \, dx,$$

which, along with the weak limit, yields, the strong convergence of  $\{\bar{y}_h\}$  in  $H_0^1(\Omega)$ .  $\square$

Next, we prove a kind of converse theorem. More precisely, we assume that  $\bar{u} \in \mathcal{U}_{ad}$  is a *strict local minimum* of (P) with associated state  $\bar{y}$ , i.e. there exists a constant  $\bar{\varepsilon} > 0$  such that

$$j(\bar{u}) < j(u) \quad \text{for all } u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\varepsilon}) \cap \mathcal{U}_{ad} \text{ with } u \neq \bar{u}.$$

We can obviously assume that  $\bar{\varepsilon} < \rho$ . Here  $\rho$  is defined in Theorem 3.2. We therefore can put the discrete operator  $S_h$  into  $(P_h)$ . Then, for any  $h \in (0, h_2)$ , we consider the discretized optimal control problem defined via

$$(P_h^{\bar{\varepsilon}}) \quad \min_{u_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\varepsilon})} j_h(u_h)$$

and the discretized cost functional given by

$$j_h : \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \rightarrow \mathbb{R}, \quad j_h(u) := \int_{\Omega} L(x, (S_h(u))(x)) \, dx + \frac{\nu}{2} \|u\|_{L^2(\Omega)}^2.$$

Using Theorem 3.7 and Assumption (A4), we can show differentiability of  $j_h$ . The proof of the following result is straightforward and therefore omitted.

**Theorem 5.2.** *For any  $h \in (0, h_2)$ , the discrete cost functional  $j_h : \bar{B}_{L^2(\Omega)}(\bar{u}, \rho) \rightarrow \mathbb{R}$  is of class  $C^1$ , and its derivative at  $u \in B_{L^2(\Omega)}(\bar{u}, \rho)$  is given by*

$$j_h'(u)w = \int_{\Omega} (\varphi_h(u) + \nu u)w \, dx \quad \text{for all } w \in L^2(\Omega),$$

where  $\varphi_h(u) \in V_h$  is the unique solution to (4.1) with  $v := \frac{\partial L}{\partial y}(\cdot, S_h(u))$ .

Since  $\mathcal{I}_h \bar{u} \rightarrow \bar{u}$  strongly in  $L^2(\Omega)$  as  $h \rightarrow 0$ , there exists a constant  $\tilde{h} = \tilde{h}(\bar{\varepsilon}) > 0$  such that the admissible set of  $(P_h^{\bar{\varepsilon}})$  is nonempty for all  $0 < h < \tilde{h}$ . We now provide a result on the existence of global minimizers and the associated optimality conditions of  $(P_h^{\bar{\varepsilon}})$ . Its proof is elementary and is thus omitted.

**Theorem 5.3.** *There exists a constant  $h_* \in (0, h_2)$  such that for any  $h \in (0, h_*)$ ,  $(P_h^{\bar{\varepsilon}})$  admits at least one global minimizer  $\bar{u}_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\varepsilon})$ . Moreover, there exists a function  $\bar{\varphi}_h \in V_h$  that together with  $\bar{u}_h$  and  $\bar{y}_h := S_h(\bar{u}_h)$  satisfies*

$$(5.10a) \quad \int_{\Omega} (b + a(\bar{y}_h)) \nabla \bar{\varphi}_h \cdot \nabla w_h + \mathbb{1}_{\{\bar{y}_h \neq \bar{\varepsilon}\}} a'(\bar{y}_h) w_h \nabla \bar{y}_h \cdot \nabla \bar{\varphi}_h \, dx = \int_{\Omega} \frac{\partial L}{\partial y}(x, \bar{y}_h) w_h \, dx,$$

$$(5.10b) \quad \int_{\Omega} (\bar{\varphi}_h + v \bar{u}_h)(u_h - \bar{u}_h) \, dx \geq 0 \quad \text{for all } w_h \in V_h, u_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\varepsilon}).$$

We now state a convergence result in  $L^2(\Omega)$ , whose proof is similar to that of Theorem 4.2 in [26] and is thus omitted here.

**Theorem 5.4.** *Let  $\{\bar{u}_h\}$  be the sequence of discrete solutions to  $(P_h^{\bar{\varepsilon}})$ , defined in Theorem 5.3. Then  $\|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} \rightarrow 0$  as  $h \rightarrow 0^+$ .*

**Remark 5.5.** By Theorem 5.4, it holds that  $\|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} \leq \frac{\bar{\varepsilon}}{2}$  for all  $h \in (0, \tilde{h})$  and for some  $\tilde{h} > 0$ . Now for any  $h \in (0, \tilde{h})$  and  $u_h \in \mathcal{U}_{ad,h}$ , we have that  $w_h := t(u_h - \bar{u}_h) + \bar{u}_h \in \bar{B}_{L^2(\Omega)}(\bar{u}_h, \frac{\bar{\varepsilon}}{2})$  for  $t > 0$  small enough and hence that  $w_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \bar{\varepsilon})$ . The variational inequality (5.10b) then implies that

$$\int_{\Omega} (\bar{\varphi}_h + v \bar{u}_h)(u_h - \bar{u}_h) \, dx \geq 0 \quad \text{for all } u_h \in \mathcal{U}_{ad,h}.$$

**Remark 5.6.** In view of Theorem 5.4, there exists a constant  $h_{\bar{\varepsilon}} > 0$  such that any solution  $\bar{u}_h$  of  $(P_h^{\bar{\varepsilon}})$  belongs to the open ball  $B_{L^2(\Omega)}(\bar{u}, \bar{\varepsilon})$ . By Theorem 3.2,  $(S_h(\bar{u}_h), \bar{u}_h)$  is thus a local minimizer of  $(P_h)$ . We have therefore shown that any strict local solution of (P) can be approximated by local optimal controls of  $(P_h)$ .

In order to show convergence in  $L^\infty(\Omega)$ , we first need the following lemma.

**Lemma 5.7.** *Let  $q > 2$  be given and let  $\bar{h}$  and  $\bar{\rho}$  be defined in Theorem 4.5. If  $\Sigma(\bar{y}) < \infty$ , then for any  $h \in (0, \bar{h})$  and  $u, v \in \mathcal{U}_{ad}$  such that  $v \in B_{L^2(\Omega)}(\bar{u}, \bar{\rho}) \cap \mathcal{U}_{ad}$ , there hold*

$$(5.11) \quad \|y_u - y_h(v)\|_{L^2(\Omega)} + h \|y_u - y_h(v)\|_{H_0^1(\Omega)} \leq C(h^2 + \|u - v\|_{L^2(\Omega)}),$$

$$(5.12) \quad \|\varphi_u - \varphi_v\|_{H^2(\Omega)} \leq C\|u - v\|_{L^2(\Omega)},$$

$$(5.13) \quad \|\varphi_u - \varphi_h(v)\|_{L^2(\Omega)} \leq C(\varepsilon_h^v + \|u - v\|_{L^2(\Omega)}),$$

$$(5.14) \quad \|\varphi_u - \varphi_h(v)\|_{H_0^1(\Omega)} \leq C(h + \|u - v\|_{L^2(\Omega)}),$$

$$(5.15) \quad \|\varphi_u - \varphi_h(v)\|_{L^\infty(\Omega)} \leq C(h^{1-\frac{2}{q}} + \|u - v\|_{L^2(\Omega)}),$$

for some constant  $C$  independent of  $u, v$ , and  $h$ . Here  $y_u := S(u)$  and  $y_h(v) := S_h(v)$ , while  $\varphi_u$  is the unique solution to (2.5) for  $v := \frac{\partial L}{\partial y}(\cdot, y_u)$  and  $\varphi_h(v)$  is the unique solution to (4.1) for  $y_h := S_h(v)$  and  $v := \frac{\partial L}{\partial y}(\cdot, y_h(v))$ .

*Proof.* First, a standard argument yields (5.11). For the other estimates, let  $\varphi_{v,h}$  be the solution to (2.5) for  $v := \frac{\partial L}{\partial y}(\cdot, y_h(v))$  and  $y_u$  replaced by  $y_v := S(v)$ . We need to show that

$$(5.16) \quad \|\varphi_{v,h} - \varphi_u\|_{H^2(\Omega)} \leq C(\|u - v\|_{L^2(\Omega)} + h^2)$$

for some constant  $C > 0$  independent of  $u, v$ , and  $h$ . To this end, we subtract the equations for  $\varphi_u$  and  $\varphi_{v,h}$  to obtain that  $\varphi_{v,h} - \varphi_u \in H_0^1(\Omega)$  and

$$(5.17) \quad -\operatorname{div}[(b + a(y_v)) \nabla(\varphi_{v,h} - \varphi_u)] + \mathbb{1}_{\{y_v \neq \bar{\varepsilon}\}} a'(y_v) \nabla y_v \cdot \nabla(\varphi_{v,h} - \varphi_u) = g_{u,v,h}$$

with

$$g_{u,v,h} := -\operatorname{div}[(a(y_u) - a(y_v))\nabla\varphi_u] + Z_{y_u,y_v} \cdot \nabla\varphi_u + \frac{\partial L}{\partial y}(\cdot, y_h(v)) - \frac{\partial L}{\partial y}(\cdot, y_u).$$

**Theorem 2.3** and **Assumption (A4)** imply that  $\varphi_u \in H^2(\Omega)$ . From this, the product formula, the chain rule [33], and the finiteness of the set  $\{\bar{I}\}$ , we deduce that

$$\operatorname{div}[(a(y_u) - a(y_v))\nabla\varphi_u] = Z_{y_u,y_v} \cdot \nabla\varphi_u + (a(y_u) - a(y_v))\Delta\varphi_u.$$

This shows that

$$g_{u,v,h} = (a(y_v) - a(y_u))\Delta\varphi_u + \frac{\partial L}{\partial y}(\cdot, y_h(v)) - \frac{\partial L}{\partial y}(\cdot, y_u) \in L^2(\Omega).$$

The standard stability estimate for the solution  $\varphi_{v,h} - \varphi_u$  to (5.17) thus gives

$$\|\varphi_{v,h} - \varphi_u\|_{H^2(\Omega)} \leq C\|g_{u,v,h}\|_{L^2(\Omega)} \leq C[\|y_u - y_v\|_{L^\infty(\Omega)} + \|y_h(v) - y_u\|_{L^2(\Omega)}]$$

for some constant  $C > 0$  not depending on  $u, v$ , and  $h$ , where we have employed the boundedness of  $\{S(w) \mid w \in \mathcal{U}_{ad}\}$  in  $C(\bar{\Omega})$ , the fact that  $\|\Delta\varphi_u\|_{L^2(\Omega)} \leq C$  due to **Theorem 2.3**, and **Assumptions (A3)** and **(A4)** to derive the last estimate. From this, (5.11), and the fact that  $\|y_u - y_v\|_{L^\infty(\Omega)} \leq C\|u - v\|_{L^2(\Omega)}$ , we obtain (5.16). The estimate (5.12) is shown by a similar argument.

We now prove (5.13)–(5.15). According to the triangle inequality, (5.16), **Theorem 4.5**, and the boundedness in  $L^2(\Omega)$  of  $\{\frac{\partial L}{\partial y}(\cdot, y_h(v)) \mid v \in B_{L^2(\Omega)}(\bar{u}, \bar{\rho}) \cap \mathcal{U}_{ad}, h \leq h_2\}$ , we obtain (5.13) and (5.14). Finally, for (5.15), we first see from the continuous embedding  $H_0^1(\Omega) \hookrightarrow L^q(\Omega)$ , the interpolation error and inverse estimates [7] for  $N = 2$  that

$$\begin{aligned} \|\varphi_{v,h} - \varphi_h(v)\|_{L^\infty(\Omega)} &\leq \|\varphi_{v,h} - \Pi_h\varphi_{v,h}\|_{L^\infty(\Omega)} + \|\Pi_h\varphi_{v,h} - \varphi_h(v)\|_{L^\infty(\Omega)} \\ &\leq Ch^1\|\varphi_{v,h}\|_{H^2(\Omega)} + Ch^{-\frac{2}{q}}\|\Pi_h\varphi_{v,h} - \varphi_h(v)\|_{L^q(\Omega)} \\ &\leq Ch^1\|\varphi_{v,h}\|_{H^2(\Omega)} + Ch^{-\frac{2}{q}}\|\Pi_h\varphi_{v,h} - \varphi_h(v)\|_{H_0^1(\Omega)} \\ &\leq C \left[ h^1\|\varphi_{v,h}\|_{H^2(\Omega)} + h^{-\frac{2}{q}}\|\Pi_h\varphi_{v,h} - \varphi_{v,h}\|_{H_0^1(\Omega)} + h^{-\frac{2}{q}}\|\varphi_{v,h} - \varphi_h(v)\|_{H_0^1(\Omega)} \right], \end{aligned}$$

which together with **Theorem 4.5** and the interpolation error estimate from **Theorem 4.4.20** in [7] yields

$$\|\varphi_{v,h} - \varphi_h(v)\|_{L^\infty(\Omega)} \leq C \left[ h^{1-\frac{2}{q}}\|\varphi_{v,h}\|_{H^2(\Omega)} + h^{1-\frac{2}{q}}\left\|\frac{\partial L}{\partial y}(\cdot, y_h(v))\right\|_{L^2(\Omega)} \right].$$

Combining this with the uniform boundedness of  $\varphi_{v,h}$  in  $H^2(\Omega)$  and of  $\frac{\partial L}{\partial y}(\cdot, y_h(v))$  in  $L^2(\Omega)$  for all  $v \in B_{L^2(\Omega)}(\bar{u}, \bar{\rho}) \cap \mathcal{U}_{ad}$  and  $h \leq h_2$ , we conclude that  $\|\varphi_{v,h} - \varphi_h(v)\|_{L^\infty(\Omega)} \leq Ch^{1-\frac{2}{q}}$ . From this, (5.16), and the embedding  $H^2(\Omega) \hookrightarrow L^\infty(\Omega)$ , the triangle inequality thus leads to (5.15).  $\square$

From **Theorem 5.4**, **Remark 5.6**, and the estimate (5.15) in **Lemma 5.7**, we obtain the desired convergence result in  $L^\infty(\Omega)$ . Its proof is similar to that of **Theorem 5.3** in [16] with some modifications and it is thus omitted.

**Theorem 5.8.** *Let  $\{(\bar{y}_h, \bar{u}_h)\}$  be the sequence of discrete solutions to  $(P_h)$  converging strongly to  $(\bar{y}, \bar{u})$  in  $H_0^1(\Omega) \times L^2(\Omega)$ . If  $\Sigma(\bar{y}) < \infty$ , then  $\|\bar{u}_h - \bar{u}\|_{L^\infty(\Omega)} \rightarrow 0$  as  $h \rightarrow 0^+$ .*

## 5.2 ERROR ESTIMATES FOR DISCRETE MINIMIZERS

We finally turn to error estimates for discrete local minimizers  $\bar{u}_h$  under the second-order sufficient optimality condition (2.18). We need the following technical lemma.

**Lemma 5.9.** *Let  $\bar{u} \in \mathcal{U}_{ad}$  and  $\bar{y} := S(\bar{u})$ . Assume that  $\{\bar{y} = \bar{t}\}$  decomposes into finitely many connected components and that on each such connected component  $C$ , either (2.16) or (2.17) is fulfilled. Let  $\{v_n\} \subset L^2(\Omega)$ ,  $v \in L^2(\Omega)$ ,  $\varphi \in C^1(\bar{\Omega}) \cap W^{2,1}(\Omega)$ , and  $\{s_n\} \in c_0^+$  be arbitrary such that  $\bar{u} + s_n v_n \in \mathcal{U}_{ad}$ ,  $\|v_n\|_{L^2(\Omega)} = 1$  and  $v_n \rightharpoonup v$  in  $L^2(\Omega)$ . Setting  $u_n := \bar{u} + s_n v_n$ ,  $y_n := S(u_n)$  for  $n \geq 1$  and  $w := S'(\bar{u})v$ , then the following assertions hold:*

(a) *If  $w_n \rightarrow w$  in  $C(\bar{\Omega})$ , then*

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}} \cdot \nabla \varphi w_n \, dx &= -[a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y}=\bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}|>0\}} w^2 \frac{\nabla \bar{y} \cdot \nabla \varphi}{|\nabla \bar{y}|} \, d\mathcal{H}^1(x) \\ &\quad + \int_{\Omega} [\mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a''(\bar{y}) w^2 \nabla \bar{y} + a'(\bar{y}; w) \nabla w] \cdot \nabla \varphi \, dx. \end{aligned}$$

(b) *If in addition (2.7b) is fulfilled for  $\bar{\varphi} := \varphi$ , then*

$$\liminf_{n \rightarrow \infty} \frac{j'(u_n)v_n - j'(\bar{u})v_n}{s_n} = 2Q(\bar{u}, \bar{y}, \varphi; v) + \nu \left(1 - \|v\|_{L^2(\Omega)}^2\right).$$

*Proof.* We first observe from Theorem 2.2 and the embedding  $L^2(\Omega) \hookrightarrow W^{-1, \bar{p}}(\Omega)$  that

$$(5.18) \quad \frac{y_n - \bar{y}}{s_n} \rightarrow w = S'(\bar{u})v \text{ in } W_0^{1, \bar{p}}(\Omega) \hookrightarrow C(\bar{\Omega}).$$

Moreover, by Theorem 2.2 and the compact embedding  $W^{2,p}(\Omega) \Subset C^1(\bar{\Omega})$  for some  $p > 2$ , we deduce from the boundedness in  $L^\infty(\Omega)$  of  $\mathcal{U}_{ad}$  that

$$(5.19) \quad y_n \rightarrow \bar{y} \text{ strongly in } C^1(\bar{\Omega}).$$

*Ad (a):* We first deduce for  $n \in \mathbb{N}$  large enough that

$$(5.20) \quad \tau_n := \|y_n - \bar{y}\|_{C(\bar{\Omega})} \leq C s_n < \delta,$$

where  $\delta$  is the constant defined in (2.9). Moreover, there exists a constant  $M > 0$  such that  $\|y_n\|_{C(\bar{\Omega})}, \|\bar{y}\|_{C(\bar{\Omega})} \leq M$  for all  $n \geq 1$ . According to Lemma 3.4, we have

$$(5.21) \quad Z_{y_n, \bar{y}} = Z_{y_n, \bar{y}}^{(1)} + Z_{y_n, \bar{y}}^{(2)} + Z_{y_n, \bar{y}}^{(3)} + Z_{y_n, \bar{y}}^{(4)}$$

with  $Z_{y_n, \bar{y}}^{(k)}$ ,  $k = 1, 2, 3, 4$ , defined in Lemma 3.4. For  $Z_{y_n, \bar{y}}^{(1)}$ , using (5.20) yields

$$\begin{aligned} Z_{y_n, \bar{y}}^{(1)} &= \mathbb{1}_{\{\bar{y} \in (-\infty, \bar{t}), y_n \in (-\infty, \bar{t})\}} [a'_0(y_n) \nabla y_n - a'_0(\bar{y}) \nabla \bar{y}] + \mathbb{1}_{\{\bar{y} \in (\bar{t}, \infty), y_n \in (\bar{t}, \infty)\}} [a'_1(y_n) \nabla y_n - a'_1(\bar{y}) \nabla \bar{y}], \\ &= [\mathbb{1}_{\{\bar{y} \in (\infty, \bar{t})\}} - \mathbb{1}_{\{\bar{y} \in (-\infty, \bar{t}), y_n \in [\bar{t}, \bar{t} + \delta]\}}] [(a'_0(y_n) - a'_0(\bar{y})) \nabla y_n + a'_0(\bar{y}) \nabla (y_n - \bar{y})] \\ &\quad + [\mathbb{1}_{\{\bar{y} \in (\bar{t}, \infty)\}} - \mathbb{1}_{\{\bar{y} \in (\bar{t}, \infty), y_n \in (\bar{t} - \delta, \bar{t})\}}] [(a'_1(y_n) - a'_1(\bar{y})) \nabla y_n + a'_1(\bar{y}) \nabla (y_n - \bar{y})]. \end{aligned}$$

Since

$$\mathbb{1}_{\{\bar{y} \in (-\infty, \bar{t}), y_n \in [\bar{t}, \bar{t} + \delta]\}}, \mathbb{1}_{\{\bar{y} \in (\bar{t}, \infty), y_n \in (\bar{t} - \delta, \bar{t})\}} \leq \mathbb{1}_{\{0 < |\bar{y} - \bar{t}| \leq \tau_n\}} \rightarrow 0$$

a.e. in  $\Omega$ , we have from (5.18), (5.19), and the Lebesgue dominated convergence theorem that

$$\begin{aligned} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(1)} \cdot \nabla \varphi w_n \, dx &\rightarrow \int_{\Omega} \mathbb{1}_{\bar{y} \in (-\infty, \bar{t})} [a_0''(\bar{y}) w \nabla \bar{y} + a_0'(\bar{y}) \nabla w] \cdot \nabla \varphi w \, dx \\ &+ \int_{\Omega} \mathbb{1}_{\bar{y} \in (\bar{t}, \infty)} [a_1''(\bar{y}) w \nabla \bar{y} + a_1'(\bar{y}) \nabla w] \cdot \nabla \varphi w \, dx \\ &= \int_{\Omega} \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} [a''(\bar{y}) w^2 \nabla \bar{y} + a'(\bar{y}) w \nabla w] \cdot \nabla \varphi \, dx. \end{aligned}$$

For  $Z_{y_n, \bar{y}}^{(2)}$ , we see from (5.20) and the fact  $\nabla \bar{y} = 0$  a.e. on  $\{\bar{y} = \bar{t}\}$  that

$$Z_{y_n, \bar{y}}^{(2)} = \mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}-\delta, \bar{t})\}} a_0'(y_n) \nabla(y_n - \bar{y}) + \mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}, \bar{t}+\delta)\}} a_1'(y_n) \nabla(y_n - \bar{y}).$$

Setting  $\hat{w}_n := \frac{y_n - \bar{y}}{s_n}$  and exploiting (5.18) yields  $w_n - \hat{w}_n \rightarrow w - w = 0$  in  $W_0^{1, \bar{p}}(\Omega) \hookrightarrow C(\bar{\Omega})$ . From this, the limit  $y_n \rightarrow \bar{y}$  in  $C(\bar{\Omega})$ , and the continuity of  $a_0'$  and  $a_1'$ , the dominated convergence theorem implies that

$$\lim_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(2)} \cdot \nabla \varphi w_n \, dx = \lim_{n \rightarrow \infty} \int_{\Omega} [\mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}-\delta, \bar{t})\}} a_0'(\bar{t}) \hat{w}_n + \mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}, \bar{t}+\delta)\}} a_1'(\bar{t}) \hat{w}_n] \nabla \hat{w}_n \cdot \nabla \varphi \, dx.$$

As a result of (2.2) and the fact that  $\hat{w}_n < 0$  on  $\{\bar{y} = \bar{t}, y_n \in (\bar{t} - \delta, \bar{t})\}$ , there holds

$$\mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}-\delta, \bar{t})\}} a_0'(\bar{t}) \hat{w}_n = \mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}-\delta, \bar{t})\}} a'(\bar{t}; \hat{w}_n).$$

Similarly, one has

$$\mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}, \bar{t}+\delta)\}} a_1'(\bar{t}) \hat{w}_n = \mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}, \bar{t}+\delta)\}} a'(\bar{t}; \hat{w}_n).$$

We thus have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(2)} \cdot \nabla \varphi w_n \, dx &= \lim_{n \rightarrow \infty} \int_{\Omega} \mathbb{1}_{\{\bar{y}=\bar{t}, y_n \in (\bar{t}-\delta, \bar{t}) \cup (\bar{t}, \bar{t}+\delta)\}} a'(\bar{t}; \hat{w}_n) \nabla \hat{w}_n \cdot \nabla \varphi \, dx \\ &= \lim_{n \rightarrow \infty} \int_{\Omega} \mathbb{1}_{\{\bar{y}=\bar{t}\}} a'(\bar{t}; \hat{w}_n) \nabla \hat{w}_n \cdot \nabla \varphi \, dx, \end{aligned}$$

where we have used (5.20) and the fact that  $\nabla y_n = \nabla \bar{y} = 0$  and so  $\nabla \hat{w}_n = 0$  a.e. on  $\{y_n = \bar{y} = \bar{t}\}$  to obtain the last identity. We thus conclude from the continuity of  $a'(\bar{t}; \cdot)$  due to Proposition 2.49 in [6], (5.18), and the dominated convergence theorem that

$$(5.22) \quad \lim_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(2)} \cdot \nabla \varphi w_n \, dx = \int_{\Omega} \mathbb{1}_{\{\bar{y}=\bar{t}\}} a'(\bar{y}; w) \nabla w \cdot \nabla \varphi \, dx.$$

For  $Z_{y_n, \bar{y}}^{(4)}$ , we have from (3.8) that

$$|Z_{y_n, \bar{y}}^{(4)}| \leq C_M [ |y_n - \bar{y}| |\nabla \bar{y}| + |\nabla(y_n - \bar{y})| ] \left( \mathbb{1}_{\Omega_{y_n, \bar{y}}^2} + \mathbb{1}_{\Omega_{y_n, \bar{y}}^3} \right) \quad \text{a.e. in } \Omega.$$

This, together with the fact that  $\mathbb{1}_{\Omega_{y_n, \bar{y}}^2} + \mathbb{1}_{\Omega_{y_n, \bar{y}}^3} \rightarrow 0$  a.e. in  $\Omega$  as well as (5.18), yields

$$(5.23) \quad \lim_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(4)} \cdot \nabla \varphi w_n \, dx = 0.$$

It remains to estimate  $Z_{y_n, \bar{y}}^{(3)}$ . To this end, we first deduce from (4.16) and the coarea formula for Lipschitz mappings (see, e.g. [30], Thm. 2, p. 117) or [1], Sec. 2.7) that

$$\begin{aligned} s_n^{-1} \left| \int_{\Omega} \mathbb{1}_{\Omega^2_{y_n, \bar{y}}} \nabla \bar{y} \cdot \nabla \varphi(w_n - \hat{w}_n) dx \right| &\leq s_n^{-1} \|w_n - \hat{w}_n\|_{L^\infty(\Omega)} \|\nabla \varphi\|_{L^\infty(\Omega)} \int_{\Omega} \mathbb{1}_{\{0 < \bar{y} - \bar{t} \leq \tau_n\}} |\nabla \bar{y}| dx \\ &= s_n^{-1} \|w_n - \hat{w}_n\|_{L^\infty(\Omega)} \|\nabla \varphi\|_{L^\infty(\Omega)} \int_{\bar{t}}^{\bar{t} + \tau_n} \int_{\{\bar{y} = t_i\}} d\mathcal{H}^1(x) dt \\ &\leq C s_n^{-1} \tau_n \|w_n - \hat{w}_n\|_{L^\infty(\Omega)} \|\nabla \varphi\|_{L^\infty(\Omega)} \rightarrow 0, \end{aligned}$$

where we have used (5.20) and the fact that  $w_n - \hat{w}_n \rightarrow 0$  in  $C(\bar{\Omega})$  to obtain the last limit. Similarly,  $s_n^{-1} \left| \int_{\Omega} \mathbb{1}_{\Omega^3_{y_n, \bar{y}}} \nabla \bar{y} \cdot \nabla \varphi(w_n - \hat{w}_n) dx \right| \rightarrow 0$ . From these limits and the definition of  $Z_{y_n, \bar{y}}^{(3)}$ , we deduce that

$$s_n^{-1} \int_{\Omega} |Z_{y_n, \bar{y}}^{(3)} \cdot \nabla \varphi(w_n - \hat{w}_n)| dx \rightarrow 0$$

and thus

$$(5.24) \quad \lim_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(3)} \cdot \nabla \varphi w_n dx = \lim_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(3)} \cdot \nabla \varphi \hat{w}_n dx = \lim_{n \rightarrow \infty} \frac{1}{s_n^2} \int_{\Omega} Z_{y_n, \bar{y}}^{(3)} \cdot \nabla \varphi (y_n - \bar{y}) dx$$

provided that one of these three limits exists. For  $\zeta_i(\bar{u}, \bar{y}; s_n, v_n)$ ,  $i = 0, 1$ , defined in (2.10), one has

$$\begin{aligned} P_n &:= Z_{y_n, \bar{y}}^{(3)} (y_n - \bar{y}) + 2 \sum_{i=0}^1 \zeta_i(\bar{u}, \bar{y}; s_n, v_n) \nabla \bar{y} \\ &= [a'_0(\bar{t}) - a'_1(\bar{t})] [\mathbb{1}_{\Omega^2_{y_n, \bar{y}}} - \mathbb{1}_{\Omega^3_{y_n, \bar{y}}}] \nabla \bar{y} (y_n - \bar{y}) + 2 [a'_0(\bar{t}) - a'_1(\bar{t})] (\bar{t} - y_n) [\mathbb{1}_{\Omega^2_{y_n, \bar{y}}} - \mathbb{1}_{\Omega^3_{y_n, \bar{y}}}] \nabla \bar{y} \\ &= [a'_0(\bar{t}) - a'_1(\bar{t})] (2\bar{t} - \bar{y} - y_n) [\mathbb{1}_{\Omega^2_{y_n, \bar{y}}} - \mathbb{1}_{\Omega^3_{y_n, \bar{y}}}] \nabla \bar{y}. \end{aligned}$$

By (2.20) and (5.19), we deduce that

$$\lim_{n \rightarrow \infty} \frac{1}{s_n^2} \int_{\Omega} P_n \cdot \nabla \varphi dx = 0.$$

Combining this with (5.24) and (2.15), we can conclude that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{s_n} \int_{\Omega} Z_{y_n, \bar{y}}^{(3)} \cdot \nabla \varphi w_n dx &= \limsup_{n \rightarrow \infty} \left[ \frac{1}{s_n^2} \int_{\Omega} P_n \cdot \nabla \varphi dx - \frac{2}{s_n^2} \int_{\Omega} \sum_{i=0}^1 \zeta_i(\bar{u}, \bar{y}; s_n, v_n) \nabla \bar{y} \cdot \nabla \varphi dx \right] \\ &= -2\tilde{Q}(\bar{u}, \bar{y}, \varphi; \{s_n\}, v). \end{aligned}$$

Together with (5.21)–(5.23), we obtain assertion (a) from (2.19).

*Ad (b):* Defining the functional  $G : L^\infty(\Omega) \rightarrow \mathbb{R}$  via  $G(y) := \int_{\Omega} L(x, y(x)) dx$  and employing Assumption (A4), we deduce that  $G$  is of class  $C^2$  and that its derivatives are given by

$$G'(y) y_1 := \int_{\Omega} \frac{\partial L}{\partial y}(x, y(x)) y_1(x) dx \quad \text{and} \quad G''(y) y_1 y_2 := \int_{\Omega} \frac{\partial^2 L}{\partial y^2}(x, y(x)) y_1(x) y_2(x) dx$$

for all  $y, y_1, y_2 \in L^\infty(\Omega)$ . We see from the chain rule that for any  $u, v \in L^2(\Omega)$ ,

$$j'(u)v = G'(S(u))S'(u)v + v \int_{\Omega} uv dx.$$

This, together with a Taylor expansion and the fact that  $\|v_n\|_{L^2(\Omega)} = 1$ , yields

$$\begin{aligned}
(5.25) \quad \frac{1}{s_n} [j'(u_n)v_n - j'(\bar{u})v_n] &= \frac{1}{s_n} [G'(y_n)S'(u_n)v_n - G'(\bar{y})S'(\bar{u})v_n] + \nu \int_{\Omega} v_n^2 dx \\
&= \frac{1}{s_n} [G'(y_n) - G'(\bar{y})]S'(\bar{u})v_n + \frac{1}{s_n} G'(y_n)[S'(u_n)v_n - S'(\bar{u})v_n] + \nu \\
&= \frac{1}{s_n} \int_0^1 G''(\bar{y} + s(y_n - \bar{y}))(y_n - \bar{y})S'(\bar{u})v_n ds + \frac{1}{s_n} G'(\bar{y})[S'(u_n)v_n - S'(\bar{u})v_n] \\
&\quad + \frac{1}{s_n} [G'(y_n) - G'(\bar{y})][S'(u_n)v_n - S'(\bar{u})v_n] + \nu.
\end{aligned}$$

Obviously, the third term on the right-hand side of (5.25) tends to 0 since  $v_n \rightarrow v$  in  $W^{-1,\tilde{p}}(\Omega)$ ,  $S'$  is continuous, and  $G$  is of class  $C^2$ . Moreover, it follows from Assumption (A4), (5.18), and the dominated convergence theorem that the first term on the right-hand side of (5.25) tends to  $G''(\bar{y})(S'(\bar{u})v)^2$ . It remains to estimate the limes inferior of the second term on the right-hand side of (5.25). Subtracting the equations for  $z_n^{(1)} := S'(u_n)v_n$  and  $z_n^{(2)} := S'(\bar{u})v_n$ , we find that  $z_n := z_n^{(1)} - z_n^{(2)} \in H_0^1(\Omega)$  satisfies

$$(5.26) \quad -\operatorname{div}[(b + a(\bar{y}))\nabla z_n + \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'(\bar{y})\nabla \bar{y} z_n] = \operatorname{div}[(a(y_n) - a(\bar{y}))\nabla z_n^{(1)} + Z_{y_n, \bar{y}} z_n^{(1)}] =: g_n.$$

We then have  $z_n = S'(\bar{u})g_n$ , which together with (2.7b) yields

$$(5.27) \quad G'(\bar{y})[z_n^{(1)} - z_n^{(2)}] = \langle G'(\bar{y}), z_n \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = \langle g_n, \varphi \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = -(B_n + C_n)$$

for  $B_n := \int_{\Omega} (a(y_n) - a(\bar{y}))\nabla z_n^{(1)} \cdot \nabla \varphi dx$  and  $C_n := \int_{\Omega} Z_{y_n, \bar{y}} \cdot \nabla \varphi z_n^{(1)} dx$ . As a result of Theorem 2.2 and the fact that  $v_n \rightarrow v$  in  $W^{-1,\tilde{p}}(\Omega)$ , there holds  $z_n^{(1)} \rightarrow S'(\bar{u})v$  in  $W_0^{1,\tilde{p}}(\Omega)$ . Besides, from (5.18) and Lemma 3.5 in [23], we have

$$\frac{a(y_n(x)) - a(\bar{y}(x))}{s_n} \rightarrow a'(\bar{y}(x); (S'(\bar{u})v)(x)) \quad \text{for all } x \in \bar{\Omega}.$$

The dominated convergence theorem thus implies that

$$\frac{1}{s_n} B_n \rightarrow \int_{\Omega} a'(\bar{y}; S'(\bar{u})v) \nabla(S'(\bar{u})v) \cdot \nabla \varphi dx.$$

This, along with (5.27) and assertion (a), ensures that

$$\begin{aligned}
\liminf_{n \rightarrow \infty} \frac{1}{s_n} G'(\bar{y})[S'(u_n)v_n - S'(\bar{u})v_n] &= [a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y}=\bar{t}\}} (S'(\bar{u})v)^2 \frac{\nabla \bar{y} \cdot \nabla \varphi}{|\nabla \bar{y}|} d\mathcal{H}^1(x) \\
&\quad - 2 \int_{\Omega} a'(\bar{y}; S'(\bar{u})v) \nabla(S'(\bar{u})v) \cdot \nabla \varphi dx - \int_{\Omega} \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a''(\bar{y})(S'(\bar{u})v)^2 \nabla \bar{y} \cdot \nabla \varphi dx.
\end{aligned}$$

Using these limits, (5.25), and the identity for  $Q$  in (2.18) (see also Theorem 3.19 in [24]), we arrive at (b).  $\square$

The following theorem is one of main results of the paper, which extends Theorem 2.14 in [17] (see, also, Lemma 5.2 in [13]) to the case where the cost functional  $j$  is of class  $C^1$  but not necessarily  $C^2$ .

**Theorem 5.10.** *Let  $\{(\bar{y}_h, \bar{u}_h)\}$  be the sequence of discrete solutions to  $(P_h)$  converging strongly to  $(\bar{y}, \bar{u})$  in  $H_0^1(\Omega) \times L^2(\Omega)$ . Assume that  $\{\bar{y} = \bar{t}\}$  decomposes into finitely many connected components and that on each such connected component  $C$ , either (2.16) or (2.17) is fulfilled. Assume further that that the second-order sufficient condition (2.18) is fulfilled. Then there exist constants  $C > 0$  and  $\hat{h} \in (0, \min\{\bar{h}, h_*\})$  such that*

$$(5.28) \quad \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}^2 \leq C[(\varepsilon_h^{\bar{u}_h})^2 + \|\bar{u} - u_h\|_{L^2(\Omega)}^2 + j'(\bar{u})(u_h - \bar{u})]$$

for all  $h \in (0, \hat{h})$  and  $u_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \hat{\varepsilon})$  with  $\hat{\varepsilon} := \min\{\bar{\varepsilon}, \bar{\rho}\}$ . Here  $\varepsilon_h^{\bar{u}}$  is defined as in (4.20).

*Proof.* We first observe from Proposition 2.9 that  $\Sigma(\bar{y}) < \infty$ . For simplicity of notation, we set  $\varepsilon_h := \varepsilon_h^{\bar{u}_h}$ . We first show that

$$(5.29) \quad [j'(\bar{u}_h) - j'(\bar{u})](\bar{u}_h - \bar{u}) \leq j'(\bar{u})(u_h - \bar{u}) + C[\varepsilon_h \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} + \varepsilon_h \|u_h - \bar{u}\|_{L^2(\Omega)} + \|u_h - \bar{u}\|_{L^2(\Omega)} \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}]$$

for some constant  $C > 0$ , for all  $u_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \hat{\varepsilon})$  and  $h \in (0, \min\{\bar{h}, h_*\})$ . To this end, let us take any  $h \in (0, \min\{\bar{h}, h_*\})$ ,  $u \in \mathcal{U}_{ad} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \hat{\varepsilon})$ , and  $u_h \in \mathcal{U}_{ad,h} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \hat{\varepsilon})$ . We deduce from (2.6), Theorem 5.2, Lemma 5.7, and the Cauchy–Schwarz inequality that

$$(5.30) \quad |[j'_h(u) - j'(u)](u_h - \bar{u})| = \left| \int_{\Omega} (\varphi_h(u) - \varphi_u)(u_h - \bar{u}) \, dx \right| \leq C\varepsilon_h^u \|u_h - \bar{u}\|_{L^2(\Omega)}.$$

Moreover, we deduce from  $j'_h(\bar{u}_h)(\bar{u}_h - u_h) \leq 0$  and  $j'(\bar{u})(\bar{u} - \bar{u}_h) \leq 0$  that

$$(5.31) \quad \begin{aligned} [j'(\bar{u}_h) - j'(\bar{u})](\bar{u}_h - \bar{u}) &= [j'_h(\bar{u}_h) - j'(\bar{u}_h)](\bar{u} - \bar{u}_h) + [j'_h(\bar{u}_h) - j'(\bar{u})](u_h - \bar{u}) + j'_h(\bar{u}_h)(\bar{u}_h - u_h) + j'(\bar{u})(u_h - \bar{u}_h) \\ &\leq [j'_h(\bar{u}_h) - j'(\bar{u}_h)](\bar{u} - \bar{u}_h) + [j'_h(\bar{u}_h) - j'(\bar{u})](u_h - \bar{u}) + j'(\bar{u})[(u_h - \bar{u}) + (\bar{u} - \bar{u}_h)] \\ &\leq [j'_h(\bar{u}_h) - j'(\bar{u}_h)](\bar{u} - \bar{u}_h) + [(j'_h(\bar{u}_h) - j'(\bar{u}_h)) + (j'(\bar{u}_h) - j'(\bar{u}))](u_h - \bar{u}) + j'(\bar{u})(u_h - \bar{u}). \end{aligned}$$

Applying (5.30) yields that

$$(5.32) \quad \begin{cases} |[j'_h(\bar{u}_h) - j'(\bar{u}_h)](\bar{u} - \bar{u}_h)| \leq C\varepsilon_h \|\bar{u} - \bar{u}_h\|_{L^2(\Omega)}, \\ |[j'_h(\bar{u}_h) - j'(\bar{u}_h)](u_h - \bar{u})| \leq C\varepsilon_h \|\bar{u} - u_h\|_{L^2(\Omega)}. \end{cases}$$

Using (2.6), (5.12),  $H^2(\Omega) \hookrightarrow L^2(\Omega)$ , and the Cauchy–Schwarz inequality yields

$$(5.33) \quad |[j'(\bar{u}_h) - j'(\bar{u})](u_h - \bar{u})| = \left| \int_{\Omega} [(\varphi_{\bar{u}_h} - \bar{\varphi}) + v(\bar{u}_h - \bar{u})](u_h - \bar{u}) \, dx \right| \leq C \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} \|u_h - \bar{u}\|_{L^2(\Omega)}.$$

From this, (5.31), and (5.32), we derive (5.29).

We now prove the conclusion of the theorem by contradiction. To that purpose, we suppose that there exist  $h_n \rightarrow 0^+$  and  $u_{h_n} \in \mathcal{U}_{ad,h_n} \cap \bar{B}_{L^2(\Omega)}(\bar{u}, \hat{\varepsilon})$  such that

$$\|\bar{u}_{h_n} - \bar{u}\|_{L^2(\Omega)}^2 > n \left[ (\varepsilon_{h_n})^2 + \|\bar{u} - u_{h_n}\|_{L^2(\Omega)}^2 + j'(\bar{u})(u_{h_n} - \bar{u}) \right] \quad \text{for all } n \geq 1$$

or, equivalently, with  $s_n := \|\bar{u}_{h_n} - \bar{u}\|_{L^2(\Omega)}$  that

$$(5.34) \quad \frac{1}{n} > \frac{(\varepsilon_{h_n})^2}{s_n^2} + \frac{\|\bar{u} - u_{h_n}\|_{L^2(\Omega)}^2}{s_n^2} + \frac{j'(\bar{u})(u_{h_n} - \bar{u})}{s_n^2} \quad \text{for all } n \geq 1.$$

By setting  $v_n := \frac{\bar{u}_{h_n} - \bar{u}}{s_n}$ , and by extracting a subsequence if necessary, we have

$$s_n \rightarrow 0^+, \quad \|v_n\|_{L^2(\Omega)} = 1, \quad \bar{u}_{h_n} = \bar{u} + s_n v_n, \quad v_n \rightharpoonup v \text{ in } L^2(\Omega) \quad \text{for some } v \in L^2(\Omega).$$

We first show that  $v$  is an element of the critical cone  $C(\mathcal{U}_{ad}; \bar{u})$  defined in (2.8). To this end, we first deduce that  $v \geq 0$  a.e. on  $\{\bar{u} = \alpha\}$  and  $v \leq 0$  a.e. on  $\{\bar{u} = \beta\}$ . Moreover, since  $j'(\bar{u})v_n \geq 0$ , there holds  $j'(\bar{u})v \geq 0$ . On the other hand, from (5.32) and (5.33) for  $u_h := \bar{u}_{h_n}$ , we obtain that

$$\limsup_{n \rightarrow \infty} \left\{ [j'(\bar{u}_{h_n}) - j'_{h_n}(\bar{u}_{h_n})]v_n + [j'(\bar{u}) - j'(\bar{u}_{h_n})]v_n \right\} \leq \lim_{n \rightarrow \infty} C(\varepsilon_{h_n} + s_n) = 0,$$

which yields

$$\begin{aligned} j'(\bar{u})v &= \lim_{n \rightarrow \infty} j'(\bar{u})v_n = \lim_{n \rightarrow \infty} [j'_{h_n}(\bar{u}_{h_n})v_n + (j'(\bar{u}_{h_n}) - j'_{h_n}(\bar{u}_{h_n}))v_n + (j'(\bar{u}) - j'(\bar{u}_{h_n}))v_n] \\ &\leq \lim_{n \rightarrow \infty} j'_{h_n}(\bar{u}_{h_n})v_n = \lim_{n \rightarrow \infty} \frac{1}{s_n} [j'_{h_n}(\bar{u}_{h_n})(u_{h_n} - \bar{u}) + j'_{h_n}(\bar{u}_{h_n})(\bar{u}_{h_n} - u_{h_n})]. \end{aligned}$$

From this and the fact that  $j'_{h_n}(\bar{u}_{h_n})(\bar{u}_{h_n} - u_{h_n}) \leq 0$ , we obtain

$$j'(\bar{u})v \leq \lim_{n \rightarrow \infty} \frac{1}{s_n} j'_{h_n}(\bar{u}_{h_n})(u_{h_n} - \bar{u}) \leq \lim_{n \rightarrow \infty} \frac{\|\varphi_{h_n}(\bar{u}_{h_n}) + v\bar{u}_{h_n}\|_{L^2(\Omega)} \|u_{h_n} - \bar{u}\|_{L^2(\Omega)}}{s_n} \rightarrow 0,$$

where we have used [Theorem 5.2](#) and the Cauchy–Schwarz inequality to derive the last estimate and the boundedness of  $\{\|\varphi_{h_n}(\bar{u}_{h_n}) + v\bar{u}_{h_n}\|_{L^2(\Omega)}\}$  (due to [Lemma 5.7](#)) as well as (5.34) to pass to the limit. There therefore holds that  $j'(\bar{u})v = 0$ . This and [Lemma 4.11](#) in [3] lead to  $v(x) = 0$  whenever  $\bar{\varphi}(x) + v\bar{u}(x) \neq 0$ . We thus have  $v \in C(\mathcal{U}_{ad}; \bar{u})$ .

We now derive a contradiction and thus complete the proof. To this end, we divide (5.29) (with  $h := h_n$ ) by  $s_n^2$  to obtain

$$\frac{1}{s_n} [j'(\bar{u}_{h_n}) - j'(\bar{u})]v_n \leq \frac{j'(\bar{u})(u_{h_n} - \bar{u})}{s_n^2} + C \left( \frac{\varepsilon_{h_n}}{s_n} + \frac{\varepsilon_{h_n}}{s_n} \frac{\|u_{h_n} - \bar{u}\|_{L^2(\Omega)}}{s_n} + \frac{\|u_{h_n} - \bar{u}\|_{L^2(\Omega)}}{s_n} \right).$$

Taking the limes inferior as  $n \rightarrow \infty$ , employing (5.34), and using [Lemma 5.9](#) (ii), we conclude that

$$(5.35) \quad 2Q(\bar{u}, \bar{y}, \bar{\varphi}; v) + \nu \left(1 - \|v\|_{L^2(\Omega)}^2\right) \leq 0.$$

Combining this with (2.18) and the fact that  $\|v\|_{L^2(\Omega)} \leq \liminf_{n \rightarrow \infty} \|v_n\|_{L^2(\Omega)} = 1$ , we have  $v = 0$ . Inserting this into (5.35) leads to  $0 < \nu \leq 0$ , which is the desired contradiction.  $\square$

**Theorem 5.11 (variational discretization).** *Assume that  $\mathcal{U}_h = L^\infty(\Omega)$ . Let  $\{(\bar{y}_h, \bar{u}_h)\}$  be the sequence of discrete solutions to  $(P_h)$  converging strongly to  $(\bar{y}, \bar{u})$  in  $H_0^1(\Omega) \times L^2(\Omega)$ . Under all assumptions of [Theorem 5.10](#), there exists a constant  $C > 0$  such that for any  $h \in (0, \hat{h})$ ,*

$$(5.36) \quad \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} \leq C \left( h^{1+\gamma} + \sigma_0^{\frac{1}{2}} \|\Sigma_{\kappa_h}(\bar{y})\|_{L^1(\Omega)}^{\frac{1}{2}} h^{\frac{3}{2}} \right) \leq Ch^{1+\gamma}$$

with  $\kappa_h := C_\infty h + \|S(\bar{u}_h) - \bar{y}\|_{L^\infty(\Omega)}$ , and  $\sigma_0, \gamma \in (0, \frac{1}{2})$ , and  $\Sigma_{\kappa_h}$  defined in (2.14), (4.18), and (4.10), respectively.

*Proof.* Choosing  $u_h := \bar{u}$  in (5.28) yields

$$(5.37) \quad \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} \leq C\varepsilon_h^{\bar{u}_h} \quad \text{for all } h \in (0, \hat{h}).$$

Setting  $r_h := \|S_h(\bar{u}_h) - S(\bar{u}_h)\|_{L^\infty(\Omega)}$  and using (4.9) yield  $r_h \leq C_\infty h$ . Exploiting (4.20), the Cauchy–Schwarz inequality, [Proposition 4.3](#), the estimate (4.12), and the monotonic growth of  $V(\bar{y}, \cdot)$ , there holds

$$\begin{aligned} (\varepsilon_h^{\bar{u}_h})^2 &\leq 3h^{2+2\gamma} + 3h^2 \|V(S(\bar{u}_h), r_h)\|_{L^2(\Omega)}^2 + 3h^2 \|Z_{S(\bar{u}_h), \bar{y}}\|_{L^2(\Omega)}^2 \\ &\leq 3h^{2+2\gamma} + Ch^2 \left[ \|V(\bar{y}, r_h + \|S(\bar{u}_h) - \bar{y}\|_{L^\infty(\Omega)})\|_{L^2(\Omega)}^2 + \|S(\bar{u}_h) - \bar{y}\|_{H_0^1(\Omega)}^2 + \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}^2 \right] \\ &\leq 3h^{2+2\gamma} + Ch^2 \left[ \|V(\bar{y}, C_\infty h + \|S(\bar{u}_h) - \bar{y}\|_{L^\infty(\Omega)})\|_{L^2(\Omega)}^2 + \|S(\bar{u}_h) - \bar{y}\|_{H_0^1(\Omega)}^2 + \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}^2 \right] \\ &\leq C \left[ h^{2+2\gamma} + h^2 \sigma_0 \|\Sigma_{\kappa_h}(\bar{y})\|_{L^1(\Omega)} (C_\infty h + \|S(\bar{u}_h) - \bar{y}\|_{L^\infty(\Omega)}) + h^2 \|S(\bar{u}_h) - \bar{y}\|_{H_0^1(\Omega)}^2 + h^2 \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}^2 \right] \\ &\leq C \left[ h^{2+2\gamma} + h^2 \sigma_0 \|\Sigma_{\kappa_h}(\bar{y})\|_{L^1(\Omega)} (h + \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}) + h^2 \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)}^2 \right], \end{aligned}$$

where we have used [Theorem 2.2](#) to derive the last inequality. From this and (5.37), a simple computation gives (5.36).  $\square$

**Remark 5.12.** If the constant  $p_*$  in [Theorem 2.2](#) is large enough, then, for arbitrary small  $\varepsilon > 0$ , we can take  $\gamma := \frac{1}{2} - \varepsilon$  in [\(4.18\)](#) by choosing  $p_0$  close to 2 enough. Therefore, the order of convergence in [Theorem 5.11](#) becomes  $O(h^{\frac{3}{2}-\varepsilon})$ . This order is less than the one for the smooth situation investigated in [\[17\]](#), there the authors showed that the order of convergence associated with the variational discretization is  $O(h^2)$ . This fact can be attributed to the nondifferentiability of the function  $a$  in the state equation as we will see later in the numerical example section.

Similarly, we obtain from [Theorem 5.10](#) error estimates for piecewise constant and continuous piecewise linear controls.

**Theorem 5.13** (piecewise constant discretization and continuous piecewise linear discretization). *Assume that  $\mathcal{U}_h = \mathcal{U}_h^i$ ,  $i = 0, 1$ . Let  $\{(\bar{y}_h, \bar{u}_h)\}$  be the sequence of discrete solutions to  $(P_h)$  converging strongly to  $(\bar{y}, \bar{u})$  in  $H_0^1(\Omega) \times L^2(\Omega)$ . Under all assumptions of [Theorem 5.10](#), there exist constants  $C > 0$  and  $\hat{h}_* \in (0, \hat{h})$  such that*

$$(5.38) \quad \|\bar{u}_h - \bar{u}\|_{L^2(\Omega)} \leq Ch \quad \text{for all } h \in (0, \hat{h}_*).$$

*Proof.* According to [Theorem 2.5](#),  $\bar{\varphi}$  and  $\bar{u}$  are Lipschitz continuous on  $\bar{\Omega}$ . Hence constants  $C, C_1 > 0$  and  $\hat{h}_* \in (0, \hat{h})$  exist such that for any  $h \in (0, \hat{h}_*)$ , there exists a  $u_h \in \mathcal{U}_{ad,h}$  satisfying  $\|\bar{u} - u_h\|_{L^\infty(\Omega)} \leq C_1 h$  and  $j'(\bar{u})(\bar{u} - u_h) = 0$  for the case  $\mathcal{U}_h = \mathcal{U}_h^0$ ; see, e.g. Lemma 4.17 in [\[12\]](#), as well as  $|j'(\bar{u})(\bar{u} - u_h)| \leq Ch^2$  for the case  $\mathcal{U}_h = \mathcal{U}_h^1$ ; see the proof of [Theorem 5.4](#) in [\[13\]](#). Combining this with [\(5.28\)](#) and the fact that  $\varepsilon_h^{\bar{u}_h} \leq C_2 h$  for all  $h \in (0, \hat{h})$  yields [\(5.38\)](#).  $\square$

### 5.3 NUMERICAL EXAMPLE

We conclude this section with a preliminary numerical example for the variational discretization of the optimal control problem. Specifically, we consider the problem

$$(5.39) \quad \begin{cases} \min_{u \in L^\infty(\Omega)} j(u) := \frac{1}{2} \int_{\Omega} (y_u(x) - y_d(x))^2 dx + \frac{\nu}{2} \int_{\Omega} u(x)^2 dx \\ \text{s.t.} \quad -\operatorname{div}[(b(x) + \max\{y_u - m, 0\})\nabla y_u] = u \quad \text{in } \Omega, \quad y_u = 0 \text{ on } \partial\Omega, \\ 0 \leq u(x) \leq 4\pi^2 \quad \text{a.e. } x \in \Omega, \end{cases}$$

where  $\Omega := (0, 1)^2 \subset \mathbb{R}^2$ ,  $m \in (0, 1]$ ,  $\nu := 1$ ,  $b(x) := b(x_1, x_2) = 2 - \max\{\sin \pi x_1 \sin \pi x_2 - m, 0\}$ , and

$$y_d(x) := (1 + 16\nu\pi^4) \sin \pi x_1 \sin \pi x_2 + 4\nu\pi^4 \mathbb{1}_{\{\sin \pi x_1 \sin \pi x_2 > m\}} (\cos^2 \pi x_1 \sin^2 \pi x_2 + \sin^2 \pi x_1 \cos^2 \pi x_2).$$

This problem fits the general setting with  $\alpha = 0$ ,  $\beta = 4\pi^2$ ,  $a(t) = \max\{t - m, 0\}$ ; i.e.,  $a_0(t) := 0$ ,  $a_1(t) := t - m$ , and  $\bar{t} := m$ . Setting  $\bar{u} := 4\pi^2 \bar{y}$  with  $\bar{y} := \sin \pi x_1 \sin \pi x_2$  and  $\bar{\varphi} := -\nu \bar{u}$ , it is straightforward to verify that  $(\bar{u}, \bar{y}, \bar{\varphi})$  satisfies the first-order optimality condition [\(2.7\)](#) associated with [\(5.39\)](#). We shall now show that there exists an  $\varepsilon > 0$  such that if  $0 \leq 1 - m < \varepsilon$  then all assumptions of [Theorem 2.7](#), and thus of [Theorem 5.11](#), are fulfilled. First, for  $\bar{t} = m = 1$  we have

$$\{\bar{y} = \bar{t}\} = \{\bar{y} = 1\} = \{(0.5, 0.5)\} \quad \text{and} \quad \{\bar{y} < \bar{t}\} = \Omega \setminus \{(0.5, 0.5)\}.$$

Moreover,  $\nabla \bar{y}$  obviously vanishes on  $\{\bar{y} = 1\}$ , and [Lemma A.1](#) in the Appendix shows that the structural assumption in [\(2.17\)](#) holds. For  $\bar{t} = m \in (0, 1)$ , a simple computation shows that

$$|\nabla \bar{y}(x)| > 0 \quad \text{for all } x = (x_1, x_2) \in \{\bar{y} = \bar{t}\},$$

which validates [\(2.16\)](#). It remains to show the existence of a number  $\varepsilon > 0$  such that the second-order sufficient optimality condition [\(2.18\)](#) is fulfilled, provided that  $0 \leq 1 - m < \varepsilon$ . To this end, by virtue of

(2.2), there holds for a.e.  $(x_1, x_2) \in \Omega$  that  $a'(\bar{y}(x_1, x_2); s) = \mathbb{1}_{\{\bar{y}(x_1, x_2) > \bar{t}\}} s$  for all  $s \in \mathbb{R}$ . From this and the fact that  $\mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'' \equiv 0$ , we have

$$\begin{aligned} Q_s(\bar{u}, \bar{y}, \bar{\varphi}; v, v) + Q_1(\bar{u}, \bar{y}, \bar{\varphi}; v, v) \\ = \frac{1}{2} \|S'(\bar{u})v\|_{L^2(\Omega)}^2 + \frac{\nu}{2} \|v\|_{L^2(\Omega)}^2 - \int_{\Omega} \mathbb{1}_{\{\bar{y} > \bar{t}\}} S'(\bar{u})v \nabla S'(\bar{u})v \cdot \nabla \bar{\varphi} \, dx \quad \text{for all } v \in L^2(\Omega). \end{aligned}$$

Since  $\bar{\varphi} = -v\bar{u} = -4\nu\pi^2\bar{y}$ , there hold

$$Q_s(\bar{u}, \bar{y}, \bar{\varphi}; v, v) + Q_1(\bar{u}, \bar{y}, \bar{\varphi}; v, v) = \frac{1}{2} \|S'(\bar{u})v\|_{L^2(\Omega)}^2 + \frac{\nu}{2} \|v\|_{L^2(\Omega)}^2 + 4\nu\pi^2 \int_{\Omega} \mathbb{1}_{\{\bar{y} > \bar{t}\}} S'(\bar{u})v \nabla S'(\bar{u})v \cdot \nabla \bar{y} \, dx$$

and

$$\begin{aligned} Q_2(\bar{u}, \bar{y}, \bar{\varphi}; v) &= \frac{1}{2} [a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y}=\bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}| > 0\}} (S'(\bar{u})v)^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} \, d\mathcal{H}^1(x) \\ &= 2\nu\pi^2 \int_{\{\bar{y}=\bar{t}\}} (S'(\bar{u})v)^2 |\nabla \bar{y}| \, d\mathcal{H}^1(x) \geq 0 \end{aligned}$$

for all  $v \in L^2(\Omega)$ . Consequently, we have

$$(5.40) \quad Q(\bar{u}, \bar{y}, \bar{\varphi}; v) \geq \frac{1}{2} \|S'(\bar{u})v\|_{L^2(\Omega)}^2 + \frac{\nu}{2} \|v\|_{L^2(\Omega)}^2 + 4\nu\pi^2 \int_{\Omega} \mathbb{1}_{\{\bar{y} > \bar{t}\}} S'(\bar{u})v \nabla S'(\bar{u})v \cdot \nabla \bar{y} \, dx$$

for all  $v \in L^2(\Omega)$ . We now estimate the last term in the right-hand side of (5.40). For that purpose, we observe that

$$\begin{aligned} \mathbb{1}_{\{\bar{y} > \bar{t}\}} |\nabla \bar{y}|^2 &= \pi^2 \mathbb{1}_{\{\sin \pi x_1 \sin \pi x_2 > m\}} (\cos^2 \pi x_1 \sin^2 \pi x_2 + \sin^2 \pi x_1 \cos^2 \pi x_2) \\ &= \pi^2 \mathbb{1}_{\{\sin \pi x_1 \sin \pi x_2 > m\}} (\sin^2 \pi x_1 + \sin^2 \pi x_2 - 2 \sin^2 \pi x_1 \sin^2 \pi x_2) \\ &\leq \pi^2 (2 - 2m^2) \end{aligned}$$

and thus

$$\mathbb{1}_{\{\bar{y} > \bar{t}\}} |\nabla \bar{y}| \leq \pi \sqrt{2(1 - m^2)} \leq 2\pi \sqrt{1 - m}$$

in  $\Omega$ , where we have exploited that  $1 < 1 + m \leq 2$ . From this and the Cauchy–Schwarz inequality, there holds

$$\begin{aligned} \left| \int_{\Omega} \mathbb{1}_{\{\bar{y} > \bar{t}\}} S'(\bar{u})v \nabla S'(\bar{u})v \cdot \nabla \bar{y} \, dx \right| &\leq \|\mathbb{1}_{\{\bar{y} > \bar{t}\}} \nabla \bar{y}\|_{L^\infty(\Omega)} \|S'(\bar{u})v\|_{L^2(\Omega)} \|\nabla S'(\bar{u})v\|_{L^2(\Omega)} \\ &\leq 2\pi C^2 \sqrt{1 - m} \|v\|_{L^2(\Omega)}^2 \end{aligned}$$

for all  $v \in L^2(\Omega)$  and for some constant  $C$  independent of  $m$  and  $v$ . Here we have used the fact that  $\|S'(\bar{u})v\|_{H^1_0(\Omega)} \leq C \|v\|_{L^2(\Omega)}$  due to [Theorem 2.2](#). Combining this with (5.40) yields

$$\begin{aligned} Q(\bar{u}, \bar{y}, \bar{\varphi}; v) &\geq \frac{1}{2} \|S'(\bar{u})v\|_{L^2(\Omega)}^2 + \frac{\nu}{2} \|v\|_{L^2(\Omega)}^2 - 8C^2 \nu \pi^3 \sqrt{1 - m} \|v\|_{L^2(\Omega)}^2 \\ &\geq \frac{1}{2} \|S'(\bar{u})v\|_{L^2(\Omega)}^2 + \frac{\nu}{4} \|v\|_{L^2(\Omega)}^2 > 0 \end{aligned}$$

for all  $v \in L^2(\Omega)$ ,  $v \neq 0$ , provided that  $0 \leq 1 - m < \varepsilon$  with positive constant  $\varepsilon$  satisfying

$$\frac{1}{4} - 8C^2 \pi^3 \sqrt{\varepsilon} = 0.$$

$h_n$	$m = 1$		$m = 0.95$	
	$\ \bar{u} - \bar{u}_{h_n}\ _{L^2(\Omega)}$	$EOC_{L^2}(n)$	$\ \bar{u} - \bar{u}_{h_n}\ _{L^2(\Omega)}$	$EOC_{L^2}(n)$
$8.84 \cdot 10^{-2}$	$1.89 \cdot 10^{-1}$	1.9842	$1.66 \cdot 10^{-1}$	2.3015
$4.42 \cdot 10^{-2}$	$4.77 \cdot 10^{-2}$	1.9960	$3.37 \cdot 10^{-2}$	1.4995
$2.21 \cdot 10^{-2}$	$1.20 \cdot 10^{-2}$	1.9990	$1.19 \cdot 10^{-2}$	1.7032
$1.10 \cdot 10^{-2}$	$2.99 \cdot 10^{-3}$	1.9998	$3.66 \cdot 10^{-3}$	2.2945
$5.52 \cdot 10^{-3}$	$7.48 \cdot 10^{-4}$	1.9999	$7.46 \cdot 10^{-4}$	1.5713
$2.76 \cdot 10^{-3}$	$1.87 \cdot 10^{-4}$	2.0000	$2.51 \cdot 10^{-4}$	2.2772
$1.38 \cdot 10^{-3}$	$4.67 \cdot 10^{-5}$		$5.18 \cdot 10^{-5}$	

**Table 1:** discretization errors and experimental orders of convergence (EOC) in  $L^2$  for the optimal control  $\bar{u}$  in dependence of  $h_n$

We have therefore verified that all hypotheses of [Theorem 2.7](#) and of [Theorem 5.11](#) are fulfilled. Finally, [Lemma A.2](#) in the Appendix shows that  $\Sigma(\bar{y}) = 0$  if  $\bar{t} = m = 1$  and  $\Sigma(\bar{y}) > 0$  if  $\bar{t} = m \in (0, 1)$ .

We now consider the discrete approximation  $\bar{u}_h$  of  $\bar{u}$ . Thanks to [Remarks 5.5](#) and [5.6](#), we have  $\bar{u}_h = \text{Proj}_{[\alpha, \beta]}(-\frac{1}{v}\bar{\varphi}_h)$  with  $\text{Proj}_{[\alpha, \beta]}$  denoting the pointwise a.e. projection mapping onto the interval  $[\alpha, \beta]$ . From this, [\(3.1\)](#), and [\(5.10a\)](#),  $\bar{y}_h$  and  $\bar{\varphi}_h$  satisfy

$$(5.41) \quad \begin{cases} \int_{\Omega} (b(x) + \max\{\bar{y}_h - m, 0\}) \nabla \bar{y}_h \cdot \nabla w_h \, dx = \int_{\Omega} \text{Proj}_{[\alpha, \beta]}(-\frac{1}{v}\bar{\varphi}_h) w_h \, dx, \\ \int_{\Omega} (b(x) + \max\{\bar{y}_h - m, 0\}) \nabla \bar{\varphi}_h \cdot \nabla v_h + \mathbb{1}_{\{\bar{y}_h > m\}} \nabla \bar{y}_h \cdot \nabla \bar{\varphi}_h v_h \, dx = \int_{\Omega} (\bar{y}_h - y_d) v_h \, dx \end{cases}$$

for all  $w_h, v_h \in V_h$ . Since the nonlinearities of [\(5.41\)](#) are semi-smooth, it is reasonable to solve this system by a semi-smooth Newton (SSN) method; see, e.g., [\[38, 44\]](#) as well as [\[23\]](#). Setting  $y_0 := 128x_1x_2(1-x_1)(1-x_2)$ , we notice that the sets  $\{y_0 > \bar{t}\}$  and  $\{y_0 < \bar{t}\}$  have positive measures for all  $m \in (1-\varepsilon, 1]$ . The starting point for the discrete SSN method solved [\(5.41\)](#) is then taken as  $(y_{h,0}, \varphi_{h,0}) := (\text{Proj}_{V_h}(y_0), 0)$  for different mesh sizes  $h$ , where  $\text{Proj}_{V_h}$  stands for the  $L^2$  projection mapping onto  $V_h$ . The integrals over elements are approximated with a quadrature scheme. This introduces a variational crime which however does not reduce the expected approximation order for piecewise linear functions. In all our tests, the SSN method converged in four or five or six iterations.

We report the resulting discretization errors  $\|\bar{u}_h - \bar{u}\|_X$  for  $X = L^2(\Omega)$  and for  $h \in \{\frac{\sqrt{2}}{2^n} \mid 4 \leq n \leq 10\}$  as well as the experimental order of convergence

$$EOC_{L^p}(n) := \frac{\log(\|\bar{u} - \bar{u}_{h_{n+1}}\|_{L^p(\Omega)}) - \log(\|\bar{u} - \bar{u}_{h_n}\|_{L^p(\Omega)})}{\log(h_{n+1}) - \log(h_n)}$$

in [Table 1](#) for both cases  $m = 1$  and  $m = 0.95$ . For the situation  $m = 1$ , the results indicate an EOC of 2, which indicates that we are observing a superconvergence property; compare [\[40\]](#). The EOC for this case is consistent with the guaranteed rate of  $O(h^2)$  shown for the smooth problem in [\[17\]](#). An suitable explanation for this could be that in the case  $m = 1$ , we have the following identity

$$a(\bar{y}(x_1, x_2)) = \max\{\bar{y}(x_1, x_2) - 1, 0\} = 0 \quad \text{for all } (x_1, x_2) \in \Omega,$$

and thus the coefficient  $a(y)$  of the state equation in [\(5.39\)](#) is in fact smooth at the optimal state. When  $m = 0.95$  (and the coefficient is nonsmooth), the values of EOC are not stable, however their minimum value is approximately equal to 1.5, which fits the theoretical study shown in [Remark 5.12](#).

## 6 CONCLUSIONS

We have studied the numerical approximation of an optimal control problem governed by a quasilinear elliptic equation with nonsmooth coefficient in the divergence part. The convergence of a sequence of minimizers of some discrete control problems to a global minimizer of the original problem is shown. A priori error estimates for three types of discretizations (variational, piecewise constant, and continuous piecewise linear discretizations) are derived under an explicit second-order sufficient condition for the continuous optimal control problem and a structural assumption on the optimal state. The estimate for variational discretization corroborates the proven rate, although the observed rate is higher, which motivates follow-up work on rate optimality or superconvergence properties for optimal control of nonsmooth quasilinear equations.

## ACKNOWLEDGMENTS

This work was supported by the DFG under the grants CL 487/2-1 and RO 2462/6-1, both within the priority programme SPP 1962 “Nonsmooth and Complementarity-based Distributed Parameter Systems: Simulation and Hierarchical Optimization”. Part of this work was completed during a visit of the second author to the Vietnam Institute for Advanced Study in Mathematics (VIASM). The second author would like to thank the VIASM for their financial support and hospitality.

## APPENDIX A VERIFICATION OF A STRUCTURAL ASSUMPTION AND COMPUTATION OF THE JUMP FUNCTIONAL

**Lemma A.1.** *Let  $\Omega := \{x = (x_1, x_2) \in \mathbb{R}^2 \mid 0 < x_1, x_2 < 1\}$  and let  $y(x_1, x_2) := \sin(\pi x_1) \sin(\pi x_2)$ . Then there exists a constant  $c_s > 0$  such that*

$$(A.1) \quad \text{meas}_{\mathbb{R}^2} (\{|y - 1| < r\}) \leq c_s r$$

for all  $r > 0$  small enough.

*Proof.* Take any sufficiently small  $r > 0$  satisfying  $(1 - r, 1) \subset (0, 1)$ . A simple computation yields

$$\{|y - 1| < r\} = \left\{ \frac{1}{\pi} \arcsin \frac{1-r}{\sin \pi x_2} < x_1 < 1 - \frac{1}{\pi} \arcsin \frac{1-r}{\sin \pi x_2}, \frac{1}{\pi} \arcsin(1-r) < x_2 < 1 - \frac{1}{\pi} \arcsin(1-r) \right\}.$$

We thus obtain

$$(A.2) \quad \text{meas}_{\mathbb{R}^2} (\{|y - \bar{t}| < r\}) = \int_{\frac{1}{\pi} \arcsin(1-r)}^{1 - \frac{1}{\pi} \arcsin(1-r)} \left(1 - \frac{2}{\pi} \arcsin \frac{1-r}{\sin \pi x_2}\right) dx_2 \leq \left[1 - 2 \frac{1}{\pi} \arcsin(1-r)\right]^2,$$

where we have used the fact that

$$0 \leq 1 - \frac{2}{\pi} \arcsin \frac{1-r}{\sin \pi x_2} \leq 1 - 2 \frac{1}{\pi} \arcsin(1-r) \quad \text{for all } 0 \leq x_2 \leq 1, r \in (0, 1).$$

L'Hospital's rule then shows that

$$\lim_{r \rightarrow 0^+} \frac{1 - 2 \frac{1}{\pi} \arcsin(1-r)}{\sqrt{r}} = \frac{4}{\pi} \lim_{r \rightarrow 0^+} \sqrt{\frac{r}{1 - (1-r)^2}} = \frac{2\sqrt{2}}{\pi},$$

which, along with (A.2), yields (A.1). □

**Lemma A.2.** Let  $\Omega := \{x = (x_1, x_2) \in \mathbb{R}^2 \mid 0 < x_1, x_2 < 1\}$  and let  $y(x_1, x_2) := \sin(\pi x_1) \sin(\pi x_2)$ . Then

$$\Sigma(y) = \begin{cases} 8\sigma_0 \left(1 - \frac{2}{\pi} \arcsin \bar{t}\right) & \text{if } \bar{t} \in (0, 1), \\ 0 & \text{if } \bar{t} = 1, \\ 4\sigma_0 & \text{if } \bar{t} = 0. \end{cases}$$

*Proof.* We distinguish the following cases:

- (i) For any  $\bar{t} \in (0, 1)$  and any sufficiently small  $r > 0$  satisfying  $(\bar{t} - r, \bar{t} + r) \subset (0, 1)$ , a simple computation shows that

$$\begin{aligned} & \{|y - \bar{t}| < r, \partial_{x_1} y > 0\} \\ &= \left\{ \frac{1}{\pi} \arcsin \frac{\bar{t}-r}{\sin \pi x_2} < x_1 < \frac{1}{\pi} \arcsin \frac{\bar{t}+r}{\sin \pi x_2}, \frac{1}{\pi} \arcsin(\bar{t} + r) < x_2 < 1 - \frac{1}{\pi} \arcsin(\bar{t} + r) \right\} \\ & \cup \left\{ \frac{1}{\pi} \arcsin \frac{\bar{t}-r}{\sin \pi x_2} < x_1 < \frac{1}{2}, \frac{1}{\pi} \arcsin(\bar{t} - r) < x_2 < \frac{1}{\pi} \arcsin(\bar{t} + r) \right\} \\ & \cup \left\{ \frac{1}{\pi} \arcsin \frac{\bar{t}-r}{\sin \pi x_2} < x_1 < \frac{1}{2}, 1 - \frac{1}{\pi} \arcsin(\bar{t} + r) < x_2 < 1 - \frac{1}{\pi} \arcsin(\bar{t} - r) \right\}. \end{aligned}$$

We thus obtain

$$\begin{aligned} \int_{\Omega} \mathbb{1}_{\{|y-\bar{t}|<r, \partial_{x_1} y > 0\}} |\partial_{x_1} y| \, dx &= -\frac{2}{\pi} [\cos(\arcsin(\bar{t} + r)) - \cos(\arcsin(\bar{t} - r))] \\ & \quad - \frac{2}{\pi} (\bar{t} - r) [\arcsin(\bar{t} + r) - \arcsin(\bar{t} - r)] + 2r \left(1 - \frac{2}{\pi} \arcsin(\bar{t} + r)\right). \end{aligned}$$

Applying L'Hospital's rule yields

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-\bar{t}|<r, \partial_{x_1} y > 0\}} |\partial_{x_1} y| \, dx = 2 \left(1 - \frac{2}{\pi} \arcsin \bar{t}\right).$$

Similarly, there hold

$$\begin{aligned} \lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-\bar{t}|<r, \partial_{x_1} y < 0\}} |\partial_{x_1} y| \, dx &= 2 \left(1 - \frac{2}{\pi} \arcsin \bar{t}\right), \\ \lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-\bar{t}|<r, \partial_{x_2} y > 0\}} |\partial_{x_2} y| \, dx &= 2 \left(1 - \frac{2}{\pi} \arcsin \bar{t}\right), \\ \lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-\bar{t}|<r, \partial_{x_2} y < 0\}} |\partial_{x_2} y| \, dx &= 2 \left(1 - \frac{2}{\pi} \arcsin \bar{t}\right). \end{aligned}$$

By adding these four limits, we obtain

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-\bar{t}|<r\}} [|\partial_{x_1} y| + |\partial_{x_2} y|] \, dx = 8 \left(1 - \frac{2}{\pi} \arcsin \bar{t}\right).$$

- (ii) For  $\bar{t} = 0$  and for any  $r \in (0, 1)$  sufficiently small, we see from a straightforward calculation that

$$\begin{aligned} \{|y - 0| < r, \partial_{x_1} y > 0\} &= \left\{ 0 < x_1 < \frac{1}{\pi} \arcsin \frac{r}{\sin \pi x_2}, \frac{1}{\pi} \arcsin r < x_2 < 1 - \frac{1}{\pi} \arcsin r \right\} \\ & \cup \left\{ 0 < x_1 < \frac{1}{2}, 0 < x_2 < \frac{1}{\pi} \arcsin r \right\} \cup \left\{ 0 < x_1 < \frac{1}{2}, 1 - \frac{1}{\pi} \arcsin r < x_2 < 1 \right\}. \end{aligned}$$

Consequently, it holds that

$$\int_{\Omega} \mathbb{1}_{\{|y-0|<r, \partial_{x_1} y > 0\}} |\partial_{x_1} y| \, dx = -\frac{2}{\pi} [\cos(\arcsin r) - 1] + r \left(1 - \frac{2}{\pi} \arcsin r\right).$$

L'Hospital's rule then shows that

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-0| < r, \partial_{x_1} y > 0\}} |\partial_{x_1} y| \, dx = 1.$$

Similarly, we can conclude that

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-0| < r\}} [|\partial_{x_1} y| + |\partial_{x_2} y|] \, dx = 4.$$

(iii) For  $\bar{t} = 1$  and for any  $r > 0$  such that  $r \in (0, 1)$ , we have

$$\{|y - 1| < r, \partial_{x_1} y > 0\} = \left\{ \frac{1}{\pi} \arcsin \frac{1-r}{\sin \pi x_2} < x_1 < \frac{1}{2}, \frac{1}{\pi} \arcsin(1-r) < x_2 < 1 - \frac{1}{\pi} \arcsin(1-r) \right\}.$$

There therefore holds that

$$\int_{\Omega} \mathbb{1}_{\{|y-1| < r, \partial_{x_1} y > 0\}} |\partial_{x_1} y| \, dx = \frac{2}{\pi} [\cos(\arcsin(1-r)) + (1-r) \arcsin(1-r)] + r - 1.$$

Again, L'Hospital's rule shows that

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-1| < r, \partial_{x_1} y > 0\}} |\partial_{x_1} y| \, dx = 0.$$

Similarly, we can deduce that

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_{\Omega} \mathbb{1}_{\{|y-1| < r\}} [|\partial_{x_1} y| + |\partial_{x_2} y|] \, dx = 0. \quad \square$$

## APPENDIX B REGULARITY OF SOLUTIONS TO THE ADJUSTED LINEARIZED STATE EQUATION

Let  $p_0, p_1$ , and  $\gamma$  be given as in (4.18). In order to show the  $W^{1+\gamma, 2}$ -regularity of solutions to the adjusted linearized state equation (4.19), we need the following result for the function

$$\mathbb{f}_{\bar{u}} := \mathbb{1}_{\{\bar{y} \neq \bar{t}\}} a'(\bar{y}) \nabla \bar{y},$$

where  $\bar{y} := S(\bar{u})$  is the optimal state corresponding to the control  $\bar{u}$ .

**Proposition B.1.** *Assume that  $\bar{u} \in L^\infty(\Omega)$  satisfies  $\Sigma(\bar{y}) < \infty$ . Then there holds*

$$\mathbb{f}_{\bar{u}} \in (W^{\gamma, p_0}(\Omega))^2 = (W_0^{\gamma, p_0}(\Omega))^2.$$

*Proof.* Since  $\gamma < \frac{1}{p_0}$ , there holds that

$$W^{\gamma, p_0}(\Omega) = W_0^{\gamma, p_0}(\Omega);$$

see, e.g. Corollary 1.4.4.5 in [34]. Thanks to Theorem 2.2, one has  $\bar{y} \in W^{2, p_1}(\Omega) \hookrightarrow C^1(\bar{\Omega})$  and thus  $\bar{y} \in C^1(\bar{\Omega})$ . Therefore, we have  $\mathbb{f}_{\bar{u}} \in L^\infty(\Omega)^2 \hookrightarrow L^{p_0}(\Omega)^2$ . From the definition of the Sobolev spaces of fractional order; see e.g. Definition 6.8.2 in [39], it then suffices to show that

$$\iint_{\Omega \times \Omega} \frac{|\mathbb{f}_{\bar{u}}(x) - \mathbb{f}_{\bar{u}}(\hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} \, dx d\hat{x} < \infty,$$

or equivalently,

$$(B.1) \quad \iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} \frac{|\mathbb{f}_{\bar{u}}(x) - \mathbb{f}_{\bar{u}}(\hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} < \infty.$$

Here  $\delta$  is the constant in (2.9) and  $L$  denotes the Lipschitz constant of  $\bar{y}$ . For any  $x, \hat{x} \in \Omega$  satisfying  $|x - \hat{x}| < \frac{\delta}{L}$ , there holds that

$$(B.2) \quad |\bar{y}(x) - \bar{y}(\hat{x})| \leq L|x - \hat{x}| < \delta.$$

Setting now

$$(B.3) \quad Z(x, \hat{x}) := \mathbb{f}_{\bar{u}}(x) - \mathbb{f}_{\bar{u}}(\hat{x}) = \mathbb{1}_{\{\bar{y} \neq \bar{t}\}}(x) a'(\bar{y}(x)) \nabla \bar{y}(x) - \mathbb{1}_{\{\bar{y} \neq \bar{t}\}}(\hat{x}) a'(\bar{y}(\hat{x})) \nabla \bar{y}(\hat{x})$$

and using the argument in the proof of Lemma 3.4 for the situation  $y(x) := \bar{y}(x)$  and  $\hat{y}(x) := \bar{y}(\hat{x})$ , we can decompose  $Z(x, \hat{x})$  as follows

$$(B.4) \quad Z(x, \hat{x}) = Z^{(1)}(x, \hat{x}) + Z^{(2)}(x, \hat{x}) + Z^{(3)}(x, \hat{x}) + Z^{(4)}(x, \hat{x}),$$

where

$$(B.5) \quad |Z^{(1)}(x, \hat{x})| + |Z^{(2)}(x, \hat{x})| + |Z^{(4)}(x, \hat{x})| \leq C[|\bar{y}(x) - \bar{y}(\hat{x})| |\nabla \bar{y}(\hat{x})| + |\nabla(\bar{y}(x) - \bar{y}(\hat{x}))|],$$

$$(B.6) \quad Z^{(3)}(x, \hat{x}) = [a'_0(\bar{t}) - a'_1(\bar{t})][\chi_2(x, \hat{x}) - \chi_3(x, \hat{x})] \nabla \bar{y}(\hat{x}),$$

and

$$\chi_2(x, \hat{x}) = \begin{cases} 1 & \text{if } \bar{y}(\hat{x}) \in (\bar{t}, \bar{t} + \delta), \bar{y}(x) \in (\bar{t} - \delta, \bar{t}), \\ 0 & \text{otherwise,} \end{cases}$$

$$\chi_3(x, \hat{x}) = \begin{cases} 1 & \text{if } \bar{y}(\hat{x}) \in (\bar{t} - \delta, \bar{t}), \bar{y}(x) \in [\bar{t}, \bar{t} + \delta), \\ 0 & \text{otherwise.} \end{cases}$$

Here we have compared the function  $Z(x, \hat{x})$  with  $Z_{y, \hat{y}}$  defined in (3.6); compared the estimates for  $Z^{(i)}(x, \hat{x})$ ,  $i = 1, 2, 4$ , with the ones in (3.8); compared the decomposition (B.4) with (3.7); and compared the functions  $\chi_2, \chi_3$  with  $\mathbb{1}_{\Omega_{y, \hat{y}}^2}, \mathbb{1}_{\Omega_{y, \hat{y}}^3}$ , respectively, for the sets  $\Omega_{y, \hat{y}}^2, \Omega_{y, \hat{y}}^3$  given in (2.9). Applying (B.2) then yields

$$\iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} \frac{|\bar{y}(x) - \bar{y}(\hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} \leq \iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} L^{p_0} |x - \hat{x}|^{-2+p_0(1-\gamma)} dx d\hat{x} < \infty,$$

due to the fact that  $2 - p_0(1 - \gamma) < 2 = N$ ; see e.g. page 331 in [39]. Moreover, from the Sobolev embedding  $W^{1, p_1}(\Omega) \hookrightarrow C^{0, 1 - \frac{2}{p_1}}(\bar{\Omega})$ , there holds  $\nabla \bar{y} \in C^{0, 1 - \frac{2}{p_1}}(\bar{\Omega})^2$ . From this, we have

$$\iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} \frac{|\nabla \bar{y}(x) - \nabla \bar{y}(\hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} \leq \iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} C^{p_0} |x - \hat{x}|^{-2+p_0(1 - \frac{2}{p_1} - \gamma)} dx d\hat{x} < \infty,$$

in the view of (4.18). By combining the last two integral estimates with (B.5) and using the fact that  $\nabla \bar{y} \in C(\bar{\Omega})^2$ , there holds

$$(B.7) \quad \iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} \frac{|Z^{(i)}(x, \hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} < \infty \quad \text{for } i = 1, 2, 4.$$

We now estimate  $Z^{(3)}(x, \hat{x})$  defined in (B.6). To this end, we deduce from the definition of  $\chi_2(x, \hat{x})$  and  $\chi_3(x, \hat{x})$  that

$$\begin{aligned} |\chi_2(x, \hat{x}) - \chi_3(x, \hat{x})| &\leq \begin{cases} 1 & \text{if } 0 < |\bar{y}(\hat{x}) - \bar{t}| \leq |\bar{y}(\hat{x}) - \bar{y}(x)|, \\ 0 & \text{otherwise} \end{cases} \\ &\leq \begin{cases} 1 & \text{if } 0 < |\bar{y}(\hat{x}) - \bar{t}| \leq L|x - \hat{x}|, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Combining the last estimate with (B.6) yields

$$\begin{aligned} |Z^{(3)}(x, \hat{x})|^{p_0} &\leq C\sigma_0^{p_0} \mathbb{1}_{\{0 < |\bar{y}(\cdot) - \bar{t}| < L|\cdot - x|\}}(\hat{x}) |\nabla \bar{y}(\hat{x})|^{p_0} \\ &\leq C(p_0)\sigma_0^{p_0} \mathbb{1}_{\{0 < |\bar{y}(\cdot) - \bar{t}| < L|\cdot - x|\}}(\hat{x}) |\nabla \bar{y}(\hat{x})|, \end{aligned}$$

thanks to the boundedness of  $|\nabla \bar{y}|$  in  $C(\bar{\Omega})$ . On the other hand, we conclude from the definition of  $\Sigma(\bar{y})$  in (2.13) that there exists a constant  $r_0 > 0$  such that

$$\frac{\sigma_0}{r} \int_{\Omega} \mathbb{1}_{\{0 < |\bar{y}(\cdot) - \bar{t}| < r\}}(\hat{x}) |\nabla \bar{y}(\hat{x})| d\hat{x} \leq \Sigma(\bar{y}) + 1 \quad \text{for all } 0 < r < r_0.$$

We then have

$$\begin{aligned} &\iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < \delta/L\}} \frac{|Z^{(3)}(x, \hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} \\ &= \iint_{\Omega \times \Omega \cap \{r_0/L \leq |x - \hat{x}| < \delta/L\}} \frac{|Z^{(3)}(x, \hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} + \iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < r_0/L\}} \frac{|Z^{(3)}(x, \hat{x})|^{p_0}}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} \\ &\leq C \iint_{\Omega \times \Omega \cap \{r_0/L \leq |x - \hat{x}| < \delta/L\}} \frac{1}{|x - \hat{x}|^{2+p_0\gamma}} dx d\hat{x} \\ &\quad + C\sigma_0^{p_0} \iint_{\Omega \times \Omega \cap \{|x - \hat{x}| < r_0/L\}} \frac{1}{|x - \hat{x}|^{2+p_0\gamma}} \mathbb{1}_{\{0 < |\bar{y}(\cdot) - \bar{t}| < L|\cdot - x|\}}(\hat{x}) |\nabla \bar{y}(\hat{x})| d\hat{x} dx \\ &\leq C(r_0, \delta) + C\sigma_0^{p_0} \int_{\{\xi \in \mathbb{R}^2; |\xi| < \frac{r_0}{L}\}} \frac{1}{|\xi|^{2+p_0\gamma}} d\xi \int_{\Omega} \mathbb{1}_{\{0 < |\bar{y}(\cdot) - \bar{t}| < L|\xi|\}}(\hat{x}) |\nabla \bar{y}(\hat{x})| d\hat{x} \\ &= C(r_0, \delta) + CL\sigma_0^{p_0-1} \int_{\{\xi \in \mathbb{R}^2; |\xi| < \frac{r_0}{L}\}} \frac{d\xi}{|\xi|^{1+p_0\gamma}} \frac{\sigma_0}{L|\xi|} \int_{\Omega} \mathbb{1}_{\{0 < |\bar{y}(\cdot) - \bar{t}| < L|\xi|\}}(\hat{x}) |\nabla \bar{y}(\hat{x})| d\hat{x} \\ &\leq C(r_0, \delta) + CL\sigma_0^{p_0-1} \frac{2\pi}{1-p_0\gamma} \left(\frac{r_0}{L}\right)^{1-p_0\gamma} (\Sigma(\bar{y}) + 1) \\ &< \infty, \end{aligned}$$

due to (4.18). From this, (B.7), (B.4), and (B.3), we have (B.1).  $\square$

**Lemma B.2.** *Assume that  $\bar{u} \in L^\infty(\Omega)$  satisfies  $\Sigma(\bar{y}) < \infty$ . Then for any  $z \in H^1(\Omega)$ , there holds*

$$\operatorname{div}(\mathbb{f}_{\bar{u}} z) \in W^{-1+\gamma, 2}(\Omega).$$

Moreover,

$$(B.8) \quad \|\operatorname{div}(\mathbb{f}_{\bar{u}} z)\|_{W^{-1+\gamma, 2}(\Omega)} \leq C \|\mathbb{f}_{\bar{u}}\|_{W^{\gamma, p_0}(\Omega)} \|z\|_{H^1(\Omega)}$$

for some constant  $C$  independent of  $z$ .

*Proof.* By Proposition B.1, one has  $\mathbb{f}_{\bar{u}} \in (W^{\gamma, p_0}(\Omega))^2$ . We now apply the multiplication theorem for Sobolev spaces; see, e.g. Theorem 1.4.4.2 in [34] and Theorem 7.4 in [4] for  $s_1 = \gamma$ ,  $s_2 = 1$ ,  $s = \gamma$ ,  $p_1 = p_0 > 2$ ,  $p_2 = 2$ ,  $p = 2$ , and  $n = 2$  to obtain

$$\mathbb{f}_{\bar{u}} z \in (W^{\gamma, 2}(\Omega))^2.$$

From this and the continuity of the divergence operator from  $(W^{\gamma, 2}(\Omega))^2$  to  $W^{-1+\gamma, 2}(\Omega)$  for  $\gamma \neq \frac{1}{2}$ ; see, e.g. Theorem 1.4.4.6 in [34], we have  $\operatorname{div}(\mathbb{f}_{\bar{u}} z) \in W^{-1+\gamma, 2}(\Omega)$ . Finally, we also have (B.8) from Theorem 7.4 in [4] and Theorem 1.4.4.6 in [34].  $\square$

**Proposition B.3.** *The following assertions hold:*

(i) *Let  $p > 2$  be fixed but arbitrary. Then, for any  $u \in L^2(\Omega)$  and  $v \in W^{-1, p}(\Omega)$ , (4.19) admits a unique solution  $\tilde{z}_{u, v} \in W_0^{1, p}(\Omega)$ . Furthermore, there exists a constant  $\rho_p > 0$  such that*

$$(B.9) \quad \|\tilde{z}_{u, v}\|_{W_0^{1, p}(\Omega)} \leq C \|v\|_{W^{-1, p}(\Omega)} \quad \text{for all } u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho_p)$$

*and for some constant  $C$  independent of  $u$  and  $v$ .*

(ii) *Assume that  $\bar{u} \in L^\infty(\Omega)$  satisfies  $\Sigma(\bar{y}) < \infty$ . Then, for any  $u \in \mathcal{U}_{ad}$  and any  $v \in W^{-1+\gamma, 2}(\Omega)$ , the equation (4.19) admits a unique solution  $\tilde{z}_{u, v}$  in  $W^{1+\gamma, 2}(\Omega)$ . Moreover, there exists a constant  $\hat{\rho}$  such that*

$$(B.10) \quad \|\tilde{z}_{u, v}\|_{W^{1+\gamma, 2}(\Omega)} \leq C(1 + \|\mathbb{f}_{\bar{u}}\|_{W^{\gamma, p_0}(\Omega)}) \|v\|_{W^{-1+\gamma, 2}(\Omega)}$$

*for all  $u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \hat{\rho}) \cap \mathcal{U}_{ad}$  and for some constant  $C$  independent of  $u$  and  $v$ .*

*Proof.* Ad (i): For any  $u \in L^2(\Omega) \hookrightarrow W^{-1, p}(\Omega)$ , by Theorem 2.2, we have  $y_u = S(u) \in W_0^{1, p}(\Omega)$ . From this, and Remark 2.9 in [15], we deduce the existence and uniqueness of  $\tilde{z}_{u, v}$  in  $H_0^1(\Omega)$  to (4.19).

We now rewrite (4.19) as follows

$$\begin{cases} -\operatorname{div}[(b + a(y_u))\nabla \tilde{z}_{u, v} + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \tilde{z}_{u, v} \nabla y_u] = v - \operatorname{div}(Z_{y_u, \bar{y}} \tilde{z}_{u, v}) & \text{in } \Omega, \\ \tilde{z}_{u, v} = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $Z_{y_u, \bar{y}}$  is defined as in (3.6). In other words, one has

$$(B.11) \quad \tilde{z}_{u, v} = S'(u)(v - \operatorname{div}(Z_{y_u, \bar{y}} \tilde{z}_{u, v})),$$

due to Theorem 2.2. By Lemma 3.4, we have that  $Z_{y_u, \bar{y}} \in L^p(\Omega)^2$ . Since  $\tilde{z}_{u, v} \in H_0^1(\Omega) \hookrightarrow L^q(\Omega)$  with  $q > \frac{2p}{p-2}$ , we then deduce that  $Z_{y_u, \bar{y}} \tilde{z}_{u, v} \in L^r(\Omega)^2$  with  $r = \frac{pq}{p+q} \in (2, p)$ . There therefore holds

$$v - \operatorname{div}(Z_{y_u, \bar{y}} \tilde{z}_{u, v}) \in W^{-1, r}(\Omega).$$

Applying Theorem 2.2 yields  $\tilde{z}_{u, v} \in W_0^{1, r}(\Omega) \hookrightarrow C(\bar{\Omega})$ . We then have  $Z_{y_u, \bar{y}} \tilde{z}_{u, v} \in L^p(\Omega)^2$  and thus

$$v - \operatorname{div}(Z_{y_u, \bar{y}} \tilde{z}_{u, v}) \in W^{-1, p}(\Omega).$$

This, together with Theorem 2.2, gives  $\tilde{z}_{u, v} \in W_0^{1, p}(\Omega)$ .

We now prove (B.9). To this end, by Theorem 2.2 and the compact embedding  $L^2(\Omega) \Subset W^{-1, p}(\Omega)$ , we first have

$$\sup\{\|S'(u)\|_{\mathbb{L}(W^{-1, p}(\Omega), W_0^{1, p}(\Omega))} \mid u \in \bar{B}_{L^2(\Omega)}(\bar{u}, 1)\} \leq C_1,$$

which gives for all  $u \in \overline{B}_{L^2(\Omega)}(\bar{u}, 1)$  that

$$(B.12) \quad \begin{aligned} \|\tilde{z}_{u,v}\|_{W_0^{1,p}(\Omega)} &\leq C_1 \|v - \operatorname{div}(Z_{y_u, \bar{y}} \tilde{z}_{u,v})\|_{W^{-1,p}(\Omega)} \leq C_1 [\|v\|_{W^{-1,p}(\Omega)} + \|Z_{y_u, \bar{y}} \tilde{z}_{u,v}\|_{L^p(\Omega)}] \\ &\leq C_1 [\|v\|_{W^{-1,p}(\Omega)} + \|\tilde{z}_{u,v}\|_{L^{2p/(p-2)}(\Omega)} \|Z_{y_u, \bar{y}}\|_{L^2(\Omega)}] \\ &\leq C_1 [\|v\|_{W^{-1,p}(\Omega)} + \|\tilde{z}_{u,v}\|_{W_0^{1,p}(\Omega)} \|Z_{y_u, \bar{y}}\|_{L^2(\Omega)}], \end{aligned}$$

where we have just used the Hölder inequality and the embeddings  $W_0^{1,p}(\Omega) \hookrightarrow H_0^1(\Omega) \hookrightarrow L^{2p/p-2}(\Omega)$ . From Lemma 3.4 and the fact that  $\|y_u - \bar{y}\|_{W_0^{1,p}(\Omega)} \rightarrow 0$  as  $\|u - \bar{u}\|_{W^{-1,p}(\Omega)} \leq C \|u - \bar{u}\|_{L^2(\Omega)} \rightarrow 0$  (see Theorem 2.2), we have

$$\|Z_{y_u, \bar{y}}\|_{L^2(\Omega)} \rightarrow 0 \quad \text{as} \quad \|u - \bar{u}\|_{L^2(\Omega)} \rightarrow 0.$$

Then, there exists a constant  $\rho_p \in (0, 1]$  such that

$$\|Z_{y_u, \bar{y}}\|_{L^2(\Omega)} \leq \frac{1}{2C_1} \quad \text{for all } u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho_p).$$

Combining this with (B.12) yields

$$\|\tilde{z}_{u,v}\|_{W_0^{1,p}(\Omega)} \leq 2C_1 \|v\|_{W^{-1,p}(\Omega)} \quad \text{for all } u \in \overline{B}_{L^2(\Omega)}(\bar{u}, \rho_p).$$

*Ad (ii):* We first show the  $W^{1+\gamma,2}$ -regularity of  $\tilde{z}_{u,v}$ . To this end, we now rewrite (4.19) as follows

$$\begin{cases} -\operatorname{div}[(b + a(y_u)) \nabla \tilde{z}_{u,v}] = v + \operatorname{div}[\mathbb{f}_{\bar{u}} \tilde{z}_{u,v}] & \text{in } \Omega, \\ \tilde{z}_{u,v} = 0 & \text{on } \partial\Omega \end{cases}$$

or, equivalently,

$$(B.13) \quad \begin{cases} -\Delta \tilde{z}_{u,v} = \frac{1}{b + a(y_u)} (\nabla b + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \nabla y_u) \cdot \nabla \tilde{z}_{u,v} + \frac{1}{b + a(y_u)} [v + \operatorname{div}(\mathbb{f}_{\bar{u}} \tilde{z}_{u,v})] & \text{in } \Omega, \\ \tilde{z}_{u,v} = 0 & \text{on } \partial\Omega. \end{cases}$$

From Proposition B.1 and Lemma B.2, there holds  $\operatorname{div}[\mathbb{f}_{\bar{u}} \tilde{z}_{u,v}] \in W^{-1+\gamma,2}(\Omega)$  and thus

$$(B.14) \quad v + \operatorname{div}[\mathbb{f}_{\bar{u}} \tilde{z}_{u,v}] \in W^{-1+\gamma,2}(\Omega).$$

Since  $\mathcal{U}_{ad}$  is bounded in  $L^\infty(\Omega)$  and thus in  $L^{p_0}(\Omega)$ , we deduce from Theorem 2.2 that  $y_u, \bar{y} \in W^{2,p_0}(\Omega) \hookrightarrow C^1(\overline{\Omega})$  and then that

$$\|y_u\|_{W^{2,p}(\Omega)} + \|y_u\|_{C^1(\overline{\Omega})} \leq M \quad \text{for all } u \in \mathcal{U}_{ad}$$

for some constant  $M > 0$ . Here  $p_0$  is the constant in (4.18). Thanks to Assumption (A2) and Assumption (A3), one has  $b + a(y_u), (b + a(y_u))^{-1} \in W^{1,\infty}(\Omega)$  and

$$(B.15) \quad \|b + a(y_u)\|_{W^{1,\infty}(\Omega)} + \|(b + a(y_u))^{-1}\|_{W^{1,\infty}(\Omega)} \leq C_M \quad \text{for all } u \in \mathcal{U}_{ad}.$$

Combining this with (B.14) yields

$$\frac{1}{b + a(y_u)} [v + \operatorname{div}[\mathbb{f}_{\bar{u}} \tilde{z}_{u,v}]] \in W^{-1+\gamma,2}(\Omega),$$

where we have employed the fact that the multiplication bilinear mapping  $W^{1,\infty}(\Omega) \times W^{-1+\gamma,2}(\Omega) \rightarrow W^{-1+\gamma,2}(\Omega)$  is continuous; see, e.g. [4]. The second term in the right hand side of (B.13) then belongs to  $W^{-1+\gamma,2}(\Omega)$ . Moreover, the first one also belongs to  $W^{-1+\gamma,2}(\Omega)$ , since it is in  $L^2(\Omega) \hookrightarrow W^{-1+\gamma,2}(\Omega)$ .

We therefore apply the  $W^{1+\gamma,2}$ -regularity of solutions to (B.13) to have that  $\tilde{z}_{u,v} \in W^{1+\gamma,2}(\Omega)$ ; see, e.g. Theorem 3 in [43] and Theorem 4.1 in [21].

It remains to show the existence of a constant  $\hat{\rho} > 0$  satisfying (B.10). To this end, we first see from (4.18) that  $1 > \frac{2}{p_0} \geq 1 - \gamma$  and  $1 < \frac{p_0}{p_0-1} < 2$ . From this and the Sobolev embedding theorem; see e.g. Theorem 3.8 in [4], we deduce that  $W_0^{1, \frac{p_0}{p_0-1}}(\Omega) \hookrightarrow W_0^{\frac{2}{p_0}, 2}(\Omega) \hookrightarrow W_0^{1-\gamma, 2}(\Omega)$  and thus

$$W^{-1+\gamma, 2}(\Omega) \hookrightarrow W^{-1, p_0}(\Omega).$$

We now apply assertion (i) for  $p := p_0$  to derive that  $\tilde{z}_{u,v} \in W_0^{1, p_0}(\Omega)$  and that

$$(B.16) \quad \|\tilde{z}_{u,v}\|_{W_0^{1, p_0}(\Omega)} \leq C\|v\|_{W^{-1, p_0}(\Omega)} \leq C\|v\|_{W^{-1+\gamma, 2}(\Omega)} \quad \text{for all } u \in \bar{B}_{L^2(\Omega)}(\bar{u}, \rho_{p_0}) \cap \mathcal{U}_{ad}.$$

Moreover, by applying Theorem 3 in [43] (see, also Theorem 4.1 in [21]) to (B.13), and using estimates (B.8) and (B.15), as well as the embedding  $W_0^{1, p_0}(\Omega) \hookrightarrow H_0^1(\Omega)$ , there is a constant  $C = C(M)$  such that

$$\begin{aligned} \|\tilde{z}_{u,v}\|_{W^{1+\gamma, 2}(\Omega)} &\leq C(M) [\|v\|_{W^{-1+\gamma, 2}(\Omega)} + (1 + \|f_{\bar{u}}\|_{W^{\gamma, p_0}(\Omega)}) \|\tilde{z}_{u,v}\|_{H_0^1(\Omega)}] \\ &\leq C(M) [\|v\|_{W^{-1+\gamma, 2}(\Omega)} + (1 + \|f_{\bar{u}}\|_{W^{\gamma, p_0}(\Omega)}) \|\tilde{z}_{u,v}\|_{W_0^{1, p_0}(\Omega)}] \quad \text{for all } u \in \mathcal{U}_{ad}. \end{aligned}$$

Setting now  $\hat{\rho} := \rho_{p_0}$  and combining the last inequality with (B.16) yields (B.10).  $\square$

## REFERENCES

- [1] G. Alberti, S. Bianchini, and G. Crippa, Structure of level sets and Sard-type properties of Lipschitz maps, *Ann. Sc. Norm. Super. Pisa Cl. Sci.(5)* XII (2013), 863–902, [doi:10.2422/2036-2145.201107\\_006](https://doi.org/10.2422/2036-2145.201107_006).
- [2] N. Arada, E. Casas, and F. Tröltzsch, Error estimates for the numerical approximation of a semilinear elliptic control problem, *Comput. Optim. Appl.* 23 (2002), 201–229, [doi:10.1023/a:1020576801966](https://doi.org/10.1023/a:1020576801966).
- [3] T. Bayen, J. F. Bonnans, and F. J. Silva, Characterization of local quadratic growth for strong minima in the optimal control of semi-linear elliptic equations, *Trans. Amer. Math. Soc.* 366 (2014), 2063–2087, [doi:10.1090/S0002-9947-2013-05961-2](https://doi.org/10.1090/S0002-9947-2013-05961-2).
- [4] A. Behzadan and M. Holst, Multiplication in Sobolev spaces, revisited, *Ark. Mat.* 59 (2021), 275–306, [doi:10.4310/arkiv.2021.v59.n2.a2](https://doi.org/10.4310/arkiv.2021.v59.n2.a2).
- [5] A. Bejan, *Convection Heat Transfer*, J. Wiley & Sons, 4 edition, 2013, [doi:10.1002/9781118671627](https://doi.org/10.1002/9781118671627).
- [6] J. F. Bonnans and A. Shapiro, *Perturbation Analysis of Optimization Problems*, Springer-Verlag, Berlin, Heidelberg, 2000, [doi:10.1007/978-1-4612-1394-9](https://doi.org/10.1007/978-1-4612-1394-9).
- [7] S. C. Brenner and L. R. Scott, *The Mathematical Theory of Finite Element Methods*, Texts in Applied Mathematics, Springer-Verlag, New York, 2008, [doi:10.1007/978-0-387-75934-0](https://doi.org/10.1007/978-0-387-75934-0).
- [8] C. Carstensen, Quasi-interpolation and a posteriori error analysis in finite element methods, *ESAIM: Mathematical Modelling and Numerical Analysis* 33 (1999), 1187–1202, [doi:10.1051/m2an:1999140](https://doi.org/10.1051/m2an:1999140).
- [9] E. Casas and V. Dhano, Error estimates for the numerical approximation of a quasilinear Neumann problem under minimal regularity of the data, *Numer. Math.* 117 (2011), 115–145, [doi:10.1007/s00211-010-0344-1](https://doi.org/10.1007/s00211-010-0344-1).

- [10] E. Casas and V. Dharmo, Error estimates for the numerical approximation of Neumann control problems governed by a class of quasilinear elliptic equations, *Comput. Optim. Appl.* 52 (2012), 719–756, [doi:10.1007/s10589-011-9440-0](https://doi.org/10.1007/s10589-011-9440-0).
- [11] E. Casas and M. Mateos, Uniform convergence of the FEM. Applications to state constrained control problems, *Comput. Appl. Math.* 21 (2002), 67–100.
- [12] E. Casas, M. Mateos, and J. P. Raymond, Error estimates for the numerical approximation of a distributed control problem for the steady-state Navier-Stokes equations, *SIAM J. Control Optim.* 46 (2007), 952–982, [doi:10.1137/060649999](https://doi.org/10.1137/060649999).
- [13] E. Casas, M. Mateos, and A. Rösch, Numerical approximation of control problems of non-monotone and non-coercive semilinear elliptic equations, *Numerische Mathematik* 149 (2021), 305–340, [doi:10.1007/s00211-021-01222-7](https://doi.org/10.1007/s00211-021-01222-7).
- [14] E. Casas, M. Mateos, and F. Tröltzsch, Error estimates for the numerical approximation of boundary semilinear elliptic control problems, *Comput. Optim. Appl.* 31 (2005), 193–219, [doi:10.1007/s10589-005-2180-2](https://doi.org/10.1007/s10589-005-2180-2).
- [15] E. Casas and F. Tröltzsch, First- and second-order optimality conditions for a class of optimal control problems with quasilinear elliptic equations, *SIAM J. Control Optim.* 48 (2009), 688–718, [doi:10.1137/080720048](https://doi.org/10.1137/080720048).
- [16] E. Casas and F. Tröltzsch, Numerical analysis of some optimal control problems governed by a class of quasilinear elliptic equations, *ESAIM: COCV* 17 (2011), 771–800, [doi:10.1051/cocv/2010025](https://doi.org/10.1051/cocv/2010025).
- [17] E. Casas and F. Tröltzsch, A general theorem on error estimates with application to a quasilinear elliptic optimal control problem, *Comput. Optim. Appl.* 53 (2012), 173–206, [doi:10.1007/s10589-011-9453-8](https://doi.org/10.1007/s10589-011-9453-8).
- [18] M. Chipot, *Elliptic Equations: An Introductory Course*, Birkhäuser Verlag, Basel, 2009, [doi:10.1007/978-3-7643-9982-5](https://doi.org/10.1007/978-3-7643-9982-5).
- [19] C. Christof and C. Meyer, A note on a priori  $L^p$ -error estimates for the obstacle problem, *Numerische Mathematik* 139 (2018), 27–45, [doi:10.1007/s00211-017-0931-5](https://doi.org/10.1007/s00211-017-0931-5).
- [20] P. G. Ciarlet, *The Finite Element Method for Elliptic Problems*, Classics in Applied Mathematics, SIAM, 2002, [doi:10.1137/1.9780898719208](https://doi.org/10.1137/1.9780898719208).
- [21] S. Clain, Elliptic operators of divergence type with Hölder coefficients in fractional Sobolev spaces, *Rend. Mat. Appl.* 17 (1997), 207–236.
- [22] C. Clason, V. H. Nhu, and A. Rösch, No-gap second-order optimality conditions for optimal control of a non-smooth quasilinear elliptic equation, *ESAIM: COCV* 27 (2021), 62, [doi:10.1051/cocv/2020092](https://doi.org/10.1051/cocv/2020092).
- [23] C. Clason, V. H. Nhu, and A. Rösch, Optimal control of a non-smooth quasilinear elliptic equation, *Mathematical Control and Related Fields* 11 (2021), 521–554, [doi:10.3934/mcrf.2020052](https://doi.org/10.3934/mcrf.2020052).
- [24] C. Clason, V. H. Nhu, and A. Rösch, Numerical analysis of a nonsmooth quasilinear elliptic control problem: I. Explicit second-order optimality conditions, *submitted to MzAN* (2023), [arXiv:2203.16865](https://arxiv.org/abs/2203.16865).
- [25] B. Dacorogna, *Direct Methods in the Calculus of Variations*, Applied Mathematical Sciences, Springer, 2 edition, 2008, [doi:10.1007/978-0-387-55249-1](https://doi.org/10.1007/978-0-387-55249-1).

- [26] J. C. De Los Reyes and V. Dharmo, Error estimates for optimal control problems of a class of quasilinear equations arising in variable viscosity fluid flow, *Numer. Math.* 132 (2016), 691–720, [doi:10.1007/s00211-015-0737-2](https://doi.org/10.1007/s00211-015-0737-2).
- [27] J. C. De Los Reyes, C. Meyer, and B. Vexler, Finite element error analysis for state-constrained optimal control of the Stokes equations, *Control Cybern.* 37 (2008), 251–284.
- [28] K. Deckelnick and M. Hinze, Convergence of a finite element approximation to a state-constrained elliptic control problem, *SIAM J. Numer. Anal.* 45 (2007), 1937–1953, [doi:10.1137/060652361](https://doi.org/10.1137/060652361).
- [29] T. Dupont and R. Scott, Polynomial approximation of functions in Sobolev spaces, *Mathematics of Computation* 34 (1980), 441–463.
- [30] L. C. Evans and R. F. Gariepy, *Measure Theory and Fine Properties of Function*, CRC Press, New York, 4th edition, 1992, [doi:10.1201/b18333](https://doi.org/10.1201/b18333).
- [31] R. S. Falk, Approximation of a class of optimal control problems with order of convergence estimates, *J. Math. Anal. Appl.* 44 (1973), 28–47, [doi:10.1016/0022-247X\(73\)90022-X](https://doi.org/10.1016/0022-247X(73)90022-X).
- [32] T. Geveci, On the approximation of the solution of an optimal control problem governed by an elliptic equation, *RAIRO. Anal. Numér.* 13 (1979), 313–328, [doi:10.1051/m2an/1979130403131](https://doi.org/10.1051/m2an/1979130403131).
- [33] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, Heidelberg, 2001, [doi:10.1007/978-3-642-61798-0](https://doi.org/10.1007/978-3-642-61798-0).
- [34] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman Advanced Pub. Program, 1985, [doi:10.1137/1.9781611972030](https://doi.org/10.1137/1.9781611972030).
- [35] M. Hinze, A variational discretization concept in control constrained optimization: The linear-quadratic case, *Comput. Optim. Appl.* 30 (2005), 45–61, [doi:10.1007/s10589-005-4559-5](https://doi.org/10.1007/s10589-005-4559-5).
- [36] M. Hinze and F. Tröltzsch, Discrete concepts versus error analysis in PDE-constrained optimization, *GAMM-Mitteilungen* 33 (2010), 148–162, [doi:10.1002/gamm.201010012](https://doi.org/10.1002/gamm.201010012).
- [37] M. Holtmannspötter and A. Rösch, A priori error estimates for the finite element approximation of a nonsmooth optimal control problem governed by a coupled semilinear PDE-ODE system, *SIAM J. Control Optim.* 59 (2021), 3329–3358, [doi:10.1137/20m1331263](https://doi.org/10.1137/20m1331263).
- [38] K. Ito and K. Kunisch, *Lagrange Multiplier Approach to Variational Problems and Applications*, Advances in Design and Control, SIAM, 2008, [doi:10.1137/1.9780898718614](https://doi.org/10.1137/1.9780898718614).
- [39] A. Kufner, O. John, and S. Fučík, *Function Spaces*, Noordhoff International Publishing, Leyden; Academia, Prague, 1977. Monographs and Textbooks on Mechanics of Solids and Fluids; Mechanics: Analysis.
- [40] C. Meyer and A. Rösch, Superconvergence Properties of Optimal Control Problems, *SIAM Journal on Control and Optimization* 43 (2004), 970–985, [doi:10.1137/s0363012903431608](https://doi.org/10.1137/s0363012903431608).
- [41] C. Meyer and O. Thoma, A priori finite element error analysis for optimal control of the obstacle problem, *SIAM J. Numer. Anal.* 51 (2013), 605–628, [doi:10.1137/110836092](https://doi.org/10.1137/110836092).
- [42] I. Neitzel, J. Pfefferer, and A. Rösch, Finite element discretization of state-constrained elliptic optimal control problems with semilinear state equation, *SIAM J. Control Optim.* 53 (2015), 874–904, [doi:10.1137/140960645](https://doi.org/10.1137/140960645).

- [43] G. Savaré, Regularity results for Elliptic equations in Lipschitz domains, *Journal of Functional Analysis* 152 (1998), 176–201, doi:10.1006/jfan.1997.3158, <https://www.sciencedirect.com/science/article/pii/S002212369793158X>.
- [44] M. Ulbrich, *Semismooth Newton Methods for Variational Inequalities and Constrained Optimization Problems in Function Spaces*, MOS-SIAM Series on Optimization, SIAM, 2011, doi:10.1137/1.9781611970692.
- [45] Y. B. Zel'dovich and Y. P. Raizer, *Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena*, Academic Press, 1966, doi:10.1115/1.3607836.