## MATH3195/5195M EXERCISE SHEET 1

## DUE: NOON, FEBRUARY 8, 2019

Warm-up exercises

**Problem 1.** Work out the example from the first lecture: What are the integer solutions of  $x^2 + y^2 = z^2$ ?

(1) First show that the rational solutions of  $X^2 + Y^2 = 1$  are of the form

$$(X,Y) = \left(\frac{-2m}{1+m^2}, \frac{1-m^2}{1+m^2}\right), m \in \mathbb{Q}, \text{ and } (X,Y) = (0,0).$$

(2) From this find the integer solutions of the original equation.

**Problem 2.** Let  $\mathbb{T} = (\mathbb{R} \cup \{\infty\}, \oplus, \odot)$  with addition defined as  $x \oplus y := \min(x, y)$  and multiplication  $x \odot y := x + y$  for all  $x, y \in \mathbb{R} \cup \{\infty\}$ .

- (a) Is  $\mathbb{T}$  a commutative ring? If yes, then show that all axioms hold, if no, then explain which axiom fails.
- (b) Calculate  $3 \odot (5 \oplus 7)$ ,  $(3 \oplus -3)^2$ , and  $(1 \oplus 8)^4$ .
- (c) Show that for any  $x, y \in \mathbb{R} \cup \{\infty\}$ , and any  $k \in \mathbb{N}$ , one has  $(x \oplus y)^k = x^k \oplus y^k$ .

**Problem 3.** (a) Prove that if  $\varphi : R \to S$  is a ring isomorphism then  $\varphi^{-1} : S \to R$  is a ring homomorphism, and hence also an isomorphism.

(b) Let *R* be a ring and  $I \subseteq R$  be an ideal and let  $\varphi : R \to R/I$  be the canonical projection. Show that ker  $\varphi = I$  and  $\varphi$  is a ring homomorphism.

**Problem 4.** Let *I*, *J* and *K* be ideals of a ring *R*. Show that

- (a)  $I \cap J$  and IJ are ideals (b)  $IJ \neq I \cap J$ ,
- (c) I(J + K) = IJ + IK,

**Problem 5.** Let *I*, *J* and *K* be ideals of a ring *R*. Recall that  $(I : J) = \{r \in R : rJ \subset I\}$ . Show that

- (a) (I : J) is an ideal of *R* and  $I \subset (I : J)$ ,
- (b)  $J \subset I$  implies that (I : J) = R,
- (c)  $IJ \subset K$  if and only if  $I \subset (K : J)$ .

**Problem 6.** Let *R* be a commutative ring and let  $I, J \subseteq R$  be ideals.

- (a) Let  $\sqrt{I} = \{r \in R : r^n \in I \text{ for some positive integer } n\}$ . Show that  $\sqrt{I}$  is an ideal that contains *I*. [Note:  $\sqrt{I}$  is called the *radical of I*.]
- (b) Prove that  $\sqrt{I \cap J} = \sqrt{I} \cap \sqrt{J}$ .
- (c) Let R = k[x, y]. Show that  $\sqrt{(x^2, y^2)} = (x, y)$  and that  $\sqrt{(x^2) \cap (y^2)} = (xy)$ .