

UNIFORM CONVERGENCE TO EQUILIBRIUM FOR A FAMILY OF DRIFT-DIFFUSION MODELS WITH TRAP-ASSISTED RECOMBINATION AND SELF-CONSISTENT POTENTIAL

KLEMENS FELLNER AND MICHAEL KNIELY

ABSTRACT. We investigate a recombination–drift–diffusion model coupled to Poisson’s equation modelling the transport of charge within certain types of semiconductors. In more detail, we study a two-level system for electrons and holes endowed with an intermediate energy level for electrons occupying trapped states. As our main result, we establish an explicit functional inequality between relative entropy and entropy production, which leads to exponential convergence to equilibrium. We stress that our approach is applied uniformly in the lifetime of electrons on the trap level assuming that this lifetime is sufficiently small.

1. INTRODUCTION AND MAIN RESULTS

We consider the following PDE–ODE recombination–drift–diffusion system coupled to Poisson’s equation on a bounded domain $\Omega \subset \mathbb{R}^3$ with boundary $\partial\Omega \in C^2$:

$$(1) \quad \begin{cases} \partial_t n = \nabla \cdot J_n(n, \psi) + R_n(n, n_{tr}), \\ \partial_t p = \nabla \cdot J_p(p, \psi) + R_p(p, n_{tr}), \\ \varepsilon \partial_t n_{tr} = R_p(p, n_{tr}) - R_n(n, n_{tr}), \\ -\lambda \Delta \psi = n - p + \varepsilon n_{tr} - D, \end{cases}$$

where the flux terms J_n , J_p and recombination terms R_n , R_p are defined as

$$\begin{aligned} J_n &:= \nabla n + n \nabla(\psi + V_n) = \mu_n \nabla \frac{n}{\mu_n} + n \nabla \psi, & \mu_n &:= e^{-V_n}, \\ J_p &:= \nabla p + p \nabla(-\psi + V_p) = \mu_p \nabla \frac{p}{\mu_p} - p \nabla \psi, & \mu_p &:= e^{-V_p}, \\ R_n &:= \frac{1}{\tau_n} \left(n_{tr} - \frac{n}{n_0 \mu_n} (1 - n_{tr}) \right), & R_p &:= \frac{1}{\tau_p} \left(1 - n_{tr} - \frac{p}{p_0 \mu_p} n_{tr} \right). \end{aligned}$$

The variables n , p , and n_{tr} denote the densities of electrons in the conduction band, holes in the valence band, and electrons on the trap level (see Fig. 1). Moreover, ψ represents the electrostatic potential generated by n , p , n_{tr} , and the time-independent doping profile $D \in L^\infty(\Omega)$. The constants $n_0, p_0, \tau_n, \tau_p > 0$ are positive recombination parameters and $\varepsilon \in (0, \varepsilon_0]$ for arbitrary but fixed $\varepsilon_0 > 0$ is a positive relaxation parameter in the sense that system (1) reduces in the limit $\varepsilon = 0$ to the famous Shockley–Read–Hall model of electron recombination [16, 20] in semiconductor drift–diffusion systems, see e.g. [15, 17]. Finally,

$$(2) \quad V_n, V_p \in W^{2,\infty}(\Omega) \quad \text{with} \quad \hat{n} \cdot \nabla V_n, \hat{n} \cdot \nabla V_p \geq 0 \quad \text{on } \partial\Omega$$

represent external time-independent potentials.

1991 *Mathematics Subject Classification.* Primary 35Q81; Secondary 78A35, 35B40, 35K57.

Key words and phrases. Semiconductor model, recombination–drift–diffusion systems, self-consistent potential, trapped states, entropy method, convergence to equilibrium, exponential rate of convergence.




FIGURE 1. An illustration of the allowed transitions of electrons between the three energy levels.

The system is equipped with no-flux boundary conditions for the flux components and homogeneous Neumann boundary data for the electrostatic potential,

$$(3) \quad \hat{n} \cdot J_n = \hat{n} \cdot J_p = \hat{n} \cdot \nabla \psi = 0 \quad \text{on } \partial\Omega$$

where \hat{n} represents the outer unit normal vector on $\partial\Omega$. As we will use a result in [15] for proving existence and uniqueness of global solutions, we choose the initial conditions $n(0, \cdot) = n_I$, $p(0, \cdot) = p_I$, and $n_{tr}(0, \cdot) = n_{tr,I}$ in accordance with [15]:

$$(4) \quad n_I, p_I \in H^1(\Omega) \cap L^\infty(\Omega), \quad n_I, p_I \geq 0, \quad 0 \leq n_{tr,I} \leq 1.$$

For convenience, we assume that the volume of Ω is normalised, i.e. $|\Omega| = 1$, and we set

$$\bar{f} := \int_{\Omega} f(x) dx$$

for any function $f \in L^1(\Omega)$. This abbreviation is consistent with the usual definition of the average of f since $|\Omega| = 1$.

The main goal of the paper is to close the gap between the models investigated in [8] and [9]. While the pure Shockley–Read–Hall model including the electrostatic potential has been considered in [8], the family of drift–diffusion models with trap-assisted recombination already appeared in [9] but without coupling to Poisson’s equation. Here, we focus on the exponential convergence to equilibrium for a PDE–ODE model including both trapped states dynamics and the self-consistent potential. More precisely, we obtain an explicit bound for the convergence rate by employing the so-called *entropy method* which amounts to deriving a functional inequality between an entropy functional and the associated entropy dissipation. For related works on exponential convergence to equilibrium for reaction–diffusion systems, see e.g. [6, 10, 11, 18, ?AMT00]. The framework of the entropy method which we shall use here to obtain explicit bounds on the convergence rate originates from [3–5], where models from reversible chemistry have been studied. An earlier application of the entropy method, but using a non-constructive compactness argument, is presented in [13, 14], where the authors prove exponential convergence for a model of electrically charged species taking the coupling to Poisson’s equation into account.

Remark 1.1. Throughout the article, we will frequently encounter the following inhomogeneous Poisson equation with right hand side $f \in L^2(\Omega)$ subject to homogeneous Neumann boundary conditions:

$$(5) \quad -\lambda \Delta \psi = f \quad \text{in } \Omega, \quad \hat{n} \cdot \nabla \psi = 0 \quad \text{on } \partial\Omega.$$

It is well-known that there exists a weak solution $\psi \in H^1(\Omega)$ if and only if $\bar{f} = 0$ holds true (compatibility condition with homogeneous Neumann boundary data). In this case, ψ is determined only up to an additive constant, which one can fix via the normalisation $\bar{\psi} = 0$ to obtain a unique solution ψ .

Due to Remark 1.1, we additionally have to demand that the initial data satisfy the charge-neutrality condition

$$(6) \quad \int_{\Omega} (n_I - p_I + \varepsilon n_{tr,I} - D) dx = 0.$$

As a consequence of the structure of system (1) and the no-flux boundary conditions in (3), we see that the total charge is preserved for all $t \geq 0$ in the sense that

$$(7) \quad \int_{\Omega} (n(t, \cdot) - p(t, \cdot) + \varepsilon n_{tr}(t, \cdot)) dx = \int_{\Omega} (n_I - p_I + \varepsilon n_{tr,I}) dx = \int_{\Omega} D dx.$$

Definition 1.2. A *global weak solution* to (1)–(4) and (6) is a quadruple $(n, p, n_{tr}, \psi) : [0, \infty) \rightarrow H^1(\Omega)^2 \times L^\infty(\Omega) \times H^1(\Omega)$ such that for all $T > 0$ the following conditions are satisfied:

- $n, p \in L^2(0, T; H^1(\Omega))$, and
- $$\begin{aligned}
 & - \int_0^T \langle u'(t), n(t) \rangle_{H^1(\Omega)^* \times H^1(\Omega)} dt - \int_{\Omega} n_I u(0) dx \\
 & \quad = - \int_0^T \int_{\Omega} J_n(n, \psi) \cdot \nabla u dx dt + \int_0^T \int_{\Omega} R_n(n, n_{tr}) u dx dt, \\
 & - \int_0^T \langle u'(t), p(t) \rangle_{H^1(\Omega)^* \times H^1(\Omega)} dt - \int_{\Omega} p_I u(0) dx \\
 & \quad = - \int_0^T \int_{\Omega} J_p(p, \psi) \cdot \nabla u dx dt + \int_0^T \int_{\Omega} R_p(p, n_{tr}) u dx dt,
 \end{aligned}$$
- for all $u \in W_2(0, T) := \{f \in L^2(0, T; H^1(\Omega)) \mid \partial_t f \in L^2(0, T; H^1(\Omega)^*)\}$ subject to $u(T) = 0$,
- $n_{tr}(T) = n_{tr,I} + \frac{1}{\varepsilon} \int_0^T (R_p(p, n_{tr}) - R_n(n, n_{tr})) dt$,
- $\lambda \int_{\Omega} \nabla \psi(T) \cdot \nabla w dx = \int_{\Omega} (n(T) - p(T) + \varepsilon n_{tr}(T) - D) w dx$ for all $w \in H^1(\Omega)$.

We further mention the embedding $W_2(0, T) \hookrightarrow C([0, T], L^2(\Omega))$ known from PDE theory (see e.g. [2]).

Proposition 1.3 (Global Solutions). *There exists a unique global weak solution $(n, p, n_{tr}, \psi) : [0, \infty) \rightarrow H^1(\Omega)^2 \times L^\infty(\Omega) \times H^1(\Omega)$ of (1)–(4) and (6) with $\psi(t, \cdot) = 0$ for all $t \geq 0$. This solution satisfies $n, p \in L^2(0, T; H^1(\Omega))$ for all $T > 0$ uniformly in $\varepsilon \in (0, \varepsilon_0]$ as well as (7). Moreover, $n, p \geq 0$, $0 \leq n_{tr} \leq 1$, and there exist positive constants $M, K(M)$ (again uniformly in $\varepsilon \in (0, \varepsilon_0]$) such that*

$$(8) \quad \|n(t)\|_{L^\infty(\Omega)} + \|p(t)\|_{L^\infty(\Omega)} \leq M \quad \text{and} \quad \|\psi(t)\|_{H^2(\Omega)} + \|\psi(t)\|_{C(\bar{\Omega})} \leq K$$

for all $t \geq 0$. In addition, there exists a positive constant $\mu(M, K)$ (uniformly in $\varepsilon \in (0, \varepsilon_0]$) such that

$$(9) \quad n(t, x), p(t, x) \geq \mu \min \{t^2, 1\}$$

and

$$(10) \quad n_{tr}(t, x) \in \left[\min \left\{ \frac{1}{2\varepsilon_0\tau_p} t, \mu \right\}, 1 - \min \left\{ \frac{1}{2\varepsilon_0\tau_n} t, \mu \right\} \right]$$

for all $t \geq 0$ and a.e. $x \in \Omega$. Finally, $n, p \in W_2(0, T) \hookrightarrow C([0, T], L^2(\Omega))$, $n_{tr} \in C([0, T], L^\infty(\Omega))$, and $\psi \in C([0, T], H^2(\Omega))$, where each inclusion holds true uniformly in $\varepsilon \in (0, \varepsilon_0]$.

Proposition 1.4 (Equilibrium States). *The stationary system*

$$(11) \quad \begin{cases} \nabla \cdot J_n(n) + R_n(n, n_{tr}) = 0, \\ \nabla \cdot J_p(p) + R_p(p, n_{tr}) = 0, \\ R_n(n, n_{tr}) = R_p(p, n_{tr}), \\ -\lambda \Delta \psi = n - p + \varepsilon n_{tr} - D, \end{cases}$$

subject to $\hat{n} \cdot J_n = \hat{n} \cdot J_p = \hat{n} \cdot \nabla \psi = 0$ admits a unique solution $(n_\infty, p_\infty, n_{tr,\infty}, \psi_\infty) \in (H^1(\Omega) \cap L^\infty(\Omega))^4$ satisfying $\overline{\psi_\infty} = 0$. The equilibrium potential ψ_∞ is continuous and there exists a positive constant K_∞ such that

$$\|\psi_\infty\|_{H^2(\Omega)} + \|\psi_\infty\|_{C(\overline{\Omega})} \leq K_\infty.$$

Moreover, there exist positive constants $M_\infty(K_\infty)$, $\mu_\infty(K_\infty)$ such that

$$(12) \quad n_\infty(x), p_\infty(x) \in (\mu_\infty, M_\infty) \quad \text{and} \quad n_{tr,\infty}(x) \in (\mu_\infty, 1 - \mu_\infty)$$

for a.e. $x \in \Omega$. The constants K_∞ , M_∞ , and μ_∞ are independent of $\varepsilon \in (0, \varepsilon_0]$. In detail, the equilibrium densities n_∞ , p_∞ , and $n_{tr,\infty}$ read

$$(13) \quad n_\infty = n_* e^{-\psi_\infty - V_n}, \quad p_\infty = p_* e^{\psi_\infty - V_p}, \quad n_{tr,\infty} = \frac{n_*}{n_* + n_0 e^{\psi_\infty}} = \frac{p_0}{p_0 + p_* e^{\psi_\infty}}$$

where the positive constants n_* and p_* are uniquely determined in terms of ψ_∞ by

$$(14) \quad n_* p_* = n_0 p_0 \quad \text{and} \quad n_* \overline{e^{-\psi_\infty - V_n}} - p_* \overline{e^{\psi_\infty - V_p}} + \varepsilon \frac{n_*}{n_* + n_0 e^{\psi_\infty}} - \overline{D} = 0.$$

Furthermore, the following relations hold true:

$$(15) \quad n_{tr,\infty} = \frac{n_\infty(1 - n_{tr,\infty})}{n_0 \mu_n} \quad \text{and} \quad 1 - n_{tr,\infty} = \frac{p_\infty n_{tr,\infty}}{p_0 \mu_p}.$$

We introduce the *entropy functional* $E(n, p, n_{tr}, \psi)$ for non-negative functions $n, p, n_{tr} \in L^2(\Omega)$ satisfying $n_{tr} \leq 1$ and $\bar{n} - \bar{p} + \varepsilon \bar{n}_{tr} = \overline{D}$ where $\psi \in H^1(\Omega)$ is the unique solution of (5) with right hand side $f = n - p + \varepsilon n_{tr} - D$ and normalisation $\overline{\psi} = 0$:

$$(16) \quad E(n, p, n_{tr}, \psi) := \int_\Omega \left(n \ln \frac{n}{n_0 \mu_n} - (n - n_0 \mu_n) + p \ln \frac{p}{p_0 \mu_p} - (p - p_0 \mu_p) + \frac{\lambda}{2} |\nabla \psi|^2 + \varepsilon \int_{1/2}^{n_{tr}} \ln \left(\frac{s}{1-s} \right) ds \right) dx.$$

The densities n and p enter via Boltzmann entropy contributions $a \ln a - (a-1) \geq 0$, whereas n_{tr} appears within the entropy functional via an integral term. We first mention that the integral $\int_{1/2}^{n_{tr}} \ln \left(\frac{s}{1-s} \right) ds$ is non-negative and finite for all $n_{tr}(x) \in [0, 1]$. In more detail, we may write

$$\begin{aligned} \int_{1/2}^{n_{tr}} \ln \left(\frac{s}{1-s} \right) ds &= [n_{tr} \ln n_{tr} - (n_{tr} - 1)] \\ &\quad + [(1 - n_{tr}) \ln(1 - n_{tr}) - ((1 - n_{tr}) - 1)] + \ln 2 - 1. \end{aligned}$$

Consequently, both the occupied and unoccupied trapped states (n_{tr} and $1 - n_{tr}$) are described via Boltzmann statistics within the entropy functional, and the integral $\int_{1/2}^{n_{tr}} \ln \left(\frac{s}{1-s} \right) ds$ allows to combine the contributions of n_{tr} and $1 - n_{tr}$ in a compact fashion.

One can further verify that the entropy functional (16) is indeed a Lyapunov functional: By defining the *entropy production functional*

$$(17) \quad P(n, p, n_{tr}, \psi) := -\frac{d}{dt} E(n, p, n_{tr}, \psi),$$

we first calculate along solutions of (1) that formally

$$(18) \quad P(n, p, n_{tr}, \psi) = \int_{\Omega} \left(\frac{|J_n|^2}{n} + \frac{|J_p|^2}{p} - R_n \ln \left(\frac{n(1 - n_{tr})}{n_0 \mu_n n_{tr}} \right) - R_p \ln \left(\frac{p n_{tr}}{p_0 \mu_p (1 - n_{tr})} \right) \right) dx.$$

The entropy production functional involves non-negative flux terms as well as recombination terms of the form $(a - 1) \ln a \geq 0$. The entropy production P is, therefore, a non-negative functional, which ensures the monotone decrease in time of the entropy E along trajectories of (1). More precisely, it will be shown in Theorem 1.6 that the global weak solutions to (1) obtained in Proposition 1.3 satisfy a suitable weak version of (17), see (19) below.

The following theorem constitutes a so-called entropy–entropy production (EEP) estimate. This is a functional inequality between entropy and entropy production for arbitrary, yet admissible non-negative functions $n, p, n_{tr} \in L^\infty(\Omega)$, $n_{tr} \leq 1$; in particular, the electrostatic potential $\psi \in H^1(\Omega)$ in the following theorem must be the unique solution of (5) subject to $f = n - p + \varepsilon n_{tr} - D$ and the normalisation $\bar{\psi} = 0$.

Theorem 1.5. *Consider all non-negative functions $n, p, n_{tr} \in L^\infty(\Omega)$ subject to $n, p \leq \mathcal{M}$, $n_{tr} \leq 1$, and $\bar{n} - \bar{p} + \varepsilon \bar{n}_{tr} = \bar{D}$ and accordingly determine $\psi \in H^1(\Omega)$ as the unique solution to (5) with $f = n - p + \varepsilon n_{tr} - D$ and $\bar{\psi} = 0$. Then, there exist explicit constants $\varepsilon_0 > 0$ and $C_{EEP} > 0$ depending on \mathcal{M} and on K_∞ (as given in Proposition 1.4) such that*

$$E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \leq C_{EEP} P(n, p, n_{tr}, \psi)$$

holds true for all $\varepsilon \in (0, \varepsilon_0]$.

Note that C_{EEP} is independent of $\varepsilon \in (0, \varepsilon_0]$ and that this abstract EEP inequality can be applied to the global solution to (1) by using $\mathcal{M} = M$ from Proposition 1.3. We are then able to prove the exponential decay of the entropy relative to the equilibrium by using a Gronwall argument.

Theorem 1.6. *Let $\varepsilon \in (0, \varepsilon_0]$ with $\varepsilon_0 > 0$ from Theorem 1.5, and let (n, p, n_{tr}, ψ) be the unique global weak solution to (1) with non-negative initial datum $(n_I, p_I, n_{tr, I}) \in (H^1(\Omega) \cap L^\infty(\Omega))^2 \times L^\infty(\Omega)$, $n_{tr, I} \leq 1$, satisfying $\bar{n}_I - \bar{p}_I + \varepsilon \bar{n}_{tr, I} = \bar{D}$ according to Proposition 1.3. In addition, let $(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty)$ be the unique equilibrium state characterised in Proposition 1.4 as a solution to (11). Then, (n, p, n_{tr}, ψ) fulfils the weak entropy production law*

$$(19) \quad E(n, p, n_{tr}, \psi)(t_1) + \int_{t_0}^{t_1} P(n, p, n_{tr}, \psi)(s) ds = E(n, p, n_{tr}, \psi)(t_0)$$

for all $0 < t_0 \leq t_1 < \infty$. As a consequence, $E(n, p, n_{tr}, \psi)$ converges exponentially to $E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty)$ with explicit rate and constant as a function of time $t \geq 0$. More precisely,

$$(20) \quad E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \leq \left(E(n_I, p_I, n_{tr, I}, \psi_I) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \right) e^{-C_{EEP}^{-1} t}$$

where $\psi_I \in H^1(\Omega)$ is the unique weak solution to (5) with $f = n_I - p_I + \varepsilon n_{tr, I} - D$ and $\bar{\psi}_I = 0$.

Corollary 1.7. *Under the hypotheses of Theorem 1.6, the following improved convergence properties with constants $0 < c, C < \infty$ both depending on M and K_∞ but not on $\varepsilon \in (0, \varepsilon_0]$ hold true for all $t \geq 0$:*

$$(21) \quad \|n - n_\infty\|_{L^\infty(\Omega)} + \|p - p_\infty\|_{L^\infty(\Omega)} + \|n_{tr} - n_{tr,\infty}\|_{L^\infty(\Omega)} + \|\psi - \psi_\infty\|_{H^2(\Omega)} \leq Ce^{-ct}.$$

In particular, $\psi \rightarrow \psi_\infty$ in $L^\infty(\Omega)$ at an exponential rate.

The remainder of this article is devoted to the proofs of the various statements above. Section 2 collects the proofs of Propositions 1.3 and 1.4. The proof of Theorem 1.5 along with the necessary prerequisites is contained in Section 3, while the results on exponential convergence to equilibrium are proven in Section 4.

2. GLOBAL SOLUTION AND EQUILIBRIUM STATE

Proof of Proposition 1.3. The existence of such a unique global solution (n, p, n_{tr}, ψ) as well as $(n, p) \in L^2(0, T; H^1(\Omega))$ for all $T > 0$ uniformly for $\varepsilon \in (0, \varepsilon_0]$ and $0 \leq n_{tr} \leq 1$ are a consequence of [15, Lemma 3.1]. The uniform-in-time L^∞ bounds for n and p follow similar to [7, Lemma 4.1], where a Nash–Moser-type iteration for L^r norms, $r \geq 1$, of n and p has been employed. But as the coupling to Poisson’s equation is missing in [7], we have to slightly modify the line of arguments.

The evolution of the L^{r+1} norm, $r \geq 1$, of n and p can be reformulated as

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} (n^{r+1} + p^{r+1}) dx &= (r+1) \int_{\Omega} \left(-rn^{r-1} \nabla n \cdot (\nabla n + n \nabla(\psi + V_n)) \right. \\ &\quad \left. - rp^{r-1} \nabla p \cdot (\nabla p + p \nabla(-\psi + V_p)) + n^r R_n + p^r R_p \right) dx \\ &\leq -\frac{4r}{r+1} \int_{\Omega} \left| \nabla n^{\frac{r+1}{2}} \right|^2 dx - \frac{4r}{r+1} \int_{\Omega} \left| \nabla p^{\frac{r+1}{2}} \right|^2 dx + r \int_{\Omega} n^{r+1} \Delta(\psi + V_n) dx \\ &\quad + r \int_{\Omega} p^{r+1} \Delta(-\psi + V_p) dx + (r+1) \int_{\Omega} (n^r R_n + p^r R_p) dx. \end{aligned}$$

To the last term, we apply the estimate $(r+1)n^r \leq \frac{1}{r} + 2rn^{r+1}$ which follows from Young’s inequality $ab \leq \frac{1}{q}a^q + \frac{1}{s}b^s$ with $a := (\frac{1}{r+1})^{\frac{1}{r+1}}$, $b := (\frac{1}{r+1})^{-\frac{1}{r+1}}n^r$, $q := r+1$, and $s := \frac{r+1}{r}$. As a consequence of $-\lambda \Delta \psi = n - p + \varepsilon n_{tr} - D$, $(n^{r+1} - p^{r+1})(n - p) \geq 0$, $|R_n| \leq C(1+n)$, and $|R_p| \leq C(1+p)$, we then deduce

$$(22) \quad \begin{aligned} \frac{d}{dt} \int_{\Omega} (n^{r+1} + p^{r+1}) dx &\leq -\frac{4r}{r+1} \int_{\Omega} \left| \nabla n^{\frac{r+1}{2}} \right|^2 dx - \frac{4r}{r+1} \int_{\Omega} \left| \nabla p^{\frac{r+1}{2}} \right|^2 dx \\ &\quad + \widehat{C}r \int_{\Omega} (n^{r+1} + p^{r+1}) dx + \frac{\widehat{C}}{r} \end{aligned}$$

with a constant $\widehat{C} > 0$ depending on ε_0 but not on r . One can now proceed as in [7, Lemma 4.1]. For completeness, we briefly collect the main arguments below and refer to [7] for the details. By utilizing the Gagliardo–Nirenberg-type inequality $\|f\|_{L^2(\Omega)} \leq C_{GN} \|f\|_{L^1(\Omega)}^{\frac{2}{5}} \|f\|_{H^1(\Omega)}^{\frac{3}{5}}$ for $f := n^{\frac{r+1}{2}}$ and $f := p^{\frac{r+1}{2}}$, one derives

$$(23) \quad \begin{aligned} \int_{\Omega} (n^{r+1} + p^{r+1}) dx &\leq \delta \int_{\Omega} \left(\left| \nabla n^{\frac{r+1}{2}} \right|^2 + \left| \nabla p^{\frac{r+1}{2}} \right|^2 \right) dx + \frac{\widetilde{C}}{\delta} \left(\int_{\Omega} (n^{\frac{r+1}{2}} + p^{\frac{r+1}{2}}) dx \right)^2 \end{aligned}$$

where $\tilde{C} > 0$ is a constant independent of r and $\delta > 0$. We now introduce $\lambda_k := 2^k - 1$ for $k \geq 1$ and set $r := \lambda_k$. Choosing a sufficiently small constant $A > 0$ and defining $\delta_k := \frac{A}{\lambda_k}$ results in

$$\delta_k(\hat{C}\lambda_k + \delta_k) \leq \frac{4\lambda_k}{\lambda_k + 1}$$

for all $k \geq 1$. By multiplying (23) with $\hat{C}\lambda_k + \delta_k$ and by combining the result with (22), we arrive at

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} (n^{\lambda_k+1} + p^{\lambda_k+1}) dx &\leq -\delta_k \int_{\Omega} (n^{\lambda_k+1} + p^{\lambda_k+1}) dx \\ &\quad + B\lambda_k(\lambda_k + \delta_k) \sup_{0 \leq \tau \leq t} \left(\int_{\Omega} \left(n^{\frac{r+1}{2}} + p^{\frac{r+1}{2}} \right) dx \right)^2 + \frac{\hat{C}}{\lambda_k} \end{aligned}$$

with the constant $B := \frac{\hat{C}\tilde{C}}{A}$. The uniform L^∞ bounds on n and p now follow from [7, Lemma 4.2].

As, in particular, $\|n(t) - p(t) + \varepsilon n_{tr}(t) - D\|_{L^2(\Omega)}$ is uniformly bounded in $t \geq 0$, we conclude that $\|\psi(t)\|_{H^2(\Omega)}$ is uniformly bounded in time by applying standard elliptic regularity theory (see e.g. [19, Chap. IV. §2. Theorem 4]). The announced bound on $\|\psi(t)\|_{C(\bar{\Omega})}$ follows from the embedding $H^2(\Omega) \hookrightarrow C(\bar{\Omega})$ valid in \mathbb{R}^3 .

The regularity $\partial_t n, \partial_t p \in L^2(0, T; H^1(\Omega)^*)$ and, hence, $n, p \in W_2(0, T) \hookrightarrow C([0, T], L^2(\Omega))$ uniformly for $\varepsilon \in (0, \varepsilon_0]$ is easily inferred from the corresponding bounds on $J_n, J_p \in L^2((0, T) \times \Omega)$ and $R_n, R_p \in L^\infty((0, T) \times \Omega)$. Likewise, $n_{tr} \in C([0, T], L^\infty(\Omega))$ and $\psi \in C([0, T], H^2(\Omega))$, where both inclusions hold true uniformly for $\varepsilon \in (0, \varepsilon_0]$.

For showing the upper and lower bound on n_{tr} in (10), we multiply the third equation in (1) with τ_p and observe that

$$\varepsilon \partial_t (\tau_p n_{tr}) \geq 1 - r n_{tr}$$

holds true with a constant $r(M) > 1$ due to $\|p\|_{L^\infty(\Omega)} \leq M$. We now distinguish the following three cases for all $t \geq 0$ and a.e. $x \in \Omega$: $n_{tr}(t, x) \geq \frac{1}{r}$, $n_{tr}(t, x) \in [\frac{1}{2r}, \frac{1}{r})$, and $n_{tr}(t, x) < \frac{1}{2r}$. In the first case, $\partial_t (\tau_p n_{tr}(t, x)) \leq 0$, while in the second case $\partial_t (\tau_p n_{tr}(t, x)) > 0$. And in the third case, $\partial_t (\tau_p n_{tr}(t, x)) > \frac{1}{2\varepsilon_0}$. Defining $t_0 := \frac{\varepsilon_0 \tau_p}{r}$, this ensures

$$(24) \quad \tau_p n_{tr}(t, x) \geq \frac{t}{2\varepsilon_0}, \quad t \in [0, t_0], \quad \text{and} \quad n_{tr}(t, x) \geq \frac{1}{2r}, \quad t \geq t_0.$$

The upper bound on n_{tr} follows by applying the same arguments to $\tau_n(1 - n_{tr})$.

Concerning the bounds on n and p in (9), we follow the lines in [9] and concentrate on the arguments for n as the result for p can be derived analogously. We begin with defining $w := e^{\psi + V_n} n$ which fulfils $\hat{n} \cdot \nabla w = 0$ on $\partial\Omega$. For simplicity, we also set w.l.o.g. $\tau_n = \tau_p = 1$ in the following calculations. The evolution of w on Ω is governed by

$$\begin{aligned} \partial_t w &= \Delta w - \nabla(\psi + V_n) \cdot \nabla w + e^{\psi + V_n} n_{tr} - \frac{1 - n_{tr}}{n_0 \mu_n} w \\ (25) \quad &\geq \Delta w - \nabla(\psi + V_n) \cdot \nabla w - \alpha w + \beta n_{tr} \end{aligned}$$

with constants $\alpha > 0$ and $\beta(K) > 0$. Employing the lower bounds on n_{tr} from (24), we first test (25) with $(w - \mu_1 t^2)_-$ for $t \in [0, t_0]$ where $\mu_1 > 0$ is a constant

specified below and where we abbreviate $(\cdot)_- := \min\{\cdot, 0\}$. This entails

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_{\Omega} (w - \mu_1 t^2)_-^2 dx &= \int_{\Omega} (w - \mu_1 t^2)_- (\partial_t w - 2\mu_1 t) dx \\ &\leq \int_{\Omega} (w - \mu_1 t^2)_- (\Delta w - \nabla(\psi + V_n) \cdot \nabla w - \alpha w + \beta n_{tr} - 2\mu_1 t) dx \\ &\leq - \int_{\Omega} \mathbb{1}_{w \leq \mu_1 t^2} |\nabla w|^2 dx - \frac{1}{2} \int_{\Omega} \nabla \left[(w - \mu_1 t^2)_-^2 \right] \cdot \nabla(\psi + V_n) dx \\ &\quad + \int_{\Omega} (w - \mu_1 t^2)_- (-\alpha w + \beta n_{tr} - 2\mu_1 t) dx \end{aligned}$$

where we also utilized $\hat{n} \cdot \nabla V_n \geq 0$ on $\partial\Omega$ from (2). Due to the bounds $t \leq t_0$ and $n_{tr}(t, x) \geq \frac{t}{2\varepsilon_0}$, we obtain

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_{\Omega} (w - \mu_1 t^2)_-^2 dx &\leq \frac{1}{2} \int_{\Omega} \Delta(\psi + V_n) (w - \mu_1 t^2)_-^2 dx \\ &\quad + \int_{\Omega} (w - \mu_1 t^2)_- \left(-\alpha \mu_1 t_0 + \frac{\beta}{2\varepsilon_0} - 2\mu_1 \right) t dx. \end{aligned}$$

Choosing $\mu_1 > 0$ according to $\mu_1(2 + \alpha t_0) \leq \frac{\beta}{2\varepsilon_0}$, we deduce

$$\frac{d}{dt} \int_{\Omega} (w - \mu_1 t^2)_-^2 dx \leq \|\Delta(\psi + V_n)\|_{L^\infty(\Omega)} \int_{\Omega} (w - \mu_1 t^2)_-^2 dx.$$

Because of $\int_{\Omega} (w(0, x))_-^2 dx = 0$, we derive $\int_{\Omega} (w - \mu_1 t^2)_-^2 dx = 0$ for all $t \in [0, t_0]$ by applying a Gronwall argument. We thus arrive at $w(t, x) \geq \mu_1 t^2$ for all $t \in [0, t_0]$ and a.e. $x \in \Omega$.

In the situation $t \geq t_0$, we test (25) with $(w - \mu_2)_-$ where $\mu_2 > 0$ is another constant to be specified. As above, we recall (2) and calculate

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_{\Omega} (w - \mu_2)_-^2 dx &= \int_{\Omega} (w - \mu_2)_- \partial_t w dx \\ &\leq \int_{\Omega} (w - \mu_2)_- (\Delta w - \nabla(\psi + V_n) \cdot \nabla w - \alpha w + \beta n_{tr}) dx \\ &\leq - \int_{\Omega} \mathbb{1}_{w \leq \mu_2} |\nabla w|^2 dx - \frac{1}{2} \int_{\Omega} \nabla \left[(w - \mu_2)_-^2 \right] \cdot \nabla(\psi + V_n) dx \\ &\quad + \int_{\Omega} (w - \mu_2)_- (-\alpha w + \beta n_{tr}) dx. \end{aligned}$$

Here, we have $n_{tr}(t, x) \geq \frac{1}{2r}$, which yields

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_{\Omega} (w - \mu_2)_-^2 dx &\leq \frac{1}{2} \int_{\Omega} \Delta(\psi + V_n) (w - \mu_2)_-^2 dx \\ &\quad + \int_{\Omega} (w - \mu_2)_- \left(-\alpha \mu_2 + \frac{\beta}{2r} \right) dx. \end{aligned}$$

If we impose the conditions $\alpha \mu_2 \leq \frac{\beta}{2r}$ and $\mu_2 \leq \mu_1 t_0^2$ on $\mu_2 > 0$, we infer

$$\frac{d}{dt} \int_{\Omega} (w - \mu_2)_-^2 dx \leq \|\Delta(\psi + V_n)\|_{L^\infty(\Omega)} \int_{\Omega} (w - \mu_2)_-^2 dx$$

as well as $\int_{\Omega} (w(t_0, x) - \mu_2)_-^2 dx = 0$. Gronwall's lemma now leads to $\int_{\Omega} (w - \mu_2)_-^2 dx = 0$ and, hence, $w(t, x) \geq \mu_2$ for all $t \geq t_0$ and a.e. $x \in \Omega$. \square

Proof of Proposition 1.4. As the entropy production vanishes at the stationary state $(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty)$, straightforward calculations show that $J_n = J_p = R_n = R_p = 0$ yields the representations for n_∞ and p_∞ as well as the two expressions for

$n_{tr,\infty}$ in (13). The relation $n_*p_* = n_0p_0$ results from a combination of the formulas arising from $R_n = 0$ and $R_p = 0$, while the fact that the conservation law is also fulfilled in the equilibrium follows from integrating Poisson's equation in (11). Note that n_* (and hence p_*) is uniquely determined from the second relation in (14) as its left hand side is strictly monotonously increasing and surjective from $(0, \infty)$ to $(-\infty, \infty)$ as a function of n_* . The identities in (15) are equivalent versions of $R_n(n_\infty, n_{tr,\infty}) = 0$ and $R_p(p_\infty, n_{tr,\infty}) = 0$.

Next, we establish the existence of the limiting potential ψ_∞ . A technical difficulty stems from the fact that the constant n_* (or equally p_*) depends non-locally on ψ_∞ , see (14). However, this can be avoided by substituting $\widetilde{\psi}_\infty := \psi_\infty - \ln n_*$ and rewriting $-\lambda \Delta \psi_\infty = n_\infty - p_\infty + \varepsilon n_{tr,\infty} - D$ as

$$(26) \quad -\lambda \Delta \widetilde{\psi}_\infty - e^{-\widetilde{\psi}_\infty - V_n} + n_0 p_0 e^{\widetilde{\psi}_\infty - V_p} - \frac{\varepsilon}{1 + n_0 e^{\widetilde{\psi}_\infty}} = -D.$$

We now aim to apply [21, Theorem 4.8] to (26), which we further reformulate as

$$(27) \quad -\lambda \Delta \widetilde{\psi}_\infty + \min\{e^{-V_n}, n_0 p_0 e^{-V_p}\} \widetilde{\psi}_\infty + d(\cdot, \widetilde{\psi}_\infty) = -D$$

where

$$d(x, y) := -e^{-y - V_n} + n_0 p_0 e^{y - V_p} - \frac{\varepsilon}{1 + n_0 e^y} - \min\{e^{-V_n}, n_0 p_0 e^{-V_p}\} y.$$

The structure of (27) is suitable to apply [21, Theorem 4.8] for the existence of a unique continuous solution provided that d is monotone increasing w.r.t. y . Indeed, direct computations show

$$e^{-y - V_n} + n_0 p_0 e^{y - V_p} \geq 2\sqrt{n_0 p_0} e^{-\frac{V_p + V_n}{2}}, \quad \forall y \in \mathbb{R},$$

(where the lower bound is attained at the unique minimum $e^y = e^{\frac{V_p - V_n}{2}} / \sqrt{n_0 p_0}$). Hence, we estimate independently of ε

$$\begin{aligned} \partial_y d(x, y) &\geq e^{-y - V_n} + n_0 p_0 e^{y - V_p} - \min\{e^{-V_n}, n_0 p_0 e^{-V_p}\} \\ &\geq 2e^{-\frac{V_n}{2}} \sqrt{n_0 p_0} e^{-\frac{V_p}{2}} - \min\{e^{-V_n}, n_0 p_0 e^{-V_p}\} > 0, \end{aligned}$$

and therefore strict monotonicity of d w.r.t. y follows.

As a consequence, (27) admits a unique solution $\widetilde{\psi}_\infty \in H^1(\Omega) \cap L^\infty(\Omega)$, which is continuous on $\overline{\Omega}$ and bounded via

$$\|\widetilde{\psi}_\infty\|_{H^1(\Omega)} + \|\widetilde{\psi}_\infty\|_{C(\overline{\Omega})} \leq \widetilde{K}_\infty$$

where the constant \widetilde{K}_∞ is independent of ε . Going back to ψ_∞ , the constraint $\overline{\psi_\infty} = 0$ implies

$$n_* = e^{-\int_\Omega \widetilde{\psi}_\infty dx},$$

which in turn uniquely determines $\psi_\infty = \widetilde{\psi}_\infty - \int_\Omega \widetilde{\psi}_\infty dx$. The bounds on $\widetilde{\psi}_\infty$ directly transfer to ψ_∞ .

As in [8], we verify the bounds (12) by solving the two equations in (14) for $n_* > 0$ abbreviating $V_\infty := \max\{\|V_n\|_{L^\infty(\Omega)}, \|V_p\|_{L^\infty(\Omega)}\}$:

$$\begin{aligned} n_* &= \frac{D - \varepsilon n_{tr,\infty}}{2e^{-\psi_\infty - V_n}} + \sqrt{\frac{D - \varepsilon n_{tr,\infty}}{4e^{-\psi_\infty - V_n}}^2 + n_0 p_0 \frac{e^{\psi_\infty - V_p}}{e^{-\psi_\infty - V_n}}} \\ &\leq e^{K_\infty + V_\infty} (\sqrt{n_0 p_0} + \varepsilon_0 + |\overline{D}|). \end{aligned}$$

We stress that the same bound is valid also for $p_* > 0$, and that the upper and lower bounds on n_∞ , p_∞ and $n_{tr,\infty}$ are a consequence of the bounds on n_* and p_* as well as $n_* p_* = n_0 p_0$. Finally, the estimate

$$\|\psi_\infty\|_{H^2(\Omega)} \leq C \|n_\infty - p_\infty + \varepsilon n_{tr,\infty} - D\|_{L^2(\Omega)} \leq C(K_\infty)$$

ensures the higher regularity of ψ_∞ . \square

3. DERIVATION OF AN EEP INEQUALITY

As an auxiliary result, we first derive a convenient expression for the entropy relative to the equilibrium.

Lemma 3.1. *The entropy relative to the equilibrium equals*

$$\begin{aligned} & E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr,\infty}, \psi_\infty) \\ &= \int_\Omega \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) + p \ln \frac{p}{p_\infty} - (p - p_\infty) \right. \\ & \quad \left. + \frac{\lambda}{2} |\nabla(\psi - \psi_\infty)|^2 + \varepsilon \int_{n_{tr,\infty}}^{n_{tr}} \left(\ln \frac{s}{1-s} - \ln \frac{n_{tr,\infty}}{1-n_{tr,\infty}} \right) ds \right) dx. \end{aligned}$$

Proof. According to the definition of $E(n, p, n_{tr}, \psi)$, one has

$$\begin{aligned} & E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr,\infty}, \psi_\infty) \\ &= \int_\Omega \left(n \ln \frac{n}{n_0 \mu_n} - n_\infty \ln \frac{n_\infty}{n_0 \mu_n} - (n - n_\infty) + p \ln \frac{p}{p_0 \mu_p} - p_\infty \ln \frac{p_\infty}{p_0 \mu_p} - (p - p_\infty) \right. \\ & \quad \left. + \frac{\lambda}{2} (|\nabla \psi|^2 - |\nabla \psi_\infty|^2) + \varepsilon \int_{n_{tr,\infty}}^{n_{tr}} \ln \frac{s}{1-s} ds \right) dx. \end{aligned}$$

We rewrite the first integrand as $n \ln \frac{n}{n_0 \mu_n} = n \ln \frac{n}{n_\infty} + n \ln \frac{n_\infty}{n_0 \mu_n}$ and use $\frac{n_\infty}{n_0 \mu_n} = \frac{n_*}{n_0} e^{-\psi_\infty}$ to find

$$\begin{aligned} & \int_\Omega \left(n \ln \frac{n}{n_0 \mu_n} - n_\infty \ln \frac{n_\infty}{n_0 \mu_n} - (n - n_\infty) \right) dx \\ &= \int_\Omega \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) + (n - n_\infty) \left(\ln \frac{n_*}{n_0} - \psi_\infty \right) \right) dx. \end{aligned}$$

Together with an analogous calculation for the p -terms, we obtain

$$\begin{aligned} & E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr,\infty}, \psi_\infty) \\ &= \int_\Omega \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) + p \ln \frac{p}{p_\infty} - (p - p_\infty) \right. \\ & \quad \left. + (n - n_\infty) \left(\ln \frac{n_*}{n_0} - \psi_\infty \right) + (p - p_\infty) \left(\ln \frac{p_*}{p_0} + \psi_\infty \right) \right. \\ & \quad \left. + \frac{\lambda}{2} |\nabla \psi|^2 - \frac{\lambda}{2} |\nabla \psi_\infty|^2 + \varepsilon \int_{n_{tr,\infty}}^{n_{tr}} \ln \frac{s}{1-s} ds \right) dx. \end{aligned}$$

We now employ the conservation law $\bar{p} - \bar{p}_\infty = \bar{n} - \bar{n}_\infty + \varepsilon(\bar{n}_{tr} - \bar{n}_{tr,\infty})$, the formula $n_* p_* = n_0 p_0$ and the representation $\frac{p_*}{p_0} = \frac{1-n_{tr,\infty}}{n_{tr,\infty}} e^{-\psi_\infty}$ to derive

$$\begin{aligned} & (\bar{n} - \bar{n}_\infty) \ln \frac{n_*}{n_0} + (\bar{p} - \bar{p}_\infty) \ln \frac{p_*}{p_0} \\ &= (\bar{n} - \bar{n}_\infty) \ln \frac{n_* p_*}{n_0 p_0} + \varepsilon \int_\Omega (n_{tr} - n_{tr,\infty}) \ln \frac{p_*}{p_0} dx \\ &= \varepsilon \int_\Omega (n_{tr} - n_{tr,\infty}) \left(\ln \frac{1-n_{tr,\infty}}{n_{tr,\infty}} - \psi_\infty \right) dx \\ &= -\varepsilon \int_\Omega \left((n_{tr} - n_{tr,\infty}) \psi_\infty + \int_{n_{tr,\infty}}^{n_{tr}} \ln \frac{n_{tr,\infty}}{1-n_{tr,\infty}} ds \right) dx. \end{aligned}$$

The relative entropy now reads

$$\begin{aligned}
& E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \\
&= \int_{\Omega} \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) + p \ln \frac{p}{p_\infty} - (p - p_\infty) \right. \\
&\quad \left. + \frac{\lambda}{2} |\nabla \psi|^2 - \frac{\lambda}{2} |\nabla \psi_\infty|^2 - (n - n_\infty - p + p_\infty + \varepsilon(n_{tr} - n_{tr, \infty})) \psi_\infty \right. \\
&\quad \left. + \varepsilon \int_{n_{tr, \infty}}^{n_{tr}} \left(\ln \frac{s}{1-s} - \ln \frac{n_{tr, \infty}}{1-n_{tr, \infty}} \right) ds \right) dx.
\end{aligned}$$

Poisson's equation $n - n_\infty - p + p_\infty + \varepsilon(n_{tr} - n_{tr, \infty}) = -\lambda \Delta(\psi - \psi_\infty)$ and an integration by parts entail

$$\begin{aligned}
& E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \\
&= \int_{\Omega} \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) + p \ln \frac{p}{p_\infty} - (p - p_\infty) + \frac{\lambda}{2} |\nabla \psi|^2 - \frac{\lambda}{2} |\nabla \psi_\infty|^2 \right. \\
&\quad \left. - \lambda \nabla(\psi - \psi_\infty) \cdot \nabla \psi_\infty + \varepsilon \int_{n_{tr, \infty}}^{n_{tr}} \left(\ln \frac{s}{1-s} - \ln \frac{n_{tr, \infty}}{1-n_{tr, \infty}} \right) ds \right) dx.
\end{aligned}$$

The claim now obviously follows from collecting the terms involving ψ and ψ_∞ . \square

Following ideas in [8, 12] and [9], we are able to bound the relative entropy essentially in terms of the squared L^2 distance between $(n, p, \sqrt{n_{tr}})$ and $(n_\infty, p_\infty, \sqrt{n_{tr, \infty}})$.

Proposition 3.2. *There exists an explicit constant $c_1(K_\infty) > 0$ satisfying*

$$\begin{aligned}
& E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \\
&\leq c_1 \int_{\Omega} \left(\frac{(n - n_\infty)^2}{n_\infty} + \frac{(p - p_\infty)^2}{p_\infty} + \varepsilon(\sqrt{n_{tr}} - \sqrt{n_{tr, \infty}})^2 \right) dx
\end{aligned}$$

for all $\varepsilon > 0$ and all non-negative $n, p, n_{tr} \in L^2(\Omega)$, $n_{tr} \leq 1$ where $\psi \in H^1(\Omega)$ is the unique solution of (5) with $f = n - p + \varepsilon n_{tr} - D$ and $\bar{\psi} = 0$.

Proof. Applying the elementary inequality $\ln x \leq x - 1$ for $x > 0$, we derive

$$n \ln \frac{n}{n_\infty} - (n - n_\infty) \leq n \left(\frac{n}{n_\infty} - 1 \right) - n + n_\infty = \frac{(n - n_\infty)^2}{n_\infty}$$

and an analogous estimate involving p and p_∞ . Integration by parts with homogeneous Neumann conditions for ψ and ψ_∞ as well as $-\lambda \Delta(\psi - \psi_\infty) = (n - n_\infty) - (p - p_\infty) + \varepsilon(n_{tr} - n_{tr, \infty})$ yield

$$\begin{aligned}
& \lambda \int_{\Omega} |\nabla(\psi - \psi_\infty)|^2 dx \\
&= \int_{\Omega} ((n - n_\infty) - (p - p_\infty) + \varepsilon(n_{tr} - n_{tr, \infty}))(\psi - \psi_\infty) dx \\
&\leq \frac{1}{2} \left(\frac{1}{\delta} \|(n - n_\infty) - (p - p_\infty) + \varepsilon(n_{tr} - n_{tr, \infty})\|^2 + \delta \|\psi - \psi_\infty\|^2 \right) \\
&\leq \frac{3L(\Omega)}{2\lambda} \left(\|n - n_\infty\|^2 + \|p - p_\infty\|^2 + \varepsilon \|n_{tr} - n_{tr, \infty}\|^2 \right) + \frac{\lambda}{2} \|\nabla(\psi - \psi_\infty)\|^2.
\end{aligned}$$

Here and below, we abbreviate $\|\cdot\| := \|\cdot\|_{L^2(\Omega)}$ and denote by $L(\Omega) > 0$ a constant such that Poincaré's estimate $\|f\|^2 \leq L(\Omega) \|\nabla f\|^2$ holds true for all $f \in H^1(\Omega)$ subject to $\bar{f} = 0$. The estimate in the last line is then a result of $\bar{\psi} - \bar{\psi}_\infty = 0$ and Poincaré's inequality together with the choice $\delta := \lambda/L(\Omega)$, whereas the previous

bound follows from Hölder's inequality and Young's inequality with some constant $\delta > 0$. We thus find

$$\begin{aligned} & \frac{\lambda}{2} \int_{\Omega} |\nabla(\psi - \psi_{\infty})|^2 dx \\ & \leq \frac{6L(\Omega)M_{\infty}}{\lambda} \int_{\Omega} \left(\frac{(n - n_{\infty})^2}{n_{\infty}} + \frac{(p - p_{\infty})^2}{p_{\infty}} + \varepsilon(\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}})^2 \right) dx, \end{aligned}$$

where we employed the bounds from (12) and $(\sqrt{n_{tr}} + \sqrt{n_{tr,\infty}})^2 \leq 4$.

The last term within the relative entropy including n_{tr} can be controlled as in [9]. For convenience, we briefly recall the main arguments. First, there exists for all $x \in \Omega$ some mean value

$$\theta(x) \in (\min\{n_{tr}(x), n_{tr,\infty}(x)\}, \max\{n_{tr}(x), n_{tr,\infty}(x)\})$$

such that

$$(28) \quad \int_{n_{tr,\infty}(x)}^{n_{tr}(x)} \ln \frac{s}{1-s} ds = (n_{tr}(x) - n_{tr,\infty}(x)) \ln \frac{\theta(x)}{1-\theta(x)}.$$

To enhance readability, we shall suppress the x -dependence of n_{tr} and $n_{tr,\infty}$ subsequently. We further use the bound $n_{tr,\infty} \in (\mu_{\infty}, 1 - \mu_{\infty})$ from (12) and observe that

$$\left| \int_{n_{tr,\infty}}^{n_{tr}} \ln \frac{s}{1-s} ds \right| \leq \int_0^1 \left| \ln \frac{s}{1-s} \right| ds = 2 \ln 2$$

for all $x \in \Omega$. In combination with (28), this estimate entails

$$\left| \ln \frac{\theta(x)}{1-\theta(x)} \right| |n_{tr} - n_{tr,\infty}| \leq 2 \ln 2.$$

By an elementary argumentation, one can now conclude that $\theta(x) \in (\xi, 1-\xi)$ where $\xi \in (0, \frac{1}{2})$ only depends on μ_{∞} . Therefore, we obtain

$$\begin{aligned} & \varepsilon \int_{\Omega} \int_{n_{tr,\infty}}^{n_{tr}} \left(\ln \frac{s}{1-s} - \ln \frac{n_{tr,\infty}}{1-n_{tr,\infty}} \right) ds dx \\ & = \varepsilon \int_{\Omega} \left(\ln \frac{\theta(x)}{1-\theta(x)} - \ln \frac{n_{tr,\infty}}{1-n_{tr,\infty}} \right) (n_{tr} - n_{tr,\infty}) dx \\ & = \varepsilon \int_{\Omega} \frac{1}{\sigma(x)(1-\sigma(x))} (\theta(x) - n_{tr,\infty})(n_{tr} - n_{tr,\infty}) dx \end{aligned}$$

with some $\sigma(x) \in (\min\{\theta(x), n_{tr,\infty}(x)\}, \max\{\theta(x), n_{tr,\infty}(x)\}) \subset [\xi, 1-\xi]$ employing the mean-value theorem and taking into account that

$$\frac{d}{ds} \ln \frac{s}{1-s} = \frac{1}{s(1-s)}.$$

As $(\sigma(x)(1-\sigma(x)))^{-1}$ is uniformly bounded in Ω in terms of $\xi(\mu_{\infty})$, there exists some $c > 0$ only depending on μ_{∞} such that

$$\begin{aligned} & \varepsilon \int_{\Omega} \int_{n_{tr,\infty}}^{n_{tr}} \left(\ln \frac{s}{1-s} - \ln \frac{n_{tr,\infty}}{1-n_{tr,\infty}} \right) ds dx \\ & \leq c\varepsilon \int_{\Omega} |\theta(x) - n_{tr,\infty}| |n_{tr} - n_{tr,\infty}| dx \\ & \leq 4c\varepsilon \int_{\Omega} (\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}})^2 dx \end{aligned}$$

where the last line results from estimating $(\sqrt{n_{tr}} + \sqrt{n_{tr,\infty}})^2 \leq 4$. This proves the claim. \square

The subsequent lemma contains rather non-intuitive estimates for bilinear terms like $(n - n_\infty)(p - p_\infty)$. These expressions will appear in the proof of Proposition 3.4 below.

Lemma 3.3. *The following estimates hold true for all non-negative $n, p, n_{tr} \in L^\infty(\Omega)$ satisfying $n, p \leq M$ and $n_{tr} \leq 1$ with explicit constants $\Gamma_1(M) > 0$ and $\Gamma_2 > 0$:*

$$\begin{aligned} (n - n_\infty)(p - p_\infty) &\leq \Gamma_1 \left(-R_n \ln \frac{n(1 - n_{tr})}{n_0 \mu_n n_{tr}} - R_p \ln \frac{pn_{tr}}{p_0 \mu_p (1 - n_{tr})} \right), \\ (n - n_\infty)(-n_{tr} + n_{tr,\infty}) &\leq \Gamma_2 \left(-R_n \ln \frac{n(1 - n_{tr})}{n_0 \mu_n n_{tr}} + (\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}})^2 \right), \\ (p - p_\infty)(n_{tr} - n_{tr,\infty}) &\leq \Gamma_2 \left(-R_p \ln \frac{pn_{tr}}{p_0 \mu_p (1 - n_{tr})} + (\sqrt{1 - n_{tr}} - \sqrt{1 - n_{tr,\infty}})^2 \right). \end{aligned}$$

Proof. As in [12] and [8], we first recall the elementary inequalities $(a - a_0)(b - b_0) \leq (\sqrt{ab} - \sqrt{a_0 b_0})^2$ for all $a, a_0, b, b_0 \geq 0$ and $4(\sqrt{x} - \sqrt{y})^2 \leq (x - y) \ln \frac{x}{y}$ for all $x \geq 0$ and $y > 0$.

Concerning the first inequality we write

$$(n - n_\infty)(p - p_\infty) \leq (\sqrt{np} - \sqrt{n_\infty p_\infty})^2 = n_\infty p_\infty \left(\sqrt{\frac{np}{n_0 \mu_n p_0 \mu_p}} - 1 \right)^2$$

and distinguish the two cases $n_{tr} > \frac{1}{2}$ and $n_{tr} \leq \frac{1}{2}$. In the case $n_{tr} > \frac{1}{2}$, we infer

$$\begin{aligned} (n - n_\infty)(p - p_\infty) &\leq n_0 p_0 \mu_n \mu_p \left(\sqrt{\frac{n}{n_0 \mu_n n_{tr}}} \left(\sqrt{\frac{p}{p_0 \mu_p} n_{tr}} - \sqrt{1 - n_{tr}} \right) \right. \\ &\quad \left. + \sqrt{\frac{1}{n_{tr}}} \left(\sqrt{\frac{n}{n_0 \mu_n} (1 - n_{tr})} - \sqrt{n_{tr}} \right) \right)^2 \\ &\leq \Gamma_1(M) \left(\left(\frac{n}{n_0 \mu_n} (1 - n_{tr}) - n_{tr} \right) \ln \frac{n(1 - n_{tr})}{n_0 \mu_n n_{tr}} \right. \\ &\quad \left. + \left(\frac{p}{p_0 \mu_p} n_{tr} - (1 - n_{tr}) \right) \ln \frac{pn_{tr}}{p_0 \mu_p (1 - n_{tr})} \right) \end{aligned}$$

employing the assumed L^∞ bound on n . Using analogous arguments we derive the same result also in the case $n_{tr} \leq \frac{1}{2}$. The second inequality arises from

$$\begin{aligned} (n - n_\infty)(-n_{tr} + n_{tr,\infty}) &\leq \left(\sqrt{n(1 - n_{tr})} - \sqrt{n_\infty(1 - n_{tr,\infty})} \right)^2 \\ &= n_0 \mu_n \left(\sqrt{\frac{n}{n_0 \mu_n} (1 - n_{tr})} - \sqrt{n_{tr,\infty}} \right)^2 \end{aligned}$$

where we used the relation $n_\infty(1 - n_{tr,\infty}) = n_0 \mu_n n_{tr,\infty}$ from Proposition 1.4. The claim is now a consequence of

$$\begin{aligned} n_0 \mu_n \left(\left(\sqrt{\frac{n}{n_0 \mu_n} (1 - n_{tr})} - \sqrt{n_{tr}} \right) + (\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}}) \right)^2 \\ \leq \Gamma_2 \left(\left(\frac{n(1 - n_{tr})}{n_0 \mu_n} - n_{tr} \right) \ln \frac{n(1 - n_{tr})}{n_0 \mu_n n_{tr}} + (\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}})^2 \right). \end{aligned}$$

Similarly, one can also verify the third inequality stated above. \square

The next result establishes an upper bound for the L^2 distance between (n, p) and (n_∞, p_∞) basically in terms of the entropy production P and $\|\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}}\|_{L^2(\Omega)}^2$. Similar arguments already appeared in [12] and [8].

Proposition 3.4. *There exists an explicit constant $c_2(M, K_\infty) > 0$ satisfying*

$$\begin{aligned} \int_{\Omega} \left(\frac{(n - n_\infty)^2}{n_\infty} + \frac{(p - p_\infty)^2}{p_\infty} \right) dx &\leq c_2 P(n, p, n_{tr}, \psi) \\ &+ c_2 \varepsilon \int_{\Omega} \left((\sqrt{n_{tr}} - \sqrt{n_{tr, \infty}})^2 + (\sqrt{1 - n_{tr}} - \sqrt{1 - n_{tr, \infty}})^2 \right) dx \end{aligned}$$

for all $\varepsilon > 0$ and all non-negative $n, p, n_{tr} \in L^\infty(\Omega)$ subject to $n, p \leq M$ and $n_{tr} \leq 1$ where $\psi \in H^1(\Omega)$ is the unique solution of (5) with $f = n - p + \varepsilon n_{tr} - D$ and $\bar{\psi} = 0$.

Proof. We start by defining intermediate equilibria $N := n_* e^{-\psi - V_n}$ and $P := p_* e^{\psi - V_p}$ which fulfill $J_n(N, \psi) = J_p(P, \psi) = 0$. Due to $J_n(n, \psi) = N \nabla(\frac{n}{N})$ and $J_p(p, \psi) = P \nabla(\frac{p}{P})$, we derive the following lower bounds for the flux terms involving J_n and J_p :

$$\begin{aligned} \frac{|J_n|^2}{n_\infty n} &= \frac{N^2}{n_\infty n} \left| \nabla \left(\frac{n}{N} \right) \right|^2 = \frac{N^2}{n_\infty n} \left| \frac{n_\infty}{N} \nabla \left(\frac{n}{n_\infty} \right) + \frac{n}{n_\infty} \nabla \left(\frac{n_\infty}{N} \right) \right|^2 \\ &= \frac{N^2}{n_\infty n} \left| e^{\psi - \psi_\infty} \nabla \left(\frac{n}{n_\infty} \right) + \frac{n}{n_\infty} e^{\psi - \psi_\infty} \nabla (\psi - \psi_\infty) \right|^2 \\ &\geq 2 \frac{N^2}{n_\infty^2} e^{2(\psi - \psi_\infty)} \nabla \left(\frac{n}{n_\infty} \right) \cdot \nabla (\psi - \psi_\infty) = 2 \nabla (\psi - \psi_\infty) \cdot \nabla \left(\frac{n - n_\infty}{n_\infty} \right). \end{aligned}$$

In the same way, we obtain $\frac{|J_p|^2}{p_\infty p} \geq -2 \nabla (\psi - \psi_\infty) \cdot \nabla \left(\frac{p - p_\infty}{p_\infty} \right)$ and, therefore,

$$\begin{aligned} \frac{\lambda}{2} \int_{\Omega} \left(\frac{|J_n|^2}{n_\infty n} + \frac{|J_p|^2}{p_\infty p} \right) dx &\geq \lambda \int_{\Omega} \nabla (\psi - \psi_\infty) \cdot \nabla \left(\frac{n - n_\infty}{n_\infty} - \frac{p - p_\infty}{p_\infty} \right) dx \\ &= \int_{\Omega} \left((n - n_\infty) - (p - p_\infty) + \varepsilon (n_{tr} - n_{tr, \infty}) \right) \left(\frac{n - n_\infty}{n_\infty} - \frac{p - p_\infty}{p_\infty} \right) dx \end{aligned}$$

via integration by parts and Poisson's equation. Rearranging this inequality now yields

$$\begin{aligned} &\int_{\Omega} \left(\frac{(n - n_\infty)^2}{n_\infty} + \frac{(p - p_\infty)^2}{p_\infty} \right) dx \\ &\leq \frac{\lambda}{2} \int_{\Omega} \left(\frac{|J_n|^2}{n_\infty n} + \frac{|J_p|^2}{p_\infty p} \right) dx + \int_{\Omega} \left(\left(\frac{1}{n_\infty} + \frac{1}{p_\infty} \right) (n - n_\infty)(p - p_\infty) \right. \\ &\quad \left. + \varepsilon (-n_{tr} + n_{tr, \infty}) \frac{n - n_\infty}{n_\infty} + \varepsilon (n_{tr} - n_{tr, \infty}) \frac{p - p_\infty}{p_\infty} \right) dx. \end{aligned}$$

Together with the bounds from (12) and Lemma 3.3, we arrive at the desired result. \square

We are now in a position to prove the EEP inequality from Theorem 1.5, where the main task is to provide an appropriate bound on $(\sqrt{n_{tr}} - \sqrt{n_{tr, \infty}})^2$.

Proof of Theorem 1.5. Step 1. Due to (15) we easily calculate

$$\begin{aligned} \sqrt{n_{tr}} - \sqrt{\frac{n}{n_0 \mu_n} (1 - n_{tr})} &= \sqrt{n_{tr}} - \sqrt{n_{tr, \infty}} - \sqrt{\frac{n}{n_0 \mu_n} (1 - n_{tr})} \\ &\quad \pm \sqrt{\frac{n_\infty}{n_0 \mu_n} (1 - n_{tr})} + \sqrt{\frac{n_\infty}{n_0 \mu_n} (1 - n_{tr, \infty})} \\ &= \sqrt{n_{tr}} - \sqrt{n_{tr, \infty}} - \sqrt{\frac{1 - n_{tr}}{n_0 \mu_n}} (\sqrt{n} - \sqrt{n_\infty}) \\ &\quad + \sqrt{\frac{n_\infty}{n_0 \mu_n}} (\sqrt{1 - n_{tr, \infty}} - \sqrt{1 - n_{tr}}). \end{aligned}$$

Observing that $\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}}$ and $\sqrt{1 - n_{tr,\infty}} - \sqrt{1 - n_{tr}}$ have the same sign and using the inequality $4(\sqrt{x} - \sqrt{y})^2 \leq (x - y) \ln \frac{x}{y}$ for $x \geq 0$ and $y > 0$, we reformulate the previous identity to find

$$\begin{aligned} (\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}})^2 &\leq \left(\sqrt{n_{tr}} - \sqrt{n_{tr,\infty}} + \sqrt{\frac{n_\infty}{n_0\mu_n}} (\sqrt{1 - n_{tr,\infty}} - \sqrt{1 - n_{tr}}) \right)^2 \\ &\leq 2 \left(\sqrt{n_{tr}} - \sqrt{\frac{n}{n_0\mu_n}(1 - n_{tr})} \right)^2 + \frac{2}{n_0\mu_n} (\sqrt{n} - \sqrt{n_\infty})^2 \\ &\leq \frac{1}{2} \left(\frac{n(1 - n_{tr})}{n_0\mu_n} - n_{tr} \right) \ln \frac{n(1 - n_{tr})}{n_0\mu_n n_{tr}} + \frac{2}{n_0\mu_n} \frac{(n - n_\infty)^2}{n_\infty}. \end{aligned}$$

Along the same lines, we also deduce

$$\begin{aligned} (\sqrt{1 - n_{tr}} - \sqrt{1 - n_{tr,\infty}})^2 &\leq \frac{1}{2} \left(\frac{pn_{tr}}{p_0\mu_p} - (1 - n_{tr}) \right) \ln \frac{pn_{tr}}{p_0\mu_p(1 - n_{tr})} + \frac{2}{p_0\mu_p} \frac{(p - p_\infty)^2}{p_\infty}. \end{aligned}$$

Step 2. We can now improve the claim of Proposition 3.2 in the sense that there exists a constant $c_1(K_\infty) > 0$ satisfying

$$\begin{aligned} E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr,\infty}, \psi_\infty) \\ \leq c_1 P(n, p, n_{tr}, \psi) + c_1 \int_\Omega \left(\frac{(n - n_\infty)^2}{n_\infty} + \frac{(p - p_\infty)^2}{p_\infty} \right) dx \end{aligned}$$

where $n, p, n_{tr} \in L^2(\Omega)$, $n_{tr} \leq 1$ are arbitrary non-negative functions and $\psi \in H^1(\Omega)$ is the unique solution of (5) with $f = n - p + \varepsilon n_{tr} - D$ and $\bar{\psi} = 0$ for any $\varepsilon > 0$. Furthermore, we notice that Proposition 3.4 now gives rise to a constant $c_2(M, K_\infty) > 0$ such that

$$\begin{aligned} \int_\Omega \left(\frac{(n - n_\infty)^2}{n_\infty} + \frac{(p - p_\infty)^2}{p_\infty} \right) dx \\ \leq c_2 P(n, p, n_{tr}, \psi) + c_2 \varepsilon \int_\Omega \left(\frac{(n - n_\infty)^2}{n_\infty} + \frac{(p - p_\infty)^2}{p_\infty} \right) dx \end{aligned}$$

holds true for all $\varepsilon > 0$, all non-negative $n, p, n_{tr} \in L^\infty(\Omega)$, $n, p \leq M$ and $n_{tr} \leq 1$, and $\psi \in H^1(\Omega)$ being the unique solution of (5) with $f = n - p + \varepsilon n_{tr} - D$ and $\bar{\psi} = 0$.

Step 3. If we restrict ε to the interval $(0, \frac{1}{2c_2})$, we finally arrive at

$$E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr,\infty}, \psi_\infty) \leq (c_1 + 2c_1c_2)P(n, p, n_{tr}, \psi)$$

employing the notation from Step 2. \square

4. PROOF OF THE EXPONENTIAL CONVERGENCE

As soon as the weak entropy production law (19) is settled, the exponential decay of the relative entropy arises from a Gronwall argument as carried out in [9] (see also [1, 22]).

Proof of Theorem 1.6. The weak entropy production law (19) readily follows from (16) and (18) for $0 < t_0 \leq t_1 < \infty$ utilizing the regularity and bounds on n, p, n_{tr} , and ψ from Proposition 1.3. The statement of the theorem is then a consequence

of Theorem 1.5 applied to the global solution (n, p, n_{tr}, ψ) . Note, however, that the weak entropy production law (19) only allows to derive

$$\begin{aligned} E(n, p, n_{tr}, \psi)(t) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \\ \leq (E(n, p, n_{tr}, \psi)(t_0) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty)) e^{-C_{EEP}^{-1}(t-t_0)} \end{aligned}$$

for all $t_0 \in (0, t]$. The assertion in (20) is then a consequence of the fact that the entropy $E(n, p, n_{tr}, \psi)(t_0)$ continuously extends to $t_0 \rightarrow 0$ since $n, p \in C([0, T], L^2(\Omega))$, $n_{tr} \in C([0, T], L^\infty(\Omega))$, and $\psi \in C([0, T], H^2(\Omega))$ for all $T > 0$ by Proposition 1.3. \square

Exponential convergence in L^∞ and H^2 for (n, p, n_{tr}) and ψ , respectively, is a consequence of standard regularity techniques, which have been partially employed already in [9].

Proof of Corollary 1.7. An immediate consequence of the exponential decay of the relative entropy as stated in (20) is the exponential convergence to the equilibrium of $n(t)$ and $p(t)$ in $L^1(\Omega)$, and of $n_{tr}(t)$ in $L^2(\Omega)$. To see this, we recall from Lemma 3.1 that the relative entropy is given by

$$\begin{aligned} E(n, p, n_{tr}, \psi) - E(n_\infty, p_\infty, n_{tr, \infty}, \psi_\infty) \\ = \int_\Omega \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) + p \ln \frac{p}{p_\infty} - (p - p_\infty) \right. \\ \left. + \frac{\lambda}{2} |\nabla(\psi - \psi_\infty)|^2 + \varepsilon \int_{n_{tr, \infty}}^{n_{tr}} \left(\ln \frac{s}{1-s} - \ln \frac{n_{tr, \infty}}{1-n_{tr, \infty}} \right) ds \right) dx. \end{aligned}$$

The well-known Csiszár–Kullback–Pinsker inequality allows us to control

$$\int_\Omega \left(n \ln \frac{n}{n_\infty} - (n - n_\infty) \right) dx \geq \frac{3}{2\bar{n} + 4n_\infty} \|n - n_\infty\|_{L^1(\Omega)}^2 \geq c \|n - n_\infty\|_{L^1(\Omega)}^2$$

and analogously $\|p - p_\infty\|_{L^1(\Omega)}^2$ in terms of a (rough) constant $c(M, K_\infty) > 0$. Next, we notice that

$$\frac{d}{ds} \ln \left(\frac{s}{1-s} \right) = \frac{1}{s(1-s)} \geq 4$$

is valid for all $s \in (0, 1)$, which enables us to estimate

$$\begin{aligned} \varepsilon \int_\Omega \int_{n_{tr, \infty}}^{n_{tr}} \left(\ln \left(\frac{s}{1-s} \right) - \ln \left(\frac{n_{tr, \infty}}{1-n_{tr, \infty}} \right) \right) ds dx \\ = \varepsilon \int_\Omega \int_{n_{tr, \infty}}^{n_{tr}} \frac{1}{\sigma(s)(1-\sigma(s))} (s - n_{tr, \infty}) ds dx \geq 2\varepsilon \|n_{tr} - n_{tr, \infty}\|_{L^2(\Omega)}^2 \end{aligned}$$

where $\sigma(s)$ serves as an intermediate point between $n_{tr, \infty}$ and s .

As a prerequisite for the exponential convergence of n and p in $L^\infty(\Omega)$, we adapt an argument from [9] to establish a polynomially growing $W^{1,q}(\Omega)$ bound on n for $q \geq 4$. The same technique is also applicable to p . As in the proof of Proposition 1.3, we introduce $w := e^{\psi+V_n} n$ (satisfying $\hat{n} \cdot \nabla w = 0$ on $\partial\Omega$) and set w.l.o.g. $\tau_n = \tau_p = 1$ leading to

$$\partial_t w = \Delta w - \nabla(\psi + V_n) \cdot \nabla w + e^{\psi+V_n} n_{tr} - \frac{1 - n_{tr}}{n_0 \mu_n} w.$$

Using $-(q-1)|\nabla w|^{q-2}\Delta w$ as a test function entails

$$\begin{aligned} \frac{1}{q} \frac{d}{dt} \int_{\Omega} |\nabla w|^q dx &= \int_{\Omega} |\nabla w|^{q-2} \nabla w \cdot \nabla \partial_t w dx = - \int_{\Omega} (q-1) |\nabla w|^{q-2} \Delta w \partial_t w dx \\ &= - \int_{\Omega} (q-1) |\nabla w|^{q-2} |\Delta w|^2 dx + \int_{\Omega} (q-1) |\nabla w|^{q-2} \nabla w \Delta w \cdot \nabla \psi dx \\ &\quad + \int_{\Omega} (q-1) |\nabla w|^{q-2} |\Delta w| C(M) (1 + w + |\nabla w|) dx. \end{aligned}$$

By applying Young's inequality to the last term on the right hand side, observing that $|\nabla w|^{q-2} \nabla w \Delta w = \frac{1}{q} \nabla (|\nabla w|^q)$, and integrating by parts the second term, we calculate

$$\begin{aligned} \frac{1}{q} \frac{d}{dt} \int_{\Omega} |\nabla w|^q dx &\leq -\frac{1}{2} \int_{\Omega} (q-1) |\nabla w|^{q-2} |\Delta w|^2 dx \\ &\quad + C(M) \int_{\Omega} (q-1) |\nabla w|^{q-2} (1 + w^2 + |\nabla w|^2) dx - \int_{\Omega} \frac{q-1}{q} |\nabla w|^q \Delta \psi dx. \end{aligned}$$

Taking into account $|\Delta \psi| \leq C(M)$, we deduce

$$\begin{aligned} \frac{1}{q(q-1)} \frac{d}{dt} \int_{\Omega} |\nabla w|^q dx &\leq -\frac{1}{2} \int_{\Omega} |\nabla w|^{q-2} |\Delta w|^2 dx \\ &\quad + C(M) \int_{\Omega} |\nabla w|^{q-2} (1 + w^2 + |\nabla w|^2) dx. \end{aligned}$$

We rewrite the very last term above by another integration by parts and Young's inequality with $C_1(M, q) > 0$, which leads us to

$$\begin{aligned} \int_{\Omega} |\nabla w|^{q-2} \nabla w \cdot \nabla w dx &= - \int_{\Omega} (q-1) |\nabla w|^{q-2} \Delta w w dx \\ &\leq \frac{1}{2C} \int_{\Omega} |\nabla w|^{q-2} |\Delta w|^2 dx + C_1 \int_{\Omega} |\nabla w|^{q-2} w^2 dx. \end{aligned}$$

The previous estimates and the bound $\|w\|_{L^\infty((0,T)\times\Omega)} \leq C(M)$ now guarantee that

$$\frac{d}{dt} \int_{\Omega} |\nabla w|^q dx \leq C_2 \int_{\Omega} |\nabla w|^{q-2} dx,$$

with a constant $C_2(M, q) > 0$. Choosing $t_0 > 0$ and $t \geq t_0$ arbitrarily and utilizing $|\Omega| = 1$, one has

$$\|\nabla w(t)\|_{L^q(\Omega)}^q \leq \|\nabla w(t_0)\|_{L^q(\Omega)}^q + C_2 \int_{t_0}^t \|\nabla w(s)\|_{L^q(\Omega)}^{q-2} ds.$$

The polynomial growth of $\|\nabla w\|_{L^q(\Omega)}$ is then obtained by an elementary Gronwall lemma (see e.g. [1]). In detail, there exists a constant $C_3(M, q) > 0$ such that

$$\|\nabla w(t)\|_{L^q(\Omega)} \leq \left(\|\nabla w(t_0)\|_{L^q(\Omega)}^2 + C_3(t - t_0) \right)^{\frac{1}{2}}.$$

The relation $\nabla n = e^{-\psi - V_n} \nabla w - n \nabla(\psi + V_n)$, in particular, ensures that

$$\begin{aligned} (29) \quad \|\nabla n\|_{L^6(\Omega)} &\leq C(M) (\|\nabla w\|_{L^6(\Omega)} + \|\nabla \psi\|_{L^6(\Omega)}) \\ &\leq C(M) (\|\nabla w\|_{L^6(\Omega)} + \|\psi\|_{H^2(\Omega)}) \leq C(M, t_0) (1 + t^{\frac{1}{2}}). \end{aligned}$$

Note that the second estimate arises from the continuous embedding $H^2(\Omega) \hookrightarrow W^{1,6}(\Omega)$ valid in \mathbb{R}^3 , whereas the last step results from the bound $\|\psi(t)\|_{H^2(\Omega)} \leq K(M)$.

Exponential convergence to the equilibrium for n in $L^q(\Omega)$, $1 < q < \infty$ is easily deduced from the exponential convergence of n in $L^1(\Omega)$ as settled above and the $L^\infty(\Omega)$ bounds on n and n_∞ in (8) and (12), respectively, by writing

$$\|n - n_\infty\|_{L^q(\Omega)}^q \leq \|n - n_\infty\|_{L^\infty(\Omega)}^{q-1} \|n - n_\infty\|_{L^1(\Omega)} \leq C e^{-ct}$$

where the constants $c(M, K_\infty, q), C(M, K_\infty, q) > 0$ are independent of ε for $\varepsilon \in (0, \varepsilon_0]$. For $q = 2$ and together with the same bound on p and the estimate

$$\|\psi - \psi_\infty\|_{H^2(\Omega)} \leq C(\|n - n_\infty\|_{L^2(\Omega)} + \|p - p_\infty\|_{L^2(\Omega)} + \varepsilon \|n_{tr} - n_{tr,\infty}\|_{L^2(\Omega)}),$$

this directly implies the exponential convergence of ψ in $H^2(\Omega)$. The Gagliardo–Nirenberg–Moser interpolation inequality now allows us to infer exponential convergence of n and p in $L^\infty(\Omega)$. In fact, the bound on $\|\nabla n\|_{L^6(\Omega)}$ in (29) entails

$$(30) \quad \|n - n_\infty\|_{L^\infty(\Omega)} \leq C \|n - n_\infty\|_{W^{1,6}(\Omega)}^{\frac{1}{2}} \|n - n_\infty\|_{L^6(\Omega)}^{\frac{1}{2}} \leq C e^{-ct},$$

with constants $c(M, K_\infty), C(M, K_\infty) > 0$.

The exponential convergence of n_{tr} in $L^\infty(\Omega)$ can be verified essentially along the same lines as in [9]. We, therefore, omit some technical details and set w.l.o.g. $\tau_n = \tau_p = 1$. By defining $u := n_{tr} - n_{tr,\infty}$, one derives the following pointwise relation by inserting $\pm n_{tr,\infty}$ several times and by applying the identities from (15):

$$\begin{aligned} \varepsilon \partial_t u &= R_p - R_n = \left(1 - n_{tr} - \frac{p}{p_0 \mu_p} n_{tr}\right) - \left(n_{tr} - \frac{n}{n_0 \mu_n} (1 - n_{tr})\right) \\ &= -u \left(2 + \frac{p_\infty}{p_0 \mu_p} + \frac{n_\infty}{n_0 \mu_n}\right) - \frac{n_{tr}}{p_0 \mu_p} (p - p_\infty) + \frac{(1 - n_{tr})}{n_0 \mu_n} (n - n_\infty). \end{aligned}$$

By recalling $0 \leq n_{tr} \leq 1$, $\mu_n = e^{-V_n}$, and $\mu_p = e^{-V_p}$ with $V_n, V_p \in L^\infty(\Omega)$, and by employing (30), we find

$$\frac{d}{dt} \|u(t, \cdot)\|_{L^\infty(\Omega)} \leq -\frac{2}{\varepsilon} \|u(t, \cdot)\|_{L^\infty(\Omega)} + \frac{C}{\varepsilon} e^{-ct}$$

where $c, C > 0$ depend on M and K_∞ but not on ε for $\varepsilon \in (0, \varepsilon_0]$. If we choose $c > 0$ sufficiently small satisfying $\varepsilon_0 c \leq 1$, we arrive at

$$\begin{aligned} \|n_{tr}(t, \cdot) - n_{tr,\infty}\|_{L^\infty(\Omega)} &\leq e^{-2t/\varepsilon} + \frac{C}{\varepsilon} \int_0^t e^{-2(t-s)/\varepsilon - cs} ds \\ &\leq e^{-2t/\varepsilon} + e^{-2t/\varepsilon} \frac{C}{2 - \varepsilon c} \left(e^{(2/\varepsilon - c)t} - 1\right) \leq e^{-2t/\varepsilon_0} + C e^{-ct}. \end{aligned}$$

Finally, estimate (21) is proven. \square

REFERENCES

- [1] P. R. Beesak, *Gronwall inequalities*, Carleton Math. Lecture Notes, vol. 11, 1975.
- [2] M. Chipot, *Elements of nonlinear analysis*, Birkhäuser advanced texts, Birkhäuser Verlag, Basel – Boston – Berlin, 2000.
- [3] L. Desvillettes and K. Fellner, *Exponential Decay toward Equilibrium via Entropy Methods for Reaction-Diffusion Equations*, J. Math. Anal. Appl. **319** (2006), 157–176.
- [4] L. Desvillettes and K. Fellner, *Entropy methods for reaction-diffusion equations: slowly growing a priori bounds*, Rev. Mat. Iberoam. **24** (2008), 407–431.
- [5] L. Desvillettes and K. Fellner, *Exponential Convergence to Equilibrium for a Nonlinear Reaction-Diffusion System Arising in Reversible Chemistry*, System Modelling and Optimization, IFIP AICT **443** (2014), 96–104.
- [6] Laurent Desvillettes, Klemens Fellner, and Bao Quoc Tang, *Trend to equilibrium for reaction-diffusion systems arising from complex balanced chemical reaction networks*, SIAM J. Math. Anal. **49** (2017), no. 4, 2666–2709.
- [7] M. Di Francesco, K. Fellner, and P. Markowich, *The entropy dissipation method for spatially inhomogeneous reaction-diffusion-type systems*, Proc. Roy. Soc. A **464** (2008), 3273–3300.

- [8] K. Fellner and M. Kniely, *On the entropy method and exponential convergence to equilibrium for a recombination-drift-diffusion system with self-consistent potential*, Applied Mathematics Letters **79** (2018), 196–204.
- [9] K. Fellner and M. Kniely, *Uniform convergence to equilibrium for a family of drift-diffusion models with trap-assisted recombination and the limiting Shockley–Read–Hall model*, Journal of Elliptic and Parabolic Equations (2020). Appeared online first, doi:10.1007/s41808-020-00068-8.
- [10] Klemens Fellner and Bao Quoc Tang, *Explicit exponential convergence to equilibrium for nonlinear reaction-diffusion systems with detailed balance condition*, Nonlinear Anal. **159** (2017), 145–180.
- [11] Klemens Fellner and Bao Quoc Tang, *Convergence to equilibrium of renormalised solutions to nonlinear chemical reaction-diffusion systems*, Z. Angew. Math. Phys. **69** (2018), no. 3, Paper No. 54, 30.
- [12] H. Gajewski and K. Gärtner, *On the Discretization of van Roosbroeck’s Equations with Magnetic Field*, ZAMM Z. angew. Math. Mech. **76** (1996), 247–264.
- [13] A. Glitzky, K. Gröger, and R. Hünlich, *Free energy and dissipation rate for reaction–diffusion processes of electrically charged species*, Appl. Anal. **60** (1996), 201–217.
- [14] A. Glitzky and R. Hünlich, *Energetic estimates and asymptotics for electro–reaction–diffusion systems*, ZAMM Z. angew. Math. Mech. **77** (1997), 823–832.
- [15] T. Goudon, V. Miljanović, and C. Schmeiser, *On the Shockley–Read–Hall Model: Generation–Recombination in Semiconductors*, SIAM J. Appl. Math. **67** (2007), 1183–1201.
- [16] R. N. Hall, *Electron-hole recombination in Germanium*, Phys. Rev. **87** (1952), 387.
- [17] P. A. Markowich, C. A. Ringhofer, and C. Schmeiser, *Semiconductor equations*, Springer-Verlag, Wien – New York, 1990.
- [18] A. Mielke, J. Haskovec, and P.A. Markowich, *On uniform decay of the entropy for reaction-diffusion systems*, J. Dynam. Differential Equations **27** (2015), 897–928.
- [19] V. P. Mikhaïlov, *Partial differential equations*, “Mir”, Moscow; distributed by Imported Publications, Inc., Chicago, Ill., 1978. Translated from the Russian by P. C. Sinha.
- [20] W. Shockley and W. T. Read, *Statistics of the recombinations of holes and electrons*, Phys. Rev. **87** (1952), 835–842.
- [21] F. Tröltzsch, *Optimal Control of Partial Differential Equations: Theory, Methods and Applications*, American Mathematical Society, 2010. Translated from German by Jürgen Sprekels.
- [22] D. Willett, *A linear generalization of Gronwall’s inequality*, Proc. Amer. Math. Soc. **16** (1965), 774–778.

INSTITUTE FOR MATHEMATICS AND SCIENTIFIC COMPUTING, UNIVERSITY OF GRAZ, HEINRICH-STRASSE 36, 8010 GRAZ, AUSTRIA

E-mail address: `klemens.fellner@uni-graz.at`

FACULTY OF MATHEMATICS, TU DORTMUND UNIVERSITY, VOGELPOTHSWEG 87, 44227 DORTMUND, GERMANY

E-mail address, Corresponding author: `michael.kniely@tu-dortmund.de`