

Hilbert Space Methods for Partial Differential Equations

a.o.Univ.Prof. Mag.Dr. Stephen Keeling

<http://math.uni-graz.at/keeling/>

Literature:

Hilbert Space Methods for Partial Differential Equations

by R.E. Showalter

Documentation:

<http://math.uni-graz.at/keeling/teaching.html>

Table of Contents I

Elements of Hilbert Space

- Linear Algebra
- Convergence and Continuity
- Completeness
- Hilbert Space
- Dual Operators and Identifications
- Uniform Boundedness and Weak Compactness
- Expansion in Eigenfunctions

Distributions and Sobolev Spaces

- Distributions
- Sobolev Spaces
- Trace
- Sobolev's Lemma and Imbedding
- Density and Compactness

Boundary Value Problems

- Introduction
- Forms, Operators and Green's Formula
- Abstract Boundary Value Problems
- Examples
- Coercivity and Elliptic Forms
- Regularity
- Closed Operators, Adjoint, Eigenfunction Expansions
- Introduction to Evolution Equations

Linear Algebra

- ▶ For F , *domain* is $\text{dom}(F)$ and *range* is $\text{Rg}(F)$.
- ▶ \mathbb{K} is field of scalars \mathbb{R} or \mathbb{C} , where \mathbb{C} are complex.
- ▶ G open in \mathbb{R}^n , not necessarily bounded.
- ▶ $K \subset\subset G$ means K is compact.
- ▶ V a linear space (not necessarily complete) over field \mathbb{K} , operations vector $+$ and scalar \cdot , zero element is θ .
- ▶ Example: (no norm) $V = C(G, \mathbb{K}) = C(G)$.
- ▶ Define with multi-indices: $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j \in \mathbb{N}$,
 $|\alpha| = \sum_{i=1}^n \alpha_i$, $D^\alpha u = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_n}^{\alpha_n} u$.
- ▶ $C^m(G) = \{f \in C(G) : D^\alpha f \in C(G), |\alpha| \leq m\}$.
- ▶ $C^\infty(G) = \bigcap_{m \geq 0} C^m(G)$.
- ▶ *Support* \underline{f} of $f \in C(G)$ is closure of $\{x \in G : f(x) \neq 0\}$.
- ▶ $C_0(G) = \{f \in C(G) : \underline{f} \subset\subset G\}$. $C_0^m(G) = C^m(G) \cap C_0(G)$.
- ▶ $C^m(\overline{G}) = \{f|_{\overline{G}} : f \in C_0^m(\mathbb{R}^n)\}$.

Linear Algebra

- ▶ Subspace $M \leq V$ when closed under operations $+/\cdot$.

- ▶ Examples:

$$C^j(G) \leq C^k(G) \leq \mathbb{K}^G$$

$$C^j(\overline{G}) \leq C^k(\overline{G})$$

$$\{\theta\} \leq C_0^j(G) \leq C_0^k(G) \quad 0 \leq k \leq j \leq \infty$$

$$C_0^k(G) \leq C^k(\overline{G}) \leq C^k(G)$$

- ▶ For $M \leq V$ and $x \in V$, coset $\hat{x} = \{x + m : m \in M\}$,
quotient set $V/M = \{\hat{x} : x \in V\}$ is a linear space:

$$\hat{x} + \hat{y} = \{(x + m_1) + (y + m_2) : m_i \in M\} = \{x + y + m : m \in M\} = \widehat{x + y}$$

$$0\hat{x} := \hat{\theta}, \quad \alpha\hat{x} = \{\alpha(x + m) : m \in M\} = \{\alpha x + m : m \in M\} = \widehat{\alpha x}$$

- ▶ Example: $V = C^1(G)$, $M = \mathbb{K}$, for $f \in C^1(G)$,
 $\hat{f} = \{g \in C^1(G) : g - f \in \mathbb{K}\}$, $V/M = \{\hat{f} : f \in C^1(G)\}$.
- ▶ V, W over \mathbb{K} , $T \in L(V, W)$ (linear) if
 $T(\alpha x + \beta y) = \alpha T(x) + \beta T(y)$, $\forall \alpha, \beta \in \mathbb{K}$, $\forall x, y \in V$.
- ▶ Kernel $K(T) = \{x \in V : Tx = \theta\}$.
- ▶ Identity $i_M : M \rightarrow V$, $i_M(m) = m$, $\forall m \in M$.
- ▶ Quotient map $q_M(x) = \hat{x}$ is a surjection $q_M \in L(V, V/M)$
with $K(q_M) = M$.

Linear Algebra

- ▶ Example: $G = (a, b)$, $d/dx = D : V \rightarrow C(\overline{G})$.
 - ▶ $V = C^1(\overline{G}) \Rightarrow D$ is linear surjection, $K(D) = \text{constants}$.
 - ▶ $V = \{\varphi \in C^1(\overline{G}) : \varphi(a) = 0\} \Rightarrow D$ isomorphism (linear bijection).
 $f \in C(\overline{G}), \varphi(x) = \int_a^x f \in V, D\varphi = f$
 - ▶ $V = \{\varphi \in C^1(\overline{G}) : \varphi(a) = \varphi(b) = 0\}$
 $\Rightarrow \text{Rg}(D) = \{f \in C(\overline{G}) : \int_a^b f = 0\}$.
- ▶ V, W over \mathbb{K} , $T : V \rightarrow W$ conjugate linear if
 $T(\alpha x + \beta y) = \overline{\alpha}T(x) + \overline{\beta}T(y), \alpha, \beta \in \mathbb{K}, x, y \in V$.
 $(\mathbb{K} = \mathbb{R} \Rightarrow \overline{\alpha} = \alpha, \overline{\beta} = \beta)$
- ▶ Algebraic dual of V is $V^* = \{T : V \rightarrow \mathbb{K}, T \text{ conjugate linear}\}$.
- ▶ $Bf = f^*, f^*(x) = \overline{f(x)}$, is bijection $B : L(V, \mathbb{K}) \rightarrow V^*$.
- ▶ Example: Define $T : C_0(G) \rightarrow C_0(G)^*$ through
 $(Tf)(\varphi) = \int_G f\overline{\varphi}, f, \varphi \in C_0(G)$.
A linear injection but not surjective: For $x_0 \in G$ define
 $\delta_{x_0}(\varphi) = \overline{\varphi(x_0)}, \varphi \in C_0(G)$
and $\delta_{x_0} \neq T_f$ for any $f \in C(G)$. Why?

Convergence and Continuity

Def: On a linear space V , a *seminorm* $p : V \rightarrow \mathbb{R}$ satisfies

- $p(x + y) \leq p(x) + p(y), \forall x, y \in V,$
- $p(\alpha x) = |\alpha|p(x), \forall x \in V, \forall \alpha \in \mathbb{K},$

and (V, p) is a *seminormed space*. If p is a *norm* it also satisfies $p(x) > 0, \forall x \neq \theta$, and (V, p) is a *normed space*.

- ▶ Example: For each $K \subset\subset G$, $P_K(f) = \sup\{|f(x)| : x \in K\}$ is a seminorm on $C(G)$ while $P_G = P_{\overline{G}}$ is a norm on $C(\overline{G})$.
- ▶ Convergence $x_n \rightarrow x$ means $p(x_n - x) \rightarrow 0$.
- ▶ Closure of $S \subset V$ is $\overline{S} = \{x \in V : \exists \{x_n\} \subset S \ni x_n \rightarrow x\}$.
- ▶ Dense if $\overline{S} = V$.
- ▶ $T : (V, p) \rightarrow (W, q)$ continuous $\Leftrightarrow x_n \xrightarrow{(V,p)} x \Rightarrow Tx_n \xrightarrow{(W,q)} Tx$.
- ▶ $|p(x) - p(y)| \leq p(x - y) \Rightarrow p : V \rightarrow \mathbb{R}$ continuous.
- ▶ p stronger than q on V : $p(x_n) \rightarrow 0 \Rightarrow q(x_n) \rightarrow 0, \{x_n\} \subset V$.
- ▶ $\mathcal{L}(V, W) = \{T \in L(V, W) \text{ continuous}\}$.

Convergence and Continuity

Theorem: Given seminormed spaces (V, p) and (W, q) ,
 $T \in \mathcal{L}(V, W) \Leftrightarrow \exists K \geq 0 \ni q(T(x)) \leq Kp(x), \forall x \in V.$

Proof: Assume $T \in \mathcal{L}(V, W)$. If there is no K above, then
 $\forall n \geq 1, \exists x_n \in V \ni q(T(x_n)) > np(x_n)$, or with $y_n = x_n/q(T(x_n))$,
 $q(T(y_n)) = 1$ and $p(y_n) \rightarrow 0$, which contradicts $T \in \mathcal{L}(V, W)$.

Assume $\exists K$ above. With $x_n \xrightarrow{(V,p)} x \in V$, $T \in \mathcal{L}(V, W)$ follows
from $q(Tx - Tx_n) = q(T(x - x_n)) \leq Kp(x - x_n) \rightarrow 0$. ■

Theorem: Let (V, p) and (W, q) be seminormed spaces. For
(nontrivial) $T \in \mathcal{L}(V, W)$,

$$|T|_{p,q} = \sup\{q(T(x)) : x \in V, p(x) \leq 1\}$$

satisfies:


$$\begin{aligned} |T|_{p,q} &= \sup\{q(T(x)) : x \in V, p(x) = 1\} && (\equiv N_1) \\ &= \inf\{K > 0 : q(T(x)) \leq Kp(x), \forall x \in V\} && (\equiv N_2) \end{aligned}$$

and $q(T(x)) \leq |T|_{p,q}p(x) \Rightarrow |\cdot|_{p,q}$ is a norm when q is.

Convergence and Continuity

Proof: Let K satisfy $q(T(x)) \leq Kp(x)$. Then $\forall x \in V \ni p(x) \leq 1$, $q(T(x)) \leq K$. Hence, $|T|_{p,q} \leq K$. K arbitrary $\Rightarrow |T|_{p,q} \leq N_2$. If $p(x) > 0$, then $y = x/p(x)$ satisfies $p(y) = 1$, so $p(y) \leq 1 \Rightarrow q(T(y)) \leq |T|_{p,q}$, i.e., for $p(x) > 0$,

$$q(T(x)) \leq |T|_{p,q}p(x). \quad (*)$$

From Theorem , $(*)$ holds trivially when $p(x) = 0$. Hence, $N_2 \leq |T|_{p,q}$ and thus $N_2 = |T|_{p,q}$.

Next, $\{x \in V : p(x) = 1\} \subset \{x \in V : p(x) \leq 1\} \Rightarrow N_1 \leq |T|_{p,q}$. Let $\{x_n\}$ be chosen with $p(x_n) \leq 1$ so that $q(T(x_n)) \rightarrow |T|_{p,q}$. From $(*)$ and $|T|_{p,q} > 0$ it can be assumed that $p(x_n) \neq 0$, so $y_n = x_n/p(x_n)$ satisfies $p(y_n) = 1$ and $q(T(y_n)) \leq N_1$. So $|T|_{p,q} \leftarrow q(T(x_n)) \leq N_1 p(x_n) \leq N_1$. Thus, $N_1 = |T|_{p,q}$. ■

- ▶ T is a *contraction* if $|T|_{p,q} \leq 1$, an *isometry* if $|T|_{p,q} = 1$.
- ▶ *Dual* of (V, p) is $V' = \{f \in V^* : \bar{f} \in \mathcal{L}(V, \mathbb{K})\}$ and
$$\|f\|_{V'} = \sup\{|f(x)| : x \in V, p(x) \leq 1\}$$

Completeness

- ▶ $\{x_n\} \subset (V, p)$ is *Cauchy* if $p(x_n - x_m) \rightarrow 0, m, n \rightarrow \infty$.
- ▶ (V, p) *complete* if every Cauchy sequence converges in V .

Def: A complete normed linear space is a *Banach space*.

- ▶ Example: For $K \subset \subset G$ and $P_K(f) = \sup\{|f(x)| : x \in K\}$, $(C(G), P_K)$ and $(C(\overline{G}), P_G)$ are complete.
- ▶ Example: For $G = (0, 1)$ and $p(x) = \int_G |x|$, $(C(\overline{G}), p)$ is not complete. Why?
- ▶ Strategy for completing a (semi)normed linear space:
 - ▶ Show that M a subspace of seminormed $(V, p) \Rightarrow \overline{M}$ also a subspace of (V, p) .
 - ▶ For seminormed (V, p) and normed (W, q) any continuous extension $T_e : \overline{D} \rightarrow W$ of $T : D \subset V \rightarrow W$ is unique.
 - ▶ For seminormed (V, p) and normed (W, q) , $\exists!$ extension $T_e \in \mathcal{L}(\overline{D}, W)$ of $T \in \mathcal{L}(D, W)$ when $D \leq V$.
 - ▶ A completion of seminormed (V, p) is (W, q) where $W = \{\text{Cauchy sequences in } V\}$ and $q(\{x_n\}) = \lim p(x_n)$.
 - ▶ A completion of normed (V, p) is $(W/K(q), \hat{q})$ where for $W/K(q) \ni \hat{x} = \{x + k : k \in K(q)\}$, $\hat{q}(\hat{x}) = \inf\{q(y) : y \in \hat{x}\}$.

Completeness

Lemma: If M is a subspace of seminormed (V, p) , then \overline{M} is also a subspace of (V, p) .

Proof: If $x, y \in \overline{M}$, $\exists x_n, y_n \in M \ni x_n \xrightarrow{(V,p)} x$ and $y_n \xrightarrow{(V,p)} y$. Then $p((x + y) - (x_n + y_n)) \leq p(x - x_n) + p(y - y_n) \Rightarrow (x_n + y_n) \xrightarrow{(V,p)} (x + y)$. Also $(x_n + y_n) \in M, \forall n \Rightarrow (x + y) \in \overline{M}$. Similarly, $\forall \alpha \in \mathbb{K}$, $p(\alpha x - \alpha x_n) = |\alpha|p(x - x_n) \rightarrow 0 \Rightarrow \alpha x_n \xrightarrow{(V,p)} \alpha x \Rightarrow \alpha x \in \overline{M}$. ■

Lemma: For seminormed (V, p) , normed (W, q) and $T : D \subset V \rightarrow W$, there is at most one continuous $T_e : \overline{D} \rightarrow W$ satisfying $T_e|_D = T$.

Proof: Suppose $\exists T_1, T_2 : \overline{D} \rightarrow W$, each continuous and $T_1|_D = T = T_2|_D$. Then $x \in \overline{D} \Rightarrow \exists \{x_n\} \subset D \ni x_n \xrightarrow{(V,p)} x$. By continuity, $T_1 x \xleftarrow{(W,q)} T_1 x_n = T_2 x_n \xrightarrow{(W,q)} T_2 x$, and $q(T_1 x - T_2 x) = 0 \Rightarrow T_1 x = T_2 x$. ■

Completeness

Theorem: Given seminormed (V, ρ) , Banach space (W, q) and $T \in \mathcal{L}(D, W)$, $D \leq V$, $\exists! T_e \in \mathcal{L}(\overline{D}, W)$ satisfying $T_e|_D = T$ and $|T_e|_{\rho, q} = |T|_{\rho, q}$.

Proof: Uniqueness follows from the above Lemma. For $x \in \overline{D}$, $\exists \{x_n\} \subset D \xrightarrow{(V, \rho)} x$ and $\{x_n\}$ is Cauchy in (V, ρ) . Then by Theorem 1, $q(Tx_m - Tx_n) \leq |T|_{\rho, q} \rho(x_m - x_n) \Rightarrow \{Tx_n\}$ is Cauchy in (W, q) with limit $y \in W$. Similarly, for any other $\{x'_n\} \subset D$ with $x'_n \xrightarrow{(V, \rho)} x$, $Tx'_n \xrightarrow{(W, q)} y'$. Then continuity, $y = y'$, follows from Theorem 1 and $q(y - y') = \lim q(Tx_n - Tx'_n) \leq |T|_{\rho, q} \rho(x_n - x'_n) \rightarrow 0$. So define unambiguously $T_e : \overline{D} \rightarrow W$ through $T_e x = y$ and otherwise $T_e|_D = T$. Then, $\forall x, x' \in \overline{D}$, $T_e(x + x') = \lim T(x_n + x'_n) = \lim Tx_n + \lim Tx'_n = T_e x + T_e x'$ and $T_e \alpha x = \lim T \alpha x_n = \alpha \lim Tx_n = \alpha T_e x$ mean that T_e is linear. Since $D \subset \overline{D}$, $|T|_{\rho, q} \leq |T_e|_{\rho, q}$. Then $\forall x \in \overline{D}$, $q(T_e(x)) = \lim q(T(x_n)) \leq |T|_{\rho, q} \lim \rho(x_n) = |T|_{\rho, q} \rho(x)$ implies through Theorem 1 that $|T_e|_{\rho, q} \leq |T|_{\rho, q}$. Thus, $|T_e|_{\rho, q} = |T|_{\rho, q}$. ■

Completeness

Def: A *completion* of seminormed (normed) (V, p) is a complete seminormed (normed, i.e., Banach space) (W, q) and a linear injection $T : V \rightarrow W$ for which $\overline{\text{Rg}(T)} = W$ and $q(Tx) = p(x), \forall x \in V$.

Theorem: Every seminormed space (V, p) has a completion (W, q) .

Proof: Define $W = \{\{x_n\} \subset V : \{x_n\} \text{ Cauchy}\}$, a seminorm $q(\{x_n\}) = \lim p(x_n)$ and a linear injection $T : (V, p) \rightarrow (W, q)$ by $Tx = \{x, x, \dots\}$ satisfying $q(Tx) = \lim p((Tx)_n) = p(x), \forall x \in V$. For $\{x_n\} \in W$, $q(\{x_n\} - T(x_N)) = \lim p(x_n - x_N)$ is arbitrarily small for N sufficiently large, so $\overline{\text{Rg}(T)} = W$. For completeness, let $\{y^n\}$ be Cauchy in W . Pick $x_n \in V$ with $q(y^n - T(x_n)) < \frac{1}{n}$. Define $y^0 = \{x_1, x_2, \dots\}$. Then $y^0 \in W$ since $p(x_m - x_n) = q(Tx_m - Tx_n) \leq q(Tx_m - y^m) + q(y^m - y^n) + q(y^n - Tx_n) \leq \frac{1}{m} + q(y^m - y^n) + \frac{1}{n} \rightarrow 0$. Also $y^n \xrightarrow{(W, q)} y^0$ since $q(y^n - y^0) \leq q(y^n - Tx_n) + q(Tx_n - y^0) < \frac{1}{n} + \lim_{m \rightarrow \infty} p(x_n - x_m) \xrightarrow{n \rightarrow \infty} 0$. ■

Completeness

Theorem: For a seminormed (V, p) and a subspace $M \leq V$ define $\hat{p}(\hat{x}) = \inf\{p(y) : y \in \hat{x}\}$. Then

- \hat{p} is a seminorm for V/M , and the quotient map $q_M : V \rightarrow V/M$ satisfies $|q_M|_{p, \hat{p}} = 1$.
- $\overline{D} = V \Rightarrow \widehat{\overline{D}} = V/M$ for $\widehat{D} = \{\hat{x} : x \in D\}$.
- \hat{p} is a norm $\Leftrightarrow M$ is closed in V .
- (V, p) complete $\Rightarrow (V/M, \hat{p})$ complete.

Proof: (a) For $\epsilon > 0$ choose $u_\epsilon \in \hat{x}$ and $v_\epsilon \in \hat{y} \ni$
 $p(u_\epsilon) \leq \epsilon + \hat{p}(\hat{x}), \quad p(v_\epsilon) - \hat{p}(\hat{y}) \leq \epsilon.$

So

$$\begin{aligned} \hat{p}(\hat{x} + \hat{y}) &= \hat{p}(\widehat{x + y}) = \inf\{p(x + y + m) : m \in M\} \\ &\leq p(u_\epsilon + v_\epsilon) \leq p(u_\epsilon) + p(v_\epsilon) \leq \hat{p}(\hat{x}) + \hat{p}(\hat{y}) + 2\epsilon \end{aligned}$$

Similarly,

$$\begin{aligned} \hat{p}(\alpha\hat{x}) &= \hat{p}(\widehat{\alpha x}) = \inf\{p(\alpha x + m) : m \in M\} \\ &\leq |\alpha| \inf\{p(x + m) : m \in M\} \leq |\alpha| p(u_\epsilon) \leq |\alpha| [\hat{p}(\hat{x}) + \epsilon] \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, \hat{p} is a seminorm.

Completeness

Since $y \in \hat{x} \Rightarrow y = x + m, m \in M$, it follows with

$$p(y) = p(x + m) \leq p(x) + p(m)$$

and $q_M(x) = \hat{x}$ that

$$\begin{aligned} |q_M|_{p, \hat{p}} &= \sup\{\hat{p}(q_M(x)) : x \in V, p(x) = 1\} \\ &= \sup\{\inf\{p(y) : y \in \hat{x}\} : x \in V, p(x) = 1\} = 1. \end{aligned}$$

(b) For $x \in V$ let $q_M(x) = \hat{x} \in V/M$. Since $\overline{D} = V, \exists\{x_n\} \subset D \ni x_n \xrightarrow{(V,p)} x$ and $q_M(x_n) = \hat{x}_n \in \hat{D}$. Then

$$\begin{aligned} \hat{p}(\hat{x} - \hat{x}_n) &= \hat{p}(q_M(x) - q_M(x_n)) = \hat{p}(q_M(x - x_n)) \\ &\leq p(x - x_n) \rightarrow 0. \end{aligned}$$

Thus $\overline{\hat{D}} = V/M$.

(c) Note that $0 = \hat{p}(\hat{x}) = \inf\{p(x - m) : m \in M\} \Leftrightarrow x \in \overline{M}$.

If \hat{p} is a norm, then $\hat{p}(\hat{x}) = 0 \Rightarrow \hat{x} = \hat{\theta} \Rightarrow x \in M$; in particular, $x \in \overline{M} \Rightarrow \hat{p}(\hat{x}) = 0 \Rightarrow x \in M$, so $\overline{M} = M$. If $\overline{M} = M$, then $\hat{p}(\hat{x}) = 0 \Rightarrow x \in \overline{M} = M \Rightarrow \hat{x} = \hat{\theta}$ means \hat{p} is a norm.



Completeness

(d) Since a Cauchy sequence converges when a subsequence converges, choose a (sub)sequence $\{\hat{x}_n\} \subset V/M$ for which $\hat{p}(\hat{x}_{n+1} - \hat{x}_n) < 2^{-(n+1)}$. Then set $y_1 = x_1$ and for given $m_n, n \geq 1$, $m_1 = \theta$, choose $m_{n+1} \ni p((x_{n+1} + m_{n+1}) - (x_n + m_n)) \leq 2^{-(n+1)} + \hat{p}(\hat{x}_{n+1} - \hat{x}_n)$. Then $y_{n+1} = x_{n+1} + m_{n+1}$ satisfies $p(y_{n+1} - y_n) \leq 2^{-n}$. For $m \geq n$, $p(y_m - y_n) \leq \dots \leq \sum_{k=n}^{m-1} 2^{-k} < 2^{1-n}$. So $\{y_n\}$ is Cauchy in (V, p) . If (V, p) is complete, $\exists y \in V \ni p(y - y_n) \rightarrow 0$ and $\hat{y} \in V/M$. Then $\hat{x}_n \rightarrow \hat{y}$ since $\hat{x}_n = \hat{y}_n$ and $\hat{p}(\hat{y} - \hat{y}_n) = \hat{p}(q_M(y - y_n)) \leq p(y - y_n) \rightarrow 0$. ■

Theorem: Every normed space (V, p) has a completion $(W/K(q), \hat{q})$, where (W, q) is the seminormed space given by Theorem [▶](#) and $\hat{q}(\hat{y}) = \inf\{q(z) : z \in \hat{y}\}$.

Proof: Suppose $K = K(q) \supset \{y_n\} \xrightarrow{(W, q)} y \in W$. By continuity of seminorms, $q(y) = \lim q(y_n) = 0$ and K is closed. By part (c) of Theorem [▶](#), \hat{q} is a norm on W/K . Since (W, q) is complete, $(W/K, \hat{q})$ is a Banach space according to part (d) of


Completeness

Theorem . It was shown for Theorem  that $\overline{\text{Rg}(T)} = W$ and

$$q(Tx) = p(x), \quad \forall x \in V.$$

With the quotient map $q_K = Q \in \mathcal{L}(W, W/K)$, it remains to show that $Q \circ T \in \mathcal{L}(V, W/K)$ is a linear injection for which $\text{Rg}(Q \circ T)$ is dense in W/K and $\hat{q}((Q \circ T)(x)) = p(x), \forall x \in V$. With $D = \text{Rg}(T)$ and

$$\hat{D} = \{T(x) + k : x \in V, k \in K\} = \text{Rg}(Q \circ T).$$

it follows with part (b) of Theorem ,

$$\overline{D} = \overline{\text{Rg}(T)} = W \quad \Rightarrow \quad \overline{\hat{D}} = \overline{\text{Rg}(Q \circ T)} = W/K.$$

Since $\forall k \in K$,


$$q(y) = q(y) - q(k) \leq q(y + k) \leq q(y) + q(k) = q(y)$$

the quotient map satisfies

$$\hat{q}(Q(y)) = \inf\{q(y + k) : k \in K\} = q(y).$$

Combining the above gives:

$$\hat{q}(Q(Tx)) = q(Tx) = p(x), \quad \forall x \in V.$$

Thus, $K(Q \circ T) \leq K(p)$. If p is a norm, then $K(p) = \{\theta\}$ and $K(Q \circ T) = \{\theta\} \Rightarrow Q \circ T$ is injective. 

Completeness

Theorem: For a seminormed (V, ρ) and a Banach space (W, q) , $\mathcal{L}(V, W)$ is a Banach space. In particular, the dual V' of a seminormed (V, ρ) is complete.

Proof: Let $\{T_n\}$ be Cauchy in $\mathcal{L}(V, W)$. Then due to

$$q(T_mx - T_nx) \leq |T_m - T_n|_{p,q} \rho(x)$$

$\{T_nx\}$ is Cauchy in W , $\forall x \in V$, with a unique limit $Tx \in W$, which defines $T : V \rightarrow W$. Also,

$$T(x + x') = \lim T_n(x + x') = \lim T_nx + \lim T_nx' = Tx + Tx'$$

and
$$T\alpha x = \lim T_n\alpha x = \alpha \lim T_nx = \alpha Tx$$

mean that $T \in L(V, W)$. From

$$q(Tx) \leftarrow q(T_nx) \leq |T_n|_{p,q} \rho(x) \leq \rho(x) \sup\{|T_n|_{p,q} : n \geq 1\}$$

It follows from Theorem  that $T \in \mathcal{L}(V, W)$ with

$|T|_{p,q} \leq \sup\{|T_n|_{p,q} : n \geq 1\}$. Finally, let m, n be large enough,

$$q(Tx - T_nx) \xleftarrow{m \rightarrow \infty} q(T_mx - T_nx) \leq |T_m - T_n|_{p,q} \rho(x) \leq \epsilon \rho(x)$$

to obtain $|T - T_n|_{p,q} \leq \epsilon$, so $T_n \xrightarrow{\mathcal{L}(V,W)} T$. ■

Hilbert Space

- ▶ Given a linear space V , a *scalar product*, $(x, y) \in \mathbb{K}$, $x, y \in V$, satisfies:
 - ▶ $x \mapsto (x, y)$ is linear $\forall y \in V$
 - ▶ $(x, y) = (y, x)$, $\forall x, y \in V$
 - ▶ $(x, x) > 0$, $\forall x \neq \theta$
 - ▶ It follows: $y \mapsto (x, y)$ is conjugate linear $\forall x \in V$
- ▶ The pair $V, (\cdot, \cdot)$ is a *scalar product space*.

Theorem: For a scalar product space $V, (\cdot, \cdot)$, the scalar product satisfies:

- Cauchy-Schwarz: $|(x, y)| \leq \|x\| \|y\|$, $\forall x, y \in V$
- $\|x\| = (x, x)^{1/2}$ is a norm on V satisfying the *parallelogram law*: $\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$
- $(\cdot, \cdot) : V \times V \rightarrow \mathbb{K}$ is continuous.

Proof: Part (a) follows with $\alpha = -\overline{(x, y)}$ and $\beta = (x, x)$ in $0 \leq (\alpha x + \beta y, \alpha x + \beta y) = \beta[\beta(y, y) - |\alpha|^2]$

Hilbert Space

For part (b), $(x, x) > 0, \forall x \neq \theta, (\alpha x, \alpha x) = |\alpha|^2(x, x)$ and with (a),

$$\|x + y\|^2 \leq \|x\|^2 + 2|(x, y)| + \|y\|^2 \leq (\|x\| + \|y\|)^2$$

A direct calculation gives the parallelogram law. For part (c),

$$|(x, y) - (x_n, y_n)| \leq \|x\| \|y - y_n\| + \|y_n\| \|x - x_n\| \rightarrow 0$$


for sequences $x_n \xrightarrow{V, \|\cdot\|} x$ and $y_n \xrightarrow{V, \|\cdot\|} y$. ■

Def: A *Hilbert Space* is a complete scalar product space. A *completion* (H, q) of a scalar product space is a completion of the same normed space for which $\exists(\cdot, \cdot)_H$ satisfying $q(\cdot)^2 = (\cdot, \cdot)_H$.

- ▶ Example: $V = C_0(G)$ with $(\varphi, \psi) = \int_G \varphi \bar{\psi}$ (Riemann integral) is a scalar product space but not a Hilbert space.
- ▶ Example: $L^2(G)$ is defined directly as L^2/M_0 , where L^2 consists of Lebesgue square-summable \mathbb{K} -valued functions and M_0 consists of Lebesgue measurable functions vanishing except on a set of measure zero. With $(\varphi, \psi) = \int_G \varphi \bar{\psi}$ (Lebesgue integral), $L^2(G)$ is a Hilbert space. (Compare with the completion below.)

Hilbert Space

Theorem: Every scalar product space has a completion which is a Hilbert space.


Proof: Let $V, (\cdot, \cdot)_V$ be a scalar product space with $\|\cdot\|_V = (\cdot, \cdot)^{1/2}$. Since $(V, \|\cdot\|_V)$ is a normed linear space, let $(H, \|\cdot\|_H)$ denote the completion of $(V, \|\cdot\|_V)$ given according to Theorem . Specifically, $H = W/K$, $K = K(\|\cdot\|_W)$ where $W = \{\{x_n\} \subset V : \{x_n\} \text{ Cauchy}\}$ and for $x = \{x_n\} \in W$, $\|x\|_W = \lim \|x_n\|_V$. Also, with the quotient map $q_K : W \rightarrow W/K$ and for $\hat{x} = q_K(x)$,

$$\|\hat{x}\|_H = \inf\{\|x + k\|_W : k \in K\} =$$

$$\inf\{\lim \|x_n + k_n\|_V : \lim \|k_n\|_V = 0\} = \lim \|x_n\|_V = \|x\|_W$$

So for $\hat{x}, \hat{y} \in H$, where similarly $\hat{y} = q_K(y)$ and $y = \{y_n\} \in W$, define a prospective scalar product on H by:

$$(\hat{x}, \hat{y})_H = \lim (x_n, y_n)_V$$

so that $(\hat{x}, \hat{x})_H = \|\hat{x}\|_H^2$. This limit is well-defined since, by Cauchy-Schwarz, $\{(x_n, y_n)\}$ is Cauchy in \mathbb{K} as $\{x_n\}$ and $\{y_n\}$ are Cauchy in V . By Theorem  $\|\cdot\|_H$ is a norm. It follows

Hilbert Space

that $(\hat{x}, \hat{x})_H = \|\hat{x}\|_H^2 > 0, \forall \hat{x} \neq \hat{\theta}$. Also,

$$\begin{aligned} |(\hat{x}, \hat{y})_H - \overline{(\hat{y}, \hat{x})_H}| &= |\lim(x_n, y_n)_V - \overline{\lim(y_n, x_n)_V}| = \\ &= |\lim(x_n, y_n)_V - \lim(\overline{y_n, x_n})_V| = |\lim(x_n, y_n)_V - \lim(x_n, y_n)_V| = 0 \end{aligned}$$


Furthermore,

$$\begin{aligned} |(\hat{x} + \hat{y}, \hat{z})_H - (\hat{x}, \hat{z})_H - (\hat{y}, \hat{z})_H| &= \\ |\lim(x_n + y_n, z_n)_V - \lim(x_n, z_n)_V - \lim(y_n, z_n)_V| &= 0 \end{aligned}$$

and

$$|(\alpha\hat{x}, \hat{z})_H - \alpha(\hat{x}, \hat{z})_H| = |\lim(\alpha x_n, z_n)_V - \alpha \lim(x_n, z_n)_V| = 0$$

So $\hat{x} \mapsto (\hat{x}, \hat{y})_H$ is linear $\forall \hat{y} \in H$. Thus, $(\cdot, \cdot)_H$ is a scalar product, and H is a Hilbert space. ■

Study Question: Prove Theorem  using the following Lemma: If V is a normed space whose norm $\|\cdot\|$ satisfies the *parallelogram law*, then the following *polarization identity* defines an inner product on V :

$4(x, y) = \|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2$
satisfying $(x, x) = \|x\|^2$. Complex terms drop for $\mathbb{K} = \mathbb{R}$.

Hilbert Space

- ▶ Example: $L^2(G)$ is the completion of $C_0(G)$ equipped with $(\varphi, \psi) = \int_G \varphi \overline{\psi}$ (Riemann integral). In this way, the elements of $L^2(G)$ are identified as sequences of $C_0(G)$ -functions which are Cauchy with respect to $(\cdot, \cdot)^{1/2}$, and such sequences are identified when their difference converges to zero.
- ▶ Using V or H for a Hilbert space and, e.g., subscript V in $(x, y)_V$ and $\|x\|_V$ dropped when unambiguous.
- ▶ An angle α between $x, y \in V$ is given by
$$(x, y) = \|x\| \|y\| \cos(\alpha).$$
- ▶ $x, y \in V$ are *orthogonal* when $(x, y) = 0$.
- ▶ For a subspace $M \leq V$, the *orthogonal complement* is
$$M^\perp = \{x \in V : (x, y) = 0, \forall y \in M\}.$$

Lemma: M^\perp is a closed subspace of V and $M \cap M^\perp = \{0\}$.

Proof: Suppose $\exists \{x_n\} \subset M^\perp \ni x_n \rightarrow x \in V \setminus M^\perp$. Since

Hilbert Space

$x \notin M^\perp, \exists y \in M \ni (x, y) \neq 0$. However, this contradicts

$$|(x, y)| = |(x, y) - (x_n, y)| \leq \|x - x_n\| \|y\| \rightarrow 0.$$

Thus, M^\perp is closed in V . Then if $x \in M \cap M^\perp$, it follows from $(x, x) = 0$ that $x = \theta$. ■

- ▶ $K \subset V$ is *convex* if $\alpha x + (1 - \alpha)y \in K, \forall x, y \in K, \forall \alpha \in [0, 1]$.

Theorem: A non-empty closed convex subset of a Hilbert space H has a unique element of minimal norm.


Proof: Let $\{x_n\} \subset K$ be a sequence chosen so that $\|x_n\|$ converges to $d = \inf\{\|x\| : x \in K\}$. Since K is convex, $\frac{1}{2}(x_n + x_m) \in K, \forall m, n$. Hence, $\|x_n + x_m\|^2 \geq 4d^2$. By the parallelogram law,

$$\|x_n - x_m\|^2 \leq 2(\|x_n\|^2 + \|x_m\|^2) - 4d^2 \rightarrow 0.$$

Hence, $\{x_n\}$ is Cauchy and converges to some $x \in H$. Since K is closed, $x \in K$. By continuity of the scalar product and hence norm, $\|x\| = \lim \|x_n\| = d$. If also $\|x'\| = d$, then $\frac{1}{2}(x + x') \in K$ and $d^2 \leq \|\frac{1}{2}(x + x')\|^2 = d^2 - \frac{1}{4}\|x - x'\|^2 \Rightarrow \|x - x'\| = 0$. ■

Hilbert Space

Theorem: Let M be a closed subspace of the Hilbert space H . Then $\forall x \in H$, $x = m + n$ where $m \in M$ and $n \in M^\perp$ are uniquely determined by x .


Proof: Uniqueness follows since $x = m_i + n_i$, $i = 1, 2$, $m_i \in M$, $n_i \in M^\perp$ means $m_2 - m_1 = n_1 - n_2 \in M \cap M^\perp = \{\theta\}$. For existence, define $K = \{x + y : y \in M\}$ and use Theorem  to obtain $n \in K$ with $\|n\| = \inf\{\|x + y\| : y \in M\}$. Then take $m = x - n$. Since $n = x + z$ for some $z \in M$, $m = x - (x + z) = -z \in M$. To show that $n \in M^\perp$, let $y \in M$. Then $\forall \alpha \in \mathbb{K}$, $n - \alpha y = x + (z - \alpha y) \in K$ and hence

$\|n - \alpha y\|^2 \geq \|n\|^2$ or

$$\|n\|^2 - 2\Re(n, \alpha y) + |\alpha|^2 \|y\|^2 \geq \|n\|^2$$

With $\alpha = \beta(n, y)$, $\beta > 0$,

$$-2\beta |(n, y)|^2 + |\beta|^2 |(n, y)|^2 \|y\|^2 \geq 0.$$

or $|(n, y)|^2 (\beta \|y\|^2 - 2) \geq 0$, which holds $\forall \beta > 0$ only if $(n, y) = 0$. 

Hilbert Space

Def: Given a Hilbert space H , a closed subspace $M \leq H$ and the unique decomposition $x = m + n$, $x \in H$, $m \in M$, $n \in M^\perp$, then $P_M : H \rightarrow M$ given by $P_M x = m$ is a *projection on M* .

▶ A projection P_M is linear:

1. $(x_1 + x_2) = (m_1 + m_2) + (n_1 + n_2) \Rightarrow P_M(x_1 + x_2) = m_1 + m_2 = P_M x_1 + P_M x_2, \forall x_i \in H, i = 1, 2$

2. $\alpha x_1 = \alpha(m_1 + n_1) \Rightarrow P_M(\alpha x_1) = \alpha P_M x_1, \forall x \in H, \forall \alpha \in \mathbb{K}$

▶ $P_M \in \mathcal{L}(H)$ since $(\mathcal{L}(H) = \mathcal{L}(H, H))$

$$\|P_M x\|^2 \leq \|P_M x\|^2 + \|(I - P_M)x\|^2 = \|x\|^2$$

▶ If $P \in \mathcal{L}(B)$ satisfies $P \circ P = P$, then P is called a *projection on the Banach space B* . General existence?

Study Question: With $\ell_1 = \{x = \{x_n\} : \|x\|_1 = \sum |x_n| < \infty\}$ define $M = \{x \in \ell_1 : \sum \frac{n}{n+1} x_n = 0\}$. With $e^m = \{\delta_{nm}\}$, show:

(a) $e^1 - \frac{1}{2} \frac{n+1}{n} e^n \in M$, (b) $\text{dist}(e^1, M) \leq \frac{1}{2}$ and (c) $y \in M \Rightarrow \|e^1 - y\|_1 > \frac{1}{2}$. Hence $\frac{1}{2} = \text{dist}(e^1, M) < \|e^1 - y\|_1, \forall y \in M$.

Hilbert Space

Theorem: (Riesz) Given a Hilbert space H and $f \in H'$, $\exists! x \in H$
 $\ni f(y) = (x, y), \forall y \in H$.

Proof: If $f = \theta_{H'}$, take $x = \theta_H$, so assume $f \neq \theta_{H'}$. Define
 $K = K(f) = \{x \in H : f(x) = 0\} \leq H$ so by continuity of f ,
 $f(y) = \lim f(y_n) = 0$, K is closed with $K^\perp \neq \{\theta\}$. Let $n \in K^\perp$ be
chosen with $\|n\| = 1$. Then $\forall z \in K^\perp, u = \overline{f(n)}z - \overline{f(z)}n$
satisfies $u \in K^\perp$, but also $f(u) = f(n)\overline{f(z)} - \overline{f(z)}f(n) = 0$
(conjugate linear) means $u \in K$, so $u = \theta_H$. Thus, K^\perp is 1D.
Then $\forall y \in H, y = P_K(y) + \lambda n$, where $(y, n) = \lambda(n, n) = \lambda$. But
 $f(y) = f(\lambda n) = \overline{\lambda}f(n) = (n, y)\overline{f(n)}$, and hence $f(y) = (f(n)n, y)$.
If $\exists\{x_1, x_2\} \subset H \ni f(y) = (x_1, y) = (x_2, y), \forall y \in H$, then with
 $y = x_2 - x_1, (x_1 - x_2, x_1 - x_2) = 0 \Rightarrow x_1 = x_2$. ■

Def: The mapping $R_H : H \rightarrow H'$ given by

$$R_H(x)(y) = (x, y), \quad x, y \in H$$

is called the *Riesz map*. Since $\|R_H(x)\|_{H'} = \sup\{|R_H(x)(y)| : \|y\| \leq 1\} = \|x\|$, R_H is an isometry, $\|R_H\|_{H, H'} = 1$. Also, $K(R_H) = \{\theta\}$.


Dual Operators and Identifications

Def: For linear spaces V, W and $T \in L(V, W)$, the *dual operator* $T' \in L(W^*, V^*)$ is $T'(f) = f \circ T, f \in W^*$.

- ▶ If V is a linear space, (W, q) a seminormed space and $T \in \mathcal{L}(V, W)$ has dense range, then T' is injective on W' :
 $f \in W'$ and $0 = T'(f)(v) = f \circ T(v), \forall v \in V \Rightarrow f = \theta_{W'}$
- ▶ If (V, p) and (W, q) are seminormed spaces and $T \in \mathcal{L}(V, W)$, then T' (henceforth understood as restricted to W') satisfies

$$|T'f(x)| \leq \|f\|_{W'} |T|_{p,q} p(x), f \in W', x \in V$$

and hence $T' \in \mathcal{L}(W', V')$ and $\|T'\|_{\mathcal{L}(W', V')} \leq |T|_{p,q}$.

Def: Let V and W be Hilbert spaces and $T \in \mathcal{L}(V, W)$. Let $F \in V'$ be given by $Fv = (u, Tv)_W, u \in W, v \in V$. Then by Theorem  $\exists! T^*u \in V$

$$(T^*u, v)_V = (u, Tv)_W, \quad \forall u \in W, \quad \forall v \in V$$

and T^* is called the *adjoint* of T .

Dual Operators and Identifications

- ▶ Given the Riesz maps R_V and R_W , it follows from $R_V \circ T^*(u)(v) = (T^*u, v)_V = (u, Tv)_W = R_W(u)(Tv) = T' \circ R_W(u)(v)$ that the dual and adjoint are related by $R_V \circ T^* = T' \circ R_W$.

Theorem: If V and W are Hilbert spaces and $T \in \mathcal{L}(V, W)$, then $T^* \in \mathcal{L}(W, V)$, $\text{Rg}(T)^\perp = K(T^*)$ and $\text{Rg}(T^*)^\perp = K(T)$. If T is an isomorphism with $T^{-1} \in \mathcal{L}(W, V)$, then T^* is an isomorphism with $(T^*)^{-1} = (T^{-1})^*$.

Proof: It follows from

$$\begin{aligned}\|T^*w\|_V &= \sup\{(T^*w, v)_V : \|v\| \leq 1\} \\ &= \sup\{(w, Tv)_W : \|v\| \leq 1\} \leq \|w\|_W \|T\|_{\mathcal{L}(V, W)}\end{aligned}$$

that $T \in \mathcal{L}(V, W) \Rightarrow T^* \in \mathcal{L}(W, V)$. Then

$0 = (w, Tv)_W = (T^*w, v)_V, \forall w \in W, \forall v \in V$
means $\text{Rg}(T)^\perp = K(T^*)$ and $\text{Rg}(T^*)^\perp = K(T)$. Finally,
 $T^{-1} \in \mathcal{L}(W, V) \Rightarrow (T^{-1})^* \in \mathcal{L}(V, W)$ and it follows from

$(u, w)_W = (T^{-1}u, T^*w)_V = (u, (T^{-1})^*T^*w)_W, \forall u, w \in W$
that $(T^*)^{-1} = (T^{-1})^* \in \mathcal{L}(V, W)$. ■

Dual Operators and Identifications

Example:

- ▶ Given the linear space $V = C_0(G)$ and the Hilbert space $W = L^2(G)$ equipped with $(f, g) = \int_G f \bar{g}$.
- ▶ For $\varphi \in C_0(G)$, let $i(\varphi)$ denote the $L^2(G)$ equivalence class containing φ . $i : C_0(G) \rightarrow L^2(G)$ is a linear injection with dense range. *Identify* domain with range, write $C_0(G) \leq L^2(G)$.
- ▶ Thus, $i' : L^2(G)' \rightarrow C_0(G)^*$ is injective on $L^2(G)'$.
 $i'(f') = f' \circ i$ is the restriction of $f' \in L^2(G)'$ to $C_0(G) \subset L^2(G)$.
- ▶ *Identify* domain and range of i' , write $L^2(G)' \leq C_0(G)^*$.
- ▶ Let $R = R_{L^2(G)}$ denote the Riesz map and identify $L^2(G)$ with $L^2(G)'$ through Theorem.
- ▶ Recall $T : C_0(G) \rightarrow C_0(G)^*$ defined by $(Tf)(\varphi) = \int_G f \bar{\varphi}$, $f, \varphi \in C_0(G)$. *Identify* domain with range, write $C_0(G) \leq C_0(G)^*$.
- ▶ Summary: $T = i' \circ R \circ i$ and
$$C_0(G) \leq L^2(G) = L^2(G)' \leq C_0(G)^*.$$

Dual Operators and Identifications

Def: A *sesquilinear form* on a linear space V is a function $a : V \times V \rightarrow \mathbb{K}$ such that $x \mapsto a(x, y)$ is linear, $\forall y \in V$, and $y \mapsto a(x, y)$ is conjugate linear, $\forall x \in V$. If $\mathbb{K} = \mathbb{R}$, then a is *bilinear*.

- ▶ There is a one-to-one correspondence between sesquilinear forms a and operators $\mathcal{A} \in L(V, V^*)$ according to $a(x, y) = \mathcal{A}x(y)$, $x, y \in V$.

Theorem: Given a normed space (V, p) and a sesquilinear form a on V , the following are equivalent:

- $a(\cdot, \cdot)$ is continuous at (θ, θ) .
- $a(\cdot, \cdot)$ is continuous on $V \times V$.
- $\exists K > 0 \ni |a(x, y)| \leq Kp(x)p(y)$, $x, y \in V$
- $\mathcal{A} \in \mathcal{L}(V, V')$.

Proof: Clearly: (c) \Leftrightarrow (d), (c) \Rightarrow (b) and (b) \Rightarrow (a). To show (a) \Rightarrow (c), choose K so that $p(\tilde{x}) \leq 1/\sqrt{K}$ and $p(\tilde{y}) \leq 1/\sqrt{K} \Rightarrow |a(\tilde{x}, \tilde{y})| \leq 1$. Then for arbitrary $x, y \in V$, set $\tilde{x} = x/[p(x)\sqrt{K}]$ and $\tilde{y} = y/[p(y)\sqrt{K}]$ to obtain (c). ■

Uniform Boundedness and Weak Compactness

Def: $\{x_n\} \subset H$ is *weakly convergent*, $x_n \xrightarrow{H} x$, if $\lim(x_n, v) = (x, v)$, $\forall v \in H$. (x is unique.) $\{x_n\}$ is *weakly bounded* if $|(x_n, v)|$ is bounded $\forall v \in H$.

Principle of Uniform Boundedness:

Theorem: $\{x_n\} \subset H$ is weakly bounded iff it is bounded.

Proof: Let $\{x_n\}$ be weakly bounded. First to show is, $\exists K, r > 0$, $\exists z \in H \ni |(x_n, y)| \leq K, \forall y \in B(z, r)$. If not, $\exists n_1 \in \mathbb{N}, y_1 \in B(\theta, 1) \ni |(x_{n_1}, y_1)| > 1$. Choose $r_1 < 1 \ni B(y_1, r_1) \subset B(\theta, 1)$ and $|(x_{n_1}, y)| > 1, \forall y \in B(y_1, r_1)$. Similarly, $\exists n_2 > n_1, y_2 \in B(y_1, r_1) \ni |(x_{n_2}, y_2)| > 2$, so choose $r_2 < \frac{1}{2} \ni B(y_2, r_2) \subset B(y_1, r_1)$ and $|(x_{n_2}, y)| > 2, \forall y \in B(y_2, r_2)$. Inductively, define $B(y_k, r_k) \subset B(y_{k-1}, r_{k-1}), r_k < \frac{1}{k} \ni |(x_{n_k}, y)| > k, \forall y \in B(y_k, r_k)$. Since $\|y_m - y_n\| < \frac{1}{n}$ if $m > n$, $\{y_n\}$ is Cauchy and $y_n \xrightarrow{H} y$. But $y \in B(y_k, r_k) \Rightarrow |(x_{n_k}, y)| > k, \forall k$, a contradiction. So, $\forall y \in B(\theta, 1), |(x_n, y)| = \frac{1}{r} |(x_n, z + ry) - (x_n, z)| \leq 2K/r$ und $\|x_n\| = \sup\{|(x_n, y)| : \|y\| \leq 1\} \leq 2K/r, \forall n$. ■

Uniform Boundedness and Weak Compactness

Lemma: Suppose $\{x_n\}$ is bounded in H and $D \subset H$ is dense.

Then $\lim(x_n, v) = (x, v), \forall v \in D \Leftrightarrow x_n \xrightarrow{H} x$.

Proof: Let $\epsilon > 0$ and fix $v \in H$. Then $\exists z \in D$ with $\|v - z\| < \epsilon$. Thus, $|(x_n - x, v)| \leq |(x_n, v - z)| + |(z, x_n - x)| + |(x, v - z)| < \epsilon\|x_n\| + |(z, x_n - x)| + \epsilon\|x\|$. Hence, $\forall n$ large enough, $|(x_n - x, v)| < 3\epsilon \sup\{\|x_m\| : m \geq 1\}$. ■

Theorem: Let the Hilbert space H have a countably dense subset $D = \{y_n\}$. If $\{x_n\} \subset H$ is bounded, then it has a weakly convergent subsequence.

Proof: Since $\{(x_n, y_1)\}$ is bounded in \mathbb{K} , there is a subsequence $\{x_{1,n}\} \ni \{(x_{1,n}, y_1)\}$ converges. Similarly, for each $j \geq 2$, $\exists \{x_{j,n}\} \subset \{x_{j-1,n}\} \ni \{(x_{j,n}, y_k)\}$ converges in \mathbb{K} for $1 \leq k \leq j$. It follows that $\{x_{n,n}\}$ is a subsequence for which $\{(x_{n,n}, y_k)\}$ converges for every $k \geq 1$. With the *span*, $\langle D \rangle \leq H$, i.e.,

$$\langle D \rangle = \left\{ \sum_{k=1}^K \alpha_k y_k : \alpha_k \in \mathbb{K}, y_k \in D, K \in \mathbb{N} \right\},$$

Uniform Boundedness and Weak Compactness

define $T : \langle D \rangle \rightarrow \mathbb{K}$ by $T(y) = \lim(x_{n,n}, y)$. According to

$$\lim(x_{n,n}, y + y') = \lim(x_{n,n}, y) + \lim(x_{n,n}, y'), \quad \forall y, y' \in \langle D \rangle$$


and


$$\lim(x_{n,n}, \alpha y) = \bar{\alpha} \lim(x_{n,n}, y), \quad \forall \alpha \in \mathbb{K}, \forall y \in \langle D \rangle$$

T is conjugate linear. Also by Theorem  and

$$|T(y)| \leq \sup\{\|x_{n,n}\| : n \geq 1\} \|y\|, \quad \forall y \in \langle D \rangle$$

\bar{T} , and hence T , is continuous on $\langle D \rangle$. Since $\overline{\langle D \rangle} = \bar{D} = H$,

it follows with Theorem , that T has a unique extension

$T_e \in H'$. By Theorem , $\exists! x \in H \ni T_e(y) = (x, y)$, $y \in H$.


Thus, $\lim(x_{n,n}, y) = T(y) = T_e(y) = (x, y)$, $\forall y \in \langle D \rangle$, and with the Lemma above, it follows that x is the weak limit of $\{x_{n,n}\}$. ■

Def: A seminormed space with a countably dense subset is called *separable*. A subset of a seminormed space is relatively *sequentially weakly compact* if every sequence in the subset has a subsequence which converges weakly in the space.

According to Theorem 6.B, every bounded set in a separable Hilbert space is relatively sequentially weakly compact.

Expansion in Eigenfunctions

Def: For a Hilbert space H , a (non-trivial) sequence $\{v_i\} \subset H$ is *orthogonal* when $(v_i, v_j) = 0$, $i \neq j$, and *orthonormal* when $(v_i, v_j) = \delta_{ij}$. The *Fourier coefficients* of $u \in H$ are $c_i = (u, v_i)/(v_i, v_i)$. (H a scalar product space?)

- ▶ For $n \geq 1$, let $M_n = \langle \{v_i\}_{i=1}^n \rangle$ and set $u_n = \sum_{i=1}^n c_i v_i$.
- ▶ By Theorem , $u_n = P_{M_n} u$ since $u = u_n + (u - u_n)$ and $((u - u_n), v_i)_{i=1}^n = 0 \Rightarrow (u - u_n) \in M_n^\perp$.

Theorem: Let $\{v_i\}$ be orthonormal in the Hilbert space H and fix $u \in H$. The Fourier coefficients of u satisfy

$$\sum_{i=1}^{\infty} |c_i|^2 \leq \|u\|^2$$

and equality holds if and only if $u = \sum_{i=1}^{\infty} c_i v_i$.

Proof: Since $u = u_n + (u - u_n)$ and $(u - u_n) \perp u_n$, it follows

$$\|u\|^2 = \|u_n\|^2 + 2\Re(u_n, u - u_n) + \|u - u_n\|^2 = \|u_n\|^2 + \|u - u_n\|^2. \quad (*)$$

So $\|u\|^2 \geq \|u_n\|^2$ and $\|u_n\|^2 = \sum_{i=1}^n |c_i|^2$ imply $\|u\|^2 \geq \sum_{i=1}^n |c_i|^2 \rightarrow \sum_{i=1}^{\infty} |c_i|^2$. Also, equality holds in $(*)$ iff $\lim \|u - u_n\| = 0$. ■

Expansion in Eigenfunctions

Def: *Bessel's Inequality* is $\sum_{i=1}^{\infty} |c_i|^2 \leq \|u\|^2$ and *Parseval's Identity* is $\sum_{i=1}^{\infty} |c_i|^2 = \|u\|^2$. The series $u = \sum_{i=1}^{\infty} c_i v_i$ is the *Fourier Series* of u with respect to the orthonormal $\{v_i\}$. $\{v_i\}$ is a *basis* for H when $\overline{\langle \{v_i\} \rangle} = H$.


Theorem: Let $\{v_i\}$ be orthonormal in the Hilbert space H . Then $\overline{\langle \{v_i\} \rangle} = H \Leftrightarrow u = \sum_{i=1}^{\infty} c_i v_i, \forall u \in H$.

Proof: (\Leftarrow) is clear. For (\Rightarrow), fix $u \in H$ and recall that $u_n = P_{M_n} u$. Then for $\epsilon > 0$, choose $n \geq 1 \ni \|u - u_n\| = \|u - P_{M_n} u\| < \epsilon$. Hence, $\lim u_n = u$. ■

Def: Let $T \in \mathcal{L}(H) = \mathcal{L}(H, H)$ for a Hilbert space H . A non-trivial $v \in H$ is an *eigenvector* of T if $Tv = \lambda v$ for some $\lambda \in \mathbb{K}$, and λ is the corresponding *eigenvalue*. T is *self-adjoint* if $(Tu, v) = (u, Tv)$, $\forall u, v \in H$. A self-adjoint T is *non-negative* if $(Tu, u) \geq 0, \forall u \in H$.

Expansion in Eigenfunctions

Lemma: Let H be a Hilbert space. If $T \in \mathcal{L}(H)$ is non-negative self-adjoint, then $\|Tu\| \leq \|T\|^{\frac{1}{2}}(Tu, u)^{\frac{1}{2}}$, $u \in H$.

Proof: Define $[u, v] = (Tu, v)$. As in the proof of Theorem , take $\alpha = -\overline{[u, v]}$ and $\beta = [u, u]$ in $0 \leq [\alpha u + \beta v, \alpha u + \beta v] = \beta(\beta[v, v] - |\alpha|^2)$ to obtain


$$|[u, v]|^2 \leq [u, u][v, v], \quad u, v \in H,$$

in case $\beta > 0$. If $[v, v] > 0$, the same inequality follows by exchanging u and v in α and β . If $[u, u] = [v, v] = 0$, then $[u, v] = 0$ follows from $0 \leq [u + tv, u + tv] = 2t[u, v]$, $\forall t \in \mathbb{R}$. The claim then follows after setting $v = Tu$ in the inequality. ■

Def: For seminormed (V, p) and (W, q) , $T \in \mathcal{L}(V, W)$ is called *compact* if for any bounded $\{u_n\} \subset V$, $\{Tu_n\} \subset W$ has a convergent subsequence.

Lemma: Let H be a Hilbert space. Suppose $T \in \mathcal{L}(H)$ is self-adjoint and compact. Then $\exists v \ni \|v\| = 1$ and $T(v) = \mu v$ where $|\mu| = \|T\|_{\mathcal{L}(H)}$.

Expansion in Eigenfunctions

Proof: The trivial case is excluded by assuming $\|T\|_{\mathcal{L}(H)} > 0$. If $\lambda = \|T\|_{\mathcal{L}(H)}$, it follows from Theorem , $\exists\{u_n\} \subset H$ with $\|u_n\| = 1$ and $\lim \|Tu_n\| = \lambda$. Then $((\lambda^2 - T^2)u_n, u_n) = \lambda^2 - \|Tu_n\|^2 \rightarrow 0$. Since $\lambda^2 - T^2$ is non-negative and self-adjoint, it follows with the previous Lemma that $(\lambda^2 - T^2)u_n \rightarrow 0$. Since T is compact, there is a subsequence, again denoted for convenience by $\{u_n\}$, for which $\{Tu_n\}$ converges to some $w \in H$. Since T is continuous, it follows, $\lim \lambda^2 u_n = \lim T^2 u_n = Tw$, so $w = \lim Tu_n = \lambda^{-2} T^2(w)$. Note that $\|w\| = \lambda$ and $T^2(w) = \lambda^2 w$. Now if $\alpha = \|(\lambda - T)w\| \neq 0$, set $v = (\lambda - T)w/\alpha$ and $\mu = -\lambda$ so that $Tv = (\lambda T - \lambda^2)w/\alpha = -\lambda v$. Otherwise, if $\alpha = 0$, set $v = w/\|w\|$ and $\mu = \lambda$. ■

Theorem: Suppose H is a Hilbert space and that $T \in \mathcal{L}(H)$ is self-adjoint and compact. Then $\exists\{v_i\}$ orthonormal eigenvectors of T for which $\text{Rg}(T) \subset \overline{\{\{v_i\}\}}$ and the corresponding eigenvalues satisfy $|\lambda_i| \geq |\lambda_{i+1}| \xrightarrow{i \rightarrow \infty} 0$.

Expansion in Eigenfunctions

Proof: By the previous Lemma, $\exists v_1$ with $\|v_1\| = 1$ and $T(v_1) = \lambda_1 v_1$ where $|\lambda_1| = \|T\|_{\mathcal{L}(H)}$. Set $H_1 = \{v_1\}^\perp$ (which is a closed subspace of H and hence a Hilbert space) and note $T(H_1) \subset H_1$ since $(Tu, v_1) = (u, Tv_1) = \lambda_1(u, v_1) = 0, \forall u \in H_1$. So $T|_{H_1} \in \mathcal{L}(H_1)$ is self-adjoint and compact, and the Lemma again gives a $v_2 \in H_1$ with $\|v_2\| = 1$ and $T(v_2) = \lambda_2 v_2$ where $|\lambda_2| = \|T|_{H_1}\|_{\mathcal{L}(H_1)} \leq \|T\|_{\mathcal{L}(H)} = |\lambda_1|$. This is continued to obtain an orthonormal sequence $\{v_i\} \subset H$ and a sequence $\{\lambda_i\} \subset \mathbb{K}$ satisfying $|\lambda_{i+1}| \leq |\lambda_i|, i \geq 1$.

Suppose $\exists n \ni \lambda_i = 0, \forall i \geq n$. Then $0 = |\lambda_n| = \|T\|_{\mathcal{L}(H_{n-1})} \Rightarrow H_{n-1} \subset K(T)$. Also $\langle \{v_i\}_{i=1}^n \rangle \subset \text{Rg}(T)$, so $\text{Rg}(T)^\perp \subset \langle \{v_i\}_{i=1}^n \rangle^\perp = H_{n-1}$, and from Theorem, it follows $K(T) = \text{Rg}(T)^\perp \subset H_{n-1}$, and hence $K(T) = H_{n-1}$. From the sandwich, $\text{Rg}(T)^\perp = \langle \{v_i\}_{i=1}^n \rangle^\perp$ and hence $\overline{\text{Rg}(T)} = \overline{\langle \{v_i\}_{i=1}^n \rangle}$. So $\langle \{v_i\}_{i=1}^n \rangle \subset \text{Rg}(T) \subset \overline{\text{Rg}(T)} = \overline{\langle \{v_i\}_{i=1}^n \rangle} = \langle \{v_i\}_{i=1}^n \rangle$ means $\text{Rg}(T) = \langle \{v_i\}_{i=1}^n \rangle$. So the proof is complete in this case.

Expansion in Eigenfunctions

From now on, assume $|\lambda_i| > 0, \forall i$. It will be shown that

$\lim \lambda_i = 0$. If not, $\exists \epsilon > 0 \ni |\lambda_i| \geq \epsilon, \forall i$. But then $\forall i \neq j$,

$$\|T(v_i) - T(v_j)\|^2 = \|\lambda_i v_i - \lambda_j v_j\|^2 = \|\lambda_i v_i\|^2 + \|\lambda_j v_j\|^2 \geq 2\epsilon^2,$$

so $\{T(v_i)\}$ has no convergent subsequence, contradicting compactness of T .

It will next be shown that $\overline{\langle v_i \rangle}$ contains $\text{Rg}(T)$. Say $w \in \text{Rg}(T)$, so $\exists u \in H \ni T(u) = w$. Define

$$w_n = \sum_{i=1}^n b_i v_i, \quad u_n = \sum_{i=1}^n c_i v_i$$

so that $\lim w_n$ and $\lim u_n$ are the Fourier series for w and u , respectively. The coefficients satisfy

$$b_i = (w, v_i) = (Tu, v_i) = (u, Tv_i) = \bar{\lambda}_i c_i = \lambda_i c_i,$$

so $T(c_i v_i) = b_i v_i$. Hence, $w - w_n = T(u - u_n)$, $n \geq 1$, and

$\|w - w_n\| \leq \|T\|_{\mathcal{L}(H_n)} \|u - u_n\|$ since $(u - u_n, v_i)_{i=1}^n = 0$. With

$\|T\|_{\mathcal{L}(H_n)} \leq |\lambda_{n+1}|$ and $\|u - u_n\|^2 = \|u\|^2 - \|u_n\|^2 \leq \|u\|^2$, it follows

$$\|w - w_n\| \leq |\lambda_{n+1}| \|u\|, \quad n \geq 1.$$

Then $\lim \lambda_i = 0 \Rightarrow w = \lim w_n$ and so $\text{Rg}(T) \subset \overline{\langle \{v_i\} \rangle}$. ■

Distributions

Def: A *mollifier* is a function $\varphi_\epsilon \in C_0^\infty(\mathbb{R}^n)$ satisfying $\forall \epsilon > 0$: $\varphi \geq 0$, $\varphi_\epsilon \subset \overline{B(0, \epsilon)}$ and $\int \varphi_\epsilon = 1$. The standard mollifier is:

$$\psi_\epsilon(x) = \begin{cases} \exp[1/(|x|^2 - \epsilon^2)], & |x| < \epsilon \\ 0, & |x| \geq \epsilon \end{cases}$$

For $G \subset \mathbb{R}^n$ and $f \in L^1(G)$ define

$$\underline{f} = \overline{G} \setminus Z_f, \quad Z_f = \cup \{ \text{open } S \subset G : \int_S f \varphi = 0, \forall \varphi \in C_0^\infty(S) \}$$

Extending $f \rightarrow 0$ in $\mathbb{R}^n \setminus G$, $\underline{f} = Z_f^c$ and the mollification of f is

$$f_\epsilon(x) = [f \star \varphi_\epsilon](x) = \int_{\mathbb{R}^n} f(x-y)\varphi_\epsilon(y)dy, \quad x \in \mathbb{R}^n$$

Lemma: If $f \in L^1(G)$, then $\forall \epsilon > 0$, $\underline{f}_\epsilon \subset \{x \in \mathbb{R}^n : \text{dist}(x, \underline{f}) \leq \epsilon\}$ and $f_\epsilon \in C^\infty(\mathbb{R}^n)$.

Proof: $f_\epsilon \in C^\infty(\mathbb{R}^n)$ follows from $f_\epsilon(x) = [f \star \varphi_\epsilon](x) = [\varphi_\epsilon \star f](x)$

Distributions

$= \int_{\mathbb{R}^n} f(z)\varphi_\epsilon(x-z)dz$. Next, choose $S \subset Z_f$ and set $S_\epsilon = \{x \in \mathbb{R}^n : \text{dist}(x, S^c) > \epsilon\}$. Then fix any mollifier φ_ϵ and define φ_ϵ^- by $\varphi_\epsilon^-(x) = \varphi_\epsilon(-x)$, so $\psi \star \varphi_\epsilon^- \in C_0^\infty(S)$, $\forall \psi \in C_0^\infty(S_\epsilon)$. With Fubini, $0 = \int_S f(z)[\psi \star \varphi_\epsilon^-](z)dz = \int_S f(z)[\int_{S_\epsilon} \psi(x)\varphi_\epsilon(x-z)dx]dz = \int_{S_\epsilon} \psi(x)[\int_S f(z)\varphi_\epsilon(x-z)dz]dx = \int_{S_\epsilon} \psi(x)[f \star \varphi_\epsilon](x)dx = \int_{S_\epsilon} \psi(x)f_\epsilon(x)dx$ and thus $S_\epsilon \subset Z_{f_\epsilon}$. Since $S \subset Z_f$ is arbitrary, $\{x \in \mathbb{R}^n : \text{dist}(x, Z_f^c) > \epsilon\} \subset Z_{f_\epsilon}$, and the final claim follows from $Z_{f_\epsilon}^c \subset \{x \in \mathbb{R}^n : \text{dist}(x, Z_f^c) > \epsilon\}^c$. ■

Def: The norm on $C^k(G)$ is

$$\|f\|_{C^k(G)} = \sup_{x \in G, |\alpha| \leq k} |D^\alpha f(x)|$$

The norm on $L^p(G)$ is

$$\|f\|_{L^p(G)} = \left[\int_G |f(x)|^p dx \right]^{\frac{1}{p}} \quad (1 \leq p < \infty)$$

$$\|f\|_{L^\infty(G)} = \text{ess sup}_{x \in G} |f(x)| = \inf_{x \in G} \{B : |f(x)| \leq B, \text{ a.e. } x\} \quad (p = \infty)$$

Distributions

Lemma: If $f \in C_0(G)$ then $\|f_\epsilon - f\|_{C(G)} \rightarrow 0$. If $f \in L^p(G)$, $1 \leq p < \infty$, then $\|f_\epsilon\|_{L^p(G)} \leq \|f\|_{L^p(G)}$ and $\|f_\epsilon - f\|_{L^p(G)} \rightarrow 0$.

Proof: By uniform continuity of f on compact sets,

$$\begin{aligned} |f_\epsilon(x) - f(x)| &\leq \int_{\mathbb{R}^n} |f(x-y) - f(x)| \varphi_\epsilon(y) dy \\ &\leq \sup\{|f(x-y) - f(x)| : x \in \underline{f}, |y| \leq \epsilon\} \end{aligned}$$

Next, for $p = 1$, it follows with Fubini (and $f \rightarrow 0$ in $\mathbb{R}^n \setminus G$)

$$\|f_\epsilon\|_{L^1(G)} \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f(x-y)| \varphi_\epsilon(y) dy dx = \underbrace{\int_{\mathbb{R}^n} \varphi_\epsilon(y) dy}_{=1} \underbrace{\int_{\mathbb{R}^n} |f(x-y)| dx}_{=\|f\|_{L^1(G)}} dy$$

For $p = 2$, since $f_\epsilon \in L^2(G)$ (why?), let $\{\phi_n\} \subset C_0(G)$ be chosen so that $\|f_\epsilon - \phi_n\|_{L^2(G)} \xrightarrow{n \rightarrow \infty} 0$. As above,

$$\left| \int_G f_\epsilon \phi_n \right| \leq \int_{\mathbb{R}^n} \varphi_\epsilon(y) \left[\int_{\mathbb{R}^n} |f(x-y) \phi_n(x)| dx \right] dy \leq \|f\|_{L^2(G)} \|\phi_n\|_{L^2(G)}.$$

Distributions

Since $\phi_n \xrightarrow{L^2(G)} f_\epsilon$ and $\|\phi_n\|_{L^2(G)} \rightarrow \|f_\epsilon\|_{L^2(G)}$, it follows that $\|f_\epsilon\|_{L^2(G)} \leq \|f\|_{L^2(G)}$.

Study Question: Show that $\|f_\epsilon\|_{L^p(G)} \leq \|f\|_{L^p(G)}$ holds for the cases other than $p = 1, 2$.

Finally, for an arbitrary $\delta > 0$, let $\varphi \in C_0(G)$ be chosen so that $\|f - \varphi\|_{L^2(G)} \leq \delta/3$. Using the first result, let $\epsilon > 0$ be chosen small enough that $\|\varphi_\epsilon - \varphi\|_{C(G)} |\varphi_\epsilon|^{1/p} \leq \delta/3$. Then

$$\|f_\epsilon - f\|_{L^p(G)} \leq \|f_\epsilon - \varphi_\epsilon\|_{L^p(G)} + \|\varphi_\epsilon - \varphi\|_{L^p(G)} + \|\varphi - f\|_{L^p(G)} \leq \delta \quad \blacksquare$$

Theorem: $C_0^\infty(G)$ is dense in $L^p(G)$ for $1 \leq p < \infty$.

Theorem: $\forall K \subset\subset G, \exists \varphi \in C_0^\infty(G) \ni 0 \leq \varphi(x) \leq 1, x \in G$, and $\varphi(x) = 1, \forall x \in K$.

Proof: Set $\epsilon = \text{dist}(K, \partial G)/4$. Then set $f(x) = 1$ if $\text{dist}(x, K) \leq 2\epsilon$ and otherwise $f(x) = 0$. Then

$f_\epsilon \subset \{x : \text{dist}(x, K) \leq 3\epsilon\}$ and $f_\epsilon = 1$ on $\{x : \text{dist}(x, K) \leq \epsilon\}$. \blacksquare

Distributions

Def: A functional $T \in C_0^\infty(G)^*$ is a *distribution* on G , and this linear space of distributions is also denoted by $\mathcal{D}^*(G)$.

Example: Identify $L_{\text{loc}}^1(G) = \cap\{L^1(K) : K \subset\subset G\} \leq \mathcal{D}^*(G)$ through $T_f(\varphi) = \int_G f\bar{\varphi}$, $\varphi \in C_0^\infty(G)$.

Def: The α th partial derivative of the distribution T is the distribution $\partial^\alpha T$ defined according to $\partial^\alpha T(\varphi) = (-1)^{|\alpha|} T(D^\alpha \varphi)$, $\varphi \in C_0^\infty(G)$.

Note: $\partial^\alpha T_f = T_{D^\alpha f}$ for $|\alpha| \leq m$ and $f \in C^m(G)$.

Examples: A function $f \in L_{\text{loc}}^1(\mathbb{R})$ may be identified with the distribution T_f . In particular, the Heaviside function, $H(x) = (1 + \text{sign}(x))/2$, and $r(x) = xH(x)$ satisfy

$$\partial r(\varphi) = H(\varphi), \quad \partial H(\varphi) = \delta(\varphi) = \bar{\varphi}(0)$$

where δ is the Dirac functional. Similarly, $\partial^m \delta(\varphi) = (-1)^m D^m \bar{\varphi}(0)$.

Distributions

Suppose $f \in C^\infty(\mathbb{R} \setminus \{x_0\})$ has one-sided limits at $\{x_0\}$ so that the jump $\sigma_0(f)$ in the direction of increasing x is well-defined.

Then $\forall \varphi \in C_0^\infty(\mathbb{R})$,

$$\partial T_f(\varphi) = -T_f(\varphi') = -\int_{\mathbb{R}} f \overline{\varphi}' = \int_{\mathbb{R} \setminus \{x_0\}} f' \overline{\varphi} + \sigma_0(f) \delta_{x_0}(\varphi)$$

Def: The support of a distribution $T \in \mathcal{D}^*(G)$ is

$$\underline{T} = \overline{G} \setminus Z_T, \quad Z_T = \cup \{\text{open } S \subset G : T(\varphi) = 0, \forall \varphi \in C_0^\infty(S)\}$$

Example: The support of the Dirac δ functional, $\delta(\varphi) = \varphi(0)$, $\varphi \in C_0^\infty(\mathbb{R}^n)$, is $\underline{\delta} = \{0\}$.

Def: A distribution $T \in \mathcal{D}^*(G)$ is *constant* if $\exists c \in \mathbb{K} \ni T(\varphi) = c \int_G \overline{\varphi}$, $\varphi \in C_0^\infty(G)$, and T may be identified with c .

Theorem: $S \in \mathcal{D}^*(\mathbb{R}) \Rightarrow \exists T \in \mathcal{D}^*(\mathbb{R}) \ni S = \partial T$. Also, $T \in \mathcal{D}^*(\mathbb{R})$ and $\partial T = 0 \Rightarrow T$ is constant.

Distributions

Proof: $S = \partial T$ holds precisely when

$$T(\psi') = -S(\psi), \quad \forall \psi \in C_0^\infty(\mathbb{R}).$$

or equivalently when

$$T(\zeta) = -S(\psi), \quad \psi(x) = \int_{-\infty}^x \zeta$$

So define T at first in this way on the subspace of $C_0^\infty(\mathbb{R})$,

$$H = \{\zeta \in C_0^\infty(\mathbb{R}) : \int \zeta = 0\}.$$

Then take $T(\psi) = 0$ when $\int \psi \neq 0$. To show that this T is well-defined on all of $C_0^\infty(\mathbb{R})$, it will be shown that every

$\psi \in C_0^\infty(\mathbb{R})$ can be written uniquely as $\psi = \zeta + c\psi_0$ where $c \in \mathbb{K}$ and $\psi_0 \in C_0^\infty(\mathbb{R})$ is fixed arbitrarily with $\int \psi_0 = 1$. For existence, choose $\psi \in C_0^\infty(\mathbb{R})$, define $c = \int \psi$ and set

$\zeta = \psi - c\psi_0 \in H$. For uniqueness, suppose

$\zeta_1 + c_1\psi_0 = \zeta_2 + c_2\psi_0$ for $\zeta_i \in H$, $c_i \in \mathbb{K}$, $i = 1, 2$. Then

$0 = \int(\zeta_2 - \zeta_1) = (c_1 - c_2) \int \psi_0 = (c_1 - c_2)$, so

$\zeta_2 - \zeta_1 = (c_1 - c_2)\psi_0 = 0$.

Then $\partial T = S = 0$ holds precisely when

$$T(\zeta) = 0, \quad \zeta \in H$$

Distributions

or when T is the constant $T(\psi_0)$ because of

$$T(\psi) = T(\zeta + c\psi_0) = \bar{c}T(\psi_0) = T(\psi_0) \int \bar{\psi}, \quad \psi \in C_0^\infty(\mathbb{R}) \quad \blacksquare$$

Theorem: If f is absolutely continuous, then $g = Df \in L_{\text{loc}}^1(\mathbb{R})$ satisfies $\partial T_f = T_g$ in $\mathcal{D}'(\mathbb{R})$. Conversely, if $T \in \mathcal{D}'(\mathbb{R})$ with $\partial T = T_g$ for $g \in L_{\text{loc}}^1(\mathbb{R})$, then $T = T_f$ for an absolutely continuous f and $T_g = \partial T_f$.

Proof: If f is absolutely continuous, then Df exists in a.e. x , $Df \in L_{\text{loc}}^1(\mathbb{R})$ and $f(x) = f(0) + \int_0^x Df$. Integration by parts shows

$$\partial T_f(\varphi) = - \int f D\bar{\varphi} = \int Df \bar{\varphi} = T_{Df}(\varphi), \quad \varphi \in C_0^\infty(\mathbb{R})$$

Conversely, suppose $T \in \mathcal{D}'(\mathbb{R})$ with $\partial T = T_g$ for $g \in L_{\text{loc}}^1(\mathbb{R})$. Then define the absolutely continuous $h(x) = \int_0^x g$. According to the first part, $\partial T_h = T_g = \partial T$, and hence $T - T_h$ is constant, say $c \in \mathbb{K}$. Then setting $f = h + c$ gives $T = T_f$ and $T_g = \partial T = \partial T_f$. \blacksquare

Distributions

Examples (Distributions in \mathbb{R}^n): Let S be an $(n - 1)$ -dimensional C^1 manifold in \mathbb{R}^n . Suppose $f \in C^\infty(\mathbb{R}^n \setminus S)$ has one-sided limits at S so that the jumps $\sigma_i(f)$ in the direction of increasing x_i are well-defined. Then $\forall \varphi \in C_0^\infty(\mathbb{R}^n)$, $1 \leq i \neq n$,
$$\partial_i T_f(\varphi) = -T_f(D_i \varphi) = -\int_{\mathbb{R}^n} f D_i \bar{\varphi} = \int_{\mathbb{R}^n \setminus S} D_i f \bar{\varphi} + \int_S \sigma_i(f) \bar{\varphi} \nu_i dS$$
where $\nu = \{\nu_i\}$ is the unit normal at S .

Suppose for $G \subset \mathbb{R}^n$ that ∂G is an $(n - 1)$ -dimensional C^1 manifold. Let $f \in C^\infty(\bar{G})$ be given and extend f by zero outside \bar{G} . Then define the distribution L_f by

$$L_f(\varphi) = \int_{\mathbb{R}^n} f \Delta \bar{\varphi} = \int_G \bar{\varphi} \Delta f + \int_{\partial G} \left[\frac{\partial \bar{\varphi}}{\partial \nu} f - \frac{\partial f}{\partial \nu} \bar{\varphi} \right] dS$$

so

$$L_f(\varphi) - T_{\Delta f}(\varphi) = \int_{\partial G} \left[\frac{\partial \bar{\varphi}}{\partial \nu} f - \frac{\partial f}{\partial \nu} \bar{\varphi} \right] dS$$

Similarly, define D_f and N_f by

$$D_f(\varphi) = \int_{\partial G} f \frac{\partial \bar{\varphi}}{\partial \nu} dS = \int_G [f \Delta \bar{\varphi} + \nabla f \cdot \nabla \bar{\varphi}]$$

and

$$N_f(\varphi) = \int_{\partial G} \bar{\varphi} \frac{\partial f}{\partial \nu} dS = \int_G [\bar{\varphi} \Delta f + \nabla f \cdot \nabla \bar{\varphi}]$$

so that $L_f - T_{\Delta f} = D_f - N_f$.

Sobolev Spaces

Def (Sobolev Spaces): For $G \subset \mathbb{R}^n$ define the scalar product,

$$(f, g)_{H^m(G)} = \sum_{|\alpha| \leq m} \int_G D^\alpha f D^\alpha \bar{g}, \quad f, g \in C^m(\bar{G})$$

with corresponding norm $\|f\|_{H^m(G)} = (f, f)_{H^m(G)}^{1/2}$. Then define $H^m(G)$ as the Hilbert space given by the completion of $C^\infty(\bar{G})$ with respect to the norm $\|\cdot\|_{H^m(G)}$. Also, define $H_0^m(G)$ as the Hilbert space given by the completion of $C_0^\infty(G)$ with respect to the norm $\|\cdot\|_{H^m(G)}$.

Note: Through the identifications $C^m(\bar{G}) \subset H^m(G)$ or $C_0^m(G) \subset H_0^m(G)$, a smooth function f will henceforth be understood also as the Cauchy sequence (f, f, \dots) in the corresponding Sobolev space.

Note: According to $C^\infty(\bar{G}) \subset C^m(\bar{G}) \subset H^m(G)$ and $C_0^\infty(G) \subset C_0^m(G) \subset H_0^m(G)$, $H^m(G)$ and $H_0^m(G)$ are also the completions of $C^m(\bar{G})$ and $C_0^m(G)$, respectively, with respect to the norm $\|\cdot\|_{H^m(G)}$.

Sobolev Spaces

Note: $L^2(G)$ is the completion of $C_0(G)$ with respect to the norm $\|\cdot\|_{L^2(G)} = \|\cdot\|_{H^0(G)}$. Since $C_0(G) \subset C(\overline{G}) \subset L^2(G)$, it follows that $H^0(G) = L^2(G) = H_0^0(G)$. For $m \geq 1$ it is generally the case that $H^m(G) \neq H_0^m(G)$.

Def: The α th distributional derivative of $f \in H^m(G)$ is given by the unique function $D^\alpha f \in L^2(G)$ satisfying

$$(D^\alpha f, \varphi) = (-1)^{|\alpha|} (f, D^\alpha \varphi), \quad \forall \varphi \in C_0^\infty(G)$$

Theorem: Let $G \subset \mathbb{R}^n$ and $m \geq 0$. Then $f \in H^m(G) \Leftrightarrow \exists \{f_n\} \subset C^m(\overline{G}) \ni \forall \alpha, |\alpha| \leq m, \{D^\alpha f\}$ is Cauchy in $L^2(G)$ and $\|D^\alpha f - D^\alpha f_n\|_{L^2(G)} \rightarrow 0$.

Proof: (\Leftarrow) is clear from the definition of completion. For (\Rightarrow), let $f \in H^m(G)$. Then $\exists \{f_n\} \subset C^m(\overline{G}) \ni \|f - f_n\|_{H^m(G)} \rightarrow 0$. Since $\forall \alpha, |\alpha| \leq m, \|D^\alpha f_n - D^\alpha f_m\|_{L^2(G)} \leq \|f_n - f_m\|_{H^m(G)}$, $\{D^\alpha f_n\}$ is Cauchy in $L^2(G)$ with limit, say, g_α . Then $\forall \varphi \in C_0^\infty(G)$,

$$(g_\alpha, \varphi) \leftarrow (D^\alpha f_n, \varphi) = (-1)^{|\alpha|} (f_n, D^\alpha \varphi) \rightarrow (-1)^{|\alpha|} (f, D^\alpha \varphi)$$

Sobolev Spaces

where $(\cdot, \cdot) = (\cdot, \cdot)_{L^2(G)}$. Hence, g_α is the α th distributional derivative of f and thus


$$\|g_\alpha - D^\alpha f_n\|_{L^2(G)} = \|D^\alpha f - D^\alpha f_n\|_{L^2(G)} \rightarrow 0. \quad \blacksquare$$

Corollary: $m \geq k \geq 0 \Rightarrow H^m(G) \subset H^k(G) \subset L^2(G)$. Also, $f \in H^m(G) \Rightarrow D^\alpha f \in L^2(G), \forall \alpha, |\alpha| \leq m$. ((\Leftarrow) shown later!)

Theorem: The dual space $H_0^m(G)'$ is (identified with) the linear span $\langle \{\partial^\alpha T_f : |\alpha| \leq m, T \in \mathcal{D}^*(G), f \in L^2(G)\} \rangle$.

Proof: If $f \in L^2(G)$ and $|\alpha| \leq m$, then

$|\partial^\alpha T_f(\varphi)| = |(f, D^\alpha \varphi)_{L^2(G)}| \leq \|f\|_{L^2(G)} \|\varphi\|_{H^m(G)}, \forall \varphi \in C_0^\infty(G)$, so $\partial^\alpha T_f$ has a continuous extension to $H_0^m(G)$. Thus, the linear span of such extensions lies in $H_0^m(G)'$.

Conversely, if $T \in H_0^m(G)'$, then by Theorem , $\exists f \in H_0^m(G) \ni T(\varphi) = (f, \varphi)_{H^m(G)}, \forall \varphi \in C_0^\infty(G)$

Sobolev Spaces

Then setting $g_\alpha = D^\alpha f \in L^2(G)$,

$$T(\varphi) = \sum_{|\alpha| \leq m} T_{g_\alpha}(D^\alpha \varphi) = \sum_{|\alpha| \leq m} (-1)^{|\alpha|} \partial^\alpha T_{g_\alpha}(\varphi)$$

shows that T lies in the above linear span. ■

Study Question: Let $G \subset \mathbb{R}^n$ be bounded. Show $\forall \mathcal{F} \in H_0^m(G)'$, $\exists u \in H_0^m(G) \ni \mathcal{F}(v) = (\nabla^m u, \nabla^m v)_{L^2(G)}$, $v \in H_0^m(G)$.

(This functional may be extended naturally for $v \in H^m(G)$.)

Show $\forall \mathcal{G} \in H^m(G)'$, $\exists w \in H^m(G) \ni \mathcal{G}(v) = (w, v)_{H^m(G)}$, $v \in H^m(G)$.

(This functional may be restricted naturally to $v \in H_0^m(G)$.)

(Hint: Show that $(\nabla^m u, \nabla^m v)_{L^2(G)}$ is a norm on $H_0^m(G)$ and otherwise use Theorem [▶](#).)

Theorem: $H_0^m(\mathbb{R}^n) = H^m(\mathbb{R}^n)$

Proof: Since $C_0^m(\mathbb{R}^n) \subset C^m(\mathbb{R}^n)$, it follows that $H_0^m(\mathbb{R}^n) \subset H^m(\mathbb{R}^n)$. For the other direction, let $u \in H^m(\mathbb{R}^n)$ be arbitrary.

Fix the cut-off function $\tau \in C_0^\infty(B(0, 2))$ given by

$\tau = \psi_\epsilon \star \chi_{B(0, \frac{3}{2})}$, $\epsilon = \frac{1}{2}$, satisfying $\tau = 1$ on $B(0, 1)$ and

$$|D^\alpha \tau(x)| \leq M, \forall x \in \mathbb{R}^n, |\alpha| \leq m$$

Sobolev Spaces

For $k = 1, 2, \dots$ define $\tau_k(x) = \tau(x/k)$ which satisfies $\tau_k = 1$ on $B(0, k)$ and $\tau_k = 0$ outside $B(0, 2k)$ while

$$|D^\alpha \tau_k(x)| \leq M k^{-|\alpha|} \leq M, \forall x \in \mathbb{R}^n, |\alpha| \leq m.$$

Then $u^k = \tau_k u$ satisfies


$$|D^\alpha u^k(x)| \leq M \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} |D^\beta u(x)|$$

and $\exists B > 0 \ni \forall G \subset \mathbb{R}^n$,

$$\|u^k\|_{H^m(G)} \leq B \|u\|_{H^m(G)}$$

Since $u = u^k$ on $B(0, k)$,

$$\|u - u^k\|_{H^m(\mathbb{R}^n)} \leq (1 + B) \|u\|_{H^m(\mathbb{R}^n \setminus B(0, k))} \xrightarrow{k \rightarrow \infty} 0.$$

Next, define $u_\epsilon^k = \psi_\epsilon \star u^k$ for $\epsilon \in (0, 1)$. Since $\underline{u}^k \subset \overline{B}(0, 2k)$, $\underline{u}_\epsilon^k \subset B(0, 2k + 1) = G_k$. From the above Lemma ,

$\|u^k - u_\epsilon^k\|_{L^2(G_k)} \xrightarrow{\epsilon \rightarrow 0} 0$, and since $D^\alpha u_\epsilon^k = (D^\alpha u^k)_\epsilon$,

$\|D^\alpha u^k - D^\alpha u_\epsilon^k\|_{L^2(G_k)} \xrightarrow{\epsilon \rightarrow 0} 0$. Hence, $\|u^k - u_\epsilon^k\|_{H^m(\mathbb{R}^n)} =$

$\|u^k - u_\epsilon^k\|_{H^m(G_k)} \xrightarrow{\epsilon \rightarrow 0} 0$. Since the right side in

$$\|u - u_\epsilon^k\|_{H^m(\mathbb{R}^n)} \leq \|u - u^k\|_{H^m(\mathbb{R}^n)} + \|u^k - u_\epsilon^k\|_{H^m(\mathbb{R}^n)}$$

Sobolev Spaces

can be made arbitrarily small for first k large enough and then ϵ small enough, $C_0^\infty(\mathbb{R}^n)$ is dense in $H^m(\mathbb{R})$. ■

Theorem: Suppose for $G \subset \mathbb{R}^n$ (open) that $\sup\{|x_1| : x \in G\} = K < \infty$. Then

$$\|\varphi\|_{L^2(G)} \leq 2K \|D_1\varphi\|_{L^2(G)}, \forall \varphi \in H_0^1(G).$$

Proof: On the basis of density in $H_0^1(G)$, assume $\varphi \in C_0^\infty(G)$. Then integrate

$$D_1(x_1|\varphi(x)|^2) = |\varphi(x)|^2 + x_1 D_1(|\varphi(x)|^2)$$

over G to obtain

$$\begin{aligned} \int_G |\varphi(x)|^2 &= \int_G D \cdot \langle x_1 |\varphi(x)|^2, 0, \dots, 0 \rangle dx - \int_G x_1 D_1 |\varphi(x)|^2 dx \\ &= \int_{\partial G} \nu \cdot \langle x_1 |\varphi(x)|^2, 0, \dots, 0 \rangle dS - \int_G x_1 D_1 |\varphi(x)|^2 dx \\ &= - \int_G x_1 [D_1\varphi(x)\bar{\varphi}(x) + \varphi(x)D_1\bar{\varphi}(x)] dx \\ &\leq 2K \|D_1\varphi\|_{L^2(G)} \|\varphi\|_{L^2(G)} \end{aligned}$$

using the divergence theorem. ■

Sobolev Spaces

Def: For $G \subset \mathbb{R}^n$ open and bounded, ∂G a C^m manifold of dimension $n - 1$ when it can be represented locally as the graph of a C^m function. Specifically, there exists the following construction:

- The hypercube $Q = B_\infty(0, 1)$, the dividing hyperplane $Q_0 = Q \cap \{y : y_n = 0\}$ and the (upper) half-hypercube $Q_+ = Q \cap \{y : y_n > 0\}$.
- The open covering $\{G_j\}_{j=1}^N$ of ∂G , i.e., $\partial G \subset \cup_{j=1}^N G_j$.
- The functions $\varphi_j \in C^m(Q, G_j)$, each a bijection of Q , Q_+ and Q_0 onto G_j , $G_j \cap G$ and $G_j \cap \partial G$ with $J(\varphi_j) = \det(\partial\varphi_j/\partial x) > 0$.
- The pair (φ_j, G_j) is called a *coordinate patch*.

Def: With $G_0 = \bar{G}$, a *partition-of-unity* subordinate to the open cover $\{G_j\}_{j=0}^N$ of \bar{G} is a collection of functions $\{\beta_j\}_{j=0}^N$ satisfying $\beta_j \in C_0^\infty(G_j)$, $\beta_j(x) \geq 0$ and $\sum_{j=0}^N \beta_j(x) = 1$, $\forall x \in \bar{G}$.

Sobolev Spaces

Also, $\{\beta_j\}_{j=1}^N$ is a *partition-of-unity* subordinate to the open cover $\{G_j\}_{j=1}^N$ of ∂G . These partitions may be constructed as follows.

- Let $\{F_j\}_{j=1}^N$ be an open covering, $\partial G \subset \cup_{j=1}^N F_j$, with $\bar{F}_j \subset G_j$. Also, choose F_0 with $\bar{F}_0 \subset G_0$ and $\bar{G} \subset \cup_{j=0}^N F_j = F$.
- For $j = 0, \dots, N$ construct $\alpha_j \in C_0^\infty(G_j)$ with $\alpha_j = 1$ in \bar{F}_j and $\alpha \in C_0^\infty(F)$ with $\alpha = 1$ in \bar{G} , where $0 \leq \alpha_j(x), \alpha(x) \leq 1, \forall x \in \mathbb{R}^n$.
- For $j = 0, \dots, N$ define $\beta_j = \alpha \alpha_j / \sum_{k=0}^N \alpha_k$ in F and $\beta_j = 0$ in $\mathbb{R}^n \setminus F$.
- $\sum_{k=0}^N \alpha_k(x) > 0, \forall x \in F$ and $\alpha = 0, \forall x \in \partial F \Rightarrow \beta_j \in C_0^\infty(F) \subset C_0^\infty(\mathbb{R}^n)$.
- $\alpha, \alpha_j \geq 0 \Rightarrow \beta_j \geq 0$ and $\alpha_j \in C_0^\infty(G_j) \Rightarrow \beta_j \in C_0^\infty(G_j)$.
 $\alpha = 1$ in $\bar{G} \Rightarrow \sum_j \beta_j = 1$ in \bar{G} .

Sobolev Spaces

Localization on subdomains:

- For $u \in H^m(G)$, $u = \sum_{j=0}^N \beta_j u$ on G .
- $u_j = \beta_j u$ satisfies $\underline{u}_j \subset G_j$ and $u_j \in H_{\Gamma_j}^m(G \cap G_j)$, where $\Gamma_j = \partial G_j \cap \overline{G}$ and $H_{\Gamma_j}^m(G \cap G_j)$ is the completion of $C_0^m(G_j)$ with respect to the norm $\|\cdot\|_{H^m(G \cap G_j)}$. (Details?)
- $u \mapsto (u_0, \dots, u_N)$ maps $H^m(G)$ to $\times_{j=0}^N H_{\Gamma_j}^m(G \cap G_j)$, and it is linear and injective since $\sum_{j=0}^N \beta_j = 1$.
- $v_j = u_j \circ \varphi_j$, $1 \leq j \leq N$, satisfies $\underline{v}_j \subset Q_+$, $v_j \in H_{\Gamma}^m(Q_+)$, where $\Gamma = \partial Q \cap \overline{Q}_+$ and $H_{\Gamma}^m(Q_+)$ is the completion of $C_0^m(Q)$ with respect to the norm $\|\cdot\|_{H^m(Q_+)}$.
- $\Lambda : u \mapsto (u_0, v_1, \dots, v_N)$ maps $H^m(G)$ to $H_0^m(G) \times [H_{\Gamma}^m(Q_+)]^N$. It is a continuous linear injection mapping onto a closed subspace, its range, where it has a continuous inverse. (Details?)

Sobolev Spaces

Localization on the boundary:

- $C^m(\partial G)$ is the set of functions $f : \partial G \rightarrow \mathbb{R}$ where $(\beta_j f) \circ \varphi_j \in C^m(Q_0)$, $1 \leq j \leq N$.

- Integrals over ∂G are given by

$$\int_{\partial G} f dS = \sum_{j=1}^N \int_{Q_0} (\beta_j f) \circ \varphi_j(y) J_j(y) dy, \quad J_j = \left[\det \left[\frac{\partial \varphi_j}{\partial y} \right]^T \frac{\partial \varphi_j}{\partial y} \right]^{\frac{1}{2}}$$

- Define the scalar product and norm on $C(\partial G) = C^0(\partial G)$,

$$(f, g)_{L^2(\partial G)} = \int_{\partial G} f \bar{g} dS, \quad \|f\|_{L^2(\partial G)} = (f, f)_{L^2(\partial G)}^{\frac{1}{2}}$$

- Define $L^2(\partial G)$ as the completion of $C(\partial G)$ with respect to this scalar product.

- $\lambda : f \mapsto ((\beta_1 f) \circ \varphi_1, \dots, (\beta_N f) \circ \varphi_N)$ maps $L^2(\partial G)$ to $[L^2(Q_0)]^N$. It is a continuous linear injection mapping onto a closed subspace, its range, where it has a continuous inverse. (Details?)

Trace

- *Traces* are a generalization of boundary values.
- For instance, functions in $L^2(G)$ have no well-defined boundary values since $|\partial G| = 0$.
- First develop traces for $G = \mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n > 0\}$ where $\partial G = \{x \in \mathbb{R}^n : x_n = 0\}$.
- Later the general case will build upon this simpler case by using localization as above.

Theorem: For $G = \mathbb{R}_+^n$ the trace mapping

$\gamma_0 : C^1(\overline{G}) \rightarrow C^0(\partial G)$ defined by

$$\gamma_0(\varphi)(x') = \varphi(x', 0), \quad \varphi \in C^1(\overline{G}), \quad x' \in \partial G$$

has a unique extension to an operator $\gamma_0 \in \mathcal{L}(H^1(G), L^2(\partial G))$ whose range is dense in $L^2(\partial G)$, and it satisfies

$$\gamma_0(\beta u) = \gamma_0(\beta)\gamma_0(u), \quad \beta \in C^1(\overline{G}), \quad u \in H^1(G).$$


Trace

Proof: For $\varphi \in C^1(\overline{G})$ and $x' \in \mathbb{R}^{n-1}$,

$$|\varphi(x', 0)|^2 = - \int_0^\infty D_n(|\varphi(x', x_n)|^2) dx_n.$$

Integrating over \mathbb{R}^{n-1} gives

$$\begin{aligned} \|\varphi(\cdot, 0)\|_{L^2(\mathbb{R}^{n-1})}^2 &\leq \int_{\mathbb{R}_+^n} |\overline{\varphi} D_n \varphi + \varphi D_n \overline{\varphi}| dx \leq 2 \|\varphi\|_{L^2(\mathbb{R}_+^n)} \|D_n \varphi\|_{L^2(\mathbb{R}_+^n)} \\ &\leq \|\varphi\|_{L^2(\mathbb{R}_+^n)}^2 + \|D_n \varphi\|_{L^2(\mathbb{R}_+^n)}^2 = \|\varphi\|_{H^1(\mathbb{R}_+^n)}^2. \end{aligned}$$

The existence of a unique continuous linear extension to $\mathcal{L}(H^1(G), L^2(\partial G))$ follows with Theorem .

For $\tau = \psi_\epsilon \star \chi_{B(0, \frac{3}{2})}$, $\epsilon = \frac{1}{2}$, and $\psi \in C_0^\infty(\mathbb{R}^{n-1})$,

$$\varphi(x) = \psi(x') \tau(x_n), \quad x = (x', x_n) \in \mathbb{R}_+^n$$

defines a $\varphi \in C^1(\overline{G})$ and $\gamma_0(\varphi) = \psi$. Thus, the range of γ_0 contains $C_0^\infty(\mathbb{R}^{n-1})$, which is dense in $L^2(\partial \mathbb{R}_+^n)$. For the last

claim, let $u_\epsilon \in C^1(\overline{G})$ satisfy $\|u - u_\epsilon\|_{H^1(G)} \xrightarrow{\epsilon \rightarrow 0} 0$, so that also


for $\beta \in C^1(\overline{G})$, $\|\beta(u - u_\epsilon)\|_{H^1(G)} \leq \|\beta\|_{C^1(G)} \|u - u_\epsilon\|_{H^1(G)} \xrightarrow{\epsilon \rightarrow 0} 0$.

Then note by the continuity of γ ,

$$\gamma_0(\beta u) \xleftarrow{\epsilon \rightarrow 0} \gamma_0(\beta u_\epsilon) = \gamma_0(\beta) \gamma_0(u_\epsilon) \xrightarrow{\epsilon \rightarrow 0} \gamma_0(\beta) \gamma_0(u). \quad \blacksquare$$

Trace

Theorem: Let $u \in H^1(\mathbb{R}_+^n)$. Then $u \in H_0^1(\mathbb{R}_+^n)$ iff $\gamma_0(u) = 0$.

Proof: If $\{u_n\} \subset C_0^\infty(\mathbb{R}_+^n)$ converges to $u \in H^1(\mathbb{R}_+^n)$, then $\gamma_0(u) = \lim \gamma_0(u_n) = 0$ by Theorem .

Let $u \in H^1(\mathbb{R}_+^n)$ with $\gamma_0 u = 0$. Recall the cut-off function $\tau \in C_0^\infty(B(0, 2))$ given by $\tau = \psi_\epsilon \star \chi_{B(0, \frac{3}{2})}$, $\epsilon = \frac{1}{2}$, satisfying $\tau = 1$ on $B(0, 1)$ and

$$|D^\alpha \tau(x/k)| \leq M k^{-|\alpha|} \leq M, \forall x \in \mathbb{R}^n, |\alpha| = 1.$$

Define the new cut-off function,

$$\begin{aligned} \sigma(t) &= t^2(3 - 2t), \quad t \in [0, 1], \\ \sigma(t) &= 0, \quad t < 0, \quad \sigma(t) = 1, \quad t > 1. \end{aligned}$$

satisfying

$$|D_{x_n} \sigma(kx_n - 1)| \leq \frac{3}{2}k < 2k.$$

Then set $\phi_k(x) = \tau(x/k)\sigma(kx_n - 1) = \tau_k(x)\sigma_k(x)$ and $u^k = \phi_k u$ so that $\underline{u}^k \subset \{x \in \mathbb{R}^n : |x| \leq 2k \text{ \& } x_n \geq 1/k\}$ while $u^k = u$ in $\{x \in \mathbb{R}^n : |x| \leq k \text{ \& } x_n \geq 2/k\} = E_k$. Since $\phi_k(x) \in [0, 1]$, $\forall x$, it follows that $|u^k| \leq |u|$ and

Trace

$$\|u - u^k\|_{L^2(\mathbb{R}_+^n)} \leq 2\|u\|_{L^2(\mathbb{R}_+^n \setminus E_k)} \xrightarrow{k \rightarrow \infty} 0.$$

For $i \neq n$, $|D_{x_i} u^k| = |\sigma_k[uD_{x_i}\tau_k + \tau_k D_{x_i}u]| \leq M|u| + |D_{x_i}u|$ so

$$\|D_{x_i}(u - u^k)\|_{L^2(\mathbb{R}_+^n)} \leq (2 + M)\|u\|_{H^1(\mathbb{R}_+^n \setminus E_k)} \xrightarrow{k \rightarrow \infty} 0$$


Then $D_{x_n} u^k = \sigma_k[uD_{x_n}\tau_k + \tau_k D_{x_n}u] + \tau_k u D_{x_n} \sigma_k$, so with the estimates $|\sigma_k[uD_{x_n}\tau_k + \tau_k D_{x_n}u]| \leq M|u| + |D_{x_n}u|$ and $|\tau_k u D_{x_n} \sigma_k| \leq 2k|u|$, the n th derivatives satisfy

$$\|D_{x_n}(u - u^k)\|_{L^2(\mathbb{R}_+^n)} \leq (2 + M)\|u\|_{H^1(\mathbb{R}_+^n \setminus E_k)} + 2k\|u\|_{L^2(F_k)}$$

where $F_k = \{x \in \mathbb{R}^n : 0 \leq x_n \leq \frac{2}{k}\}$ contains $\{x \in \mathbb{R}^n : \frac{1}{k} \leq x_n \leq \frac{2}{k}\}$ in which $D_{x_n} \sigma_k \neq 0$.

Study Question: Show that $u(x', x_n) = \int_0^{x_n} D_{x_n} u(x', t) dt$, a.e. $x' \in \mathbb{R}^{n-1}$, $x_n \in \mathbb{R}_+^1$.

Answer: Choose $\{\phi_\epsilon\} \subset C^\infty(\overline{R_+^n})$ so that $\|u - \phi_\epsilon\|_{H^1(\mathbb{R}_+^n)} \xrightarrow{\epsilon \rightarrow 0} 0$.

Then with Theorem  $|\phi_\epsilon(\cdot, 0)| \leq |\gamma_0(u)| + |\gamma_0(u - \phi_\epsilon)| \leq$

$c\|u - \phi_\epsilon\|_{H^1(\mathbb{R}_+^n)} \xrightarrow{\epsilon \rightarrow 0} 0$. Since convergence in $L^2(\mathbb{R}_+^n)$ gives a.e. pointwise convergence,

Trace

$$u(x', x_n) \xrightarrow{\epsilon \rightarrow 0} \phi_\epsilon(x', x_n) - \phi_\epsilon(x', 0) = \int_0^{x_n} D_{x_n} \phi_\epsilon(x', t) dt \\ \xrightarrow{\epsilon \rightarrow 0} \int_0^{x_n} D_{x_n} u(x', t) dt, \text{ a.e. } x' \in \mathbb{R}^{n-1}, x_n \in \mathbb{R}_+^1.$$

Then with Cauchy-Schwarz,

$$|u(x', x_n)|^2 \leq [\int_0^{x_n} 1^2 dt] [\int_0^{x_n} |D_{x_n} u(x', t)|^2 dt] = x_n \int_0^{x_n} |D_{x_n} u(x', t)|^2 dt$$

$$\text{So } \int_0^{2/k} |u(x', x_n)|^2 dx_n \leq \int_0^{2/k} x_n [\int_0^{x_n} |D_{x_n} u(x', t)|^2 dt] dx_n \\ \leq \frac{2}{k} \int_0^{2/k} [\int_0^{x_n} |D_{x_n} u(x', t)|^2 dt] dx_n = \\ \frac{2}{k} \int_0^{2/k} \int_t^{2/k} |D_{x_n} u(x', t)|^2 dx_n dt = \frac{4}{k^2} \int_0^{2/k} |D_{x_n} u(x', t)|^2 dt$$

and hence,

$$2k \|u\|_{L^2(F_k)} \leq 4 \|D_{x_n} u\|_{L^2(F_k)}$$

Combining the above estimates gives

$$\|D_{x_n}(u - u^k)\|_{L^2(\mathbb{R}_+^n)} \leq (2+M) \|u\|_{H^1(\mathbb{R}_+^n \setminus E_k)} + 8 \|D_{x_n} u\|_{L^2(F_k)} \xrightarrow{k \rightarrow \infty} 0$$


Thus, $\|u - u^k\|_{H^1(\mathbb{R}_+^n)} \xrightarrow{k \rightarrow \infty} 0$. Finally, define $u_\epsilon^k = \psi_\epsilon \star u^k$ for

$\epsilon \in (0, 1/k)$ so that $u_\epsilon^k \in C_0^\infty(\mathbb{R}_+^n)$. Since

$$\underline{u}^k \subset \{x \in \mathbb{R}^n : |x| \leq 2k \text{ \& } x_n \geq 1/k\},$$

$$\underline{u}_\epsilon^k \subset \{x \in \mathbb{R}^n : |x| < 2k + 1/k \text{ \& } x_n > 0\} = G_k. \text{ From the above}$$

Trace

Lemma , $\|u^k - u_\epsilon^k\|_{L^2(G_k)} \xrightarrow{\epsilon \rightarrow 0} 0$, and since $D^\alpha u_\epsilon^k = (D^\alpha u^k)_\epsilon$,

$\|D^\alpha u^k - D^\alpha u_\epsilon^k\|_{L^2(G_k)} \xrightarrow{\epsilon \rightarrow 0} 0$. Hence, $\|u^k - u_\epsilon^k\|_{H^1(\mathbb{R}^n)} =$

$\|u^k - u_\epsilon^k\|_{H^1(G_k)} \xrightarrow{\epsilon \rightarrow 0} 0$. Since the right side in

$$\|u - u_\epsilon^k\|_{H^1(\mathbb{R}^n)} \leq \|u - u^k\|_{H^1(\mathbb{R}^n)} + \|u^k - u_\epsilon^k\|_{H^1(\mathbb{R}^n)}$$

can be made arbitrarily small for first k large enough and then ϵ small enough, u can be approximated in $H^1(G)$ arbitrarily well with $C_0^\infty(\mathbb{R}^n)$ and hence $u \in H_0^1(G)$. ■

For ∂G sufficiently smooth, $\gamma_0 : H^1(G) \rightarrow L^2(\partial G)$ is defined as follows in terms of the natural formulation above for a curvature-free boundary:

$$\gamma_0(u) = \sum_{j=1}^N (\gamma_0((\beta_j u) \circ \varphi_j)) \circ \varphi_j^{-1}$$

where $\{\beta_j\}_{j=1}^N$ gives a partition-of-unity subordinate to the open cover $\{G_j\}_{j=1}^N$ of ∂G and $\{(\varphi_j, G_j)\}_{j=1}^N$ are corresponding coordinate patches.

Estimating γ_0 and extending by continuity gives the following.

Trace

Theorem: Let $G \subset \mathbb{R}^n$ be bounded and open with ∂G a C^1 manifold where G lies only on one side of ∂G . Then there is a unique $\gamma_0 \in \mathcal{L}(H^1(G), L^2(\partial G))$ such that $\gamma_0(u) = u|_{\partial G}$ for each $u \in C^1(\bar{G})$. Also, $K(\gamma_0) = H_0^1(G)$ and $\overline{\text{Rg}(\gamma_0)} = L^2(\partial G)$.

Higher order traces of normal derivatives are first defined in terms of usual boundary values for sufficiently smooth functions and then extended by continuity with the following result.

Theorem: Let $G \subset \mathbb{R}^n$ be bounded and open with ∂G a C^1 manifold where G lies only on one side of ∂G . Then there is a unique $\gamma \in \mathcal{L}(H^m(G), \prod_{j=0}^{m-1} H^{m-1-j}(\partial G))$ such that

$$\gamma(u) = (\gamma_0(u), \dots, \gamma_{m-1}(u))$$

and $\gamma_j(u) = \partial^j u / \partial \nu^j |_{\partial G}$, $0 \leq j \leq m-1$, for $u \in C^m(\bar{G})$.

Also, $K(\gamma) = H_0^m(G)$ and $\overline{\text{Rg}(\gamma)} = \prod_{j=0}^{m-1} H^{m-1-j}(\partial G)$.

Note that $\text{Rg}(\gamma)$ can be characterized in terms of fractional order Sobolev spaces, e.g., $\gamma_0 \in \mathcal{L}(H^1(G), H^{\frac{1}{2}}(\partial G))$. However, the presented results are sufficient in this work.

Sobolev's Lemma and Imbedding

Goal: Identify $C_u^k(G)$ with $H^m(G)$ for certain k and m , where

Def: $(C_u^k(G), \|\cdot\|_{C^k(G)})$ is the Banach space of functions with uniformly continuous derivatives up to order k . Note for $G \subset \mathbb{R}^n$ bounded, $C_u^k(G) = C^k(\overline{G})$.

For this goal, G must possess a certain regularity:

Def: Let a cone with vertex y be denoted by $K(y) = K(y; \rho, \Omega) = \{x = y + \lambda\omega, \lambda \in (0, \rho), \omega \in \Omega\}$ where $\Omega = \partial B(0, 1) \cap B(x, r)$ for some $x \in \partial B(0, 1)$ and $r > 0$. Then $|K(y)| = \gamma\rho^n$ where $\gamma = |\Omega|_{\partial B(0,1)}$ is the *solid angle* of Ω .

A domain G satisfies a *cone condition* if $\exists \rho, \gamma > 0 \ni \forall y \in \overline{G}$, $\exists K(y; \rho, \Omega) \subset \overline{G}$ with $\gamma = |\Omega|_{\partial B(0,1)}$.

Theorem: Suppose $G \subset \mathbb{R}^n$ is open and bounded and satisfies a cone condition. Then for $m > n/2$, $\exists C > 0 \ni$

$$\|u\|_{C_u(G)} = \|u\|_{C(G)} \leq C \|u\|_{H^m(G)}, \quad \forall u \in C^m(\overline{G})$$

Sobolev's Lemma and Imbedding

Proof: Fix $g \in C_0^\infty(\mathbb{R})$ by $g = \psi_\epsilon \star \chi_{[-\frac{3}{4}, \frac{3}{4}]}$, $\epsilon = \frac{1}{4}$. Define

$\tau(t) = g(t/\rho)$, $\rho > 0$, satisfying

$$|\tau^{(k)}(t)| \leq A_k \rho^{-k}$$

for some constants A_k . Let $u \in C^m(\overline{G})$ and assume $2m > n$.

For $y \in \overline{G}$, $K(y) \subset \overline{G}$, integrate along a ray

$\{x = y + r\omega, r \in (0, \rho)\} \subset K(y)$, $\omega \in \Omega$, emanating from y :

$$\int_0^\rho D_r[\tau(r)u(y + r\omega)]dr = -u(y)$$

So integrating over all of $K(y)$,

$$\int_\Omega \int_0^\rho D_r[\tau(r)u(y + r\omega)]drd\omega = -u(y) \int_\Omega d\omega = -u(y)\gamma$$

Integrate by parts $m - 1$ times to obtain

$$u(y) = \frac{(-1)^m}{\gamma(m-1)!} \int_\Omega \int_0^\rho D_r^m(\tau u) r^{m-1} drd\omega$$

Then with $x = y + r\omega$, $dx = r^{n-1} drd\omega$,

$$|u(y)| = \frac{1}{\gamma(m-1)!} \left| \int_{K(y)} D_r^m(\tau u) |x - y|^{m-n} dx \right|$$

With Cauchy-Schwarz,

$$|u(y)|^2 \leq \frac{1}{(\gamma(m-1)!)^2} \left[\int_{K(y)} |D_r^m(\tau u)|^2 dx \right] \left[\int_{K(y)} |x - y|^{2(m-n)} dx \right]$$

Sobolev's Lemma and Imbedding

Using

$$\int_{K(y)} |x - y|^{2(m-n)} dx = \int_{\Omega} \int_0^{\rho} r^{2m-n-1} dr d\omega = \frac{\gamma \rho^{2m-n}}{2m-n}$$

the previous estimate becomes

$$|u(y)|^2 \leq C_{m,n} \rho^{2m-n} \int_{K(y)} |D_r^m(\tau u)|^2 dx$$

where $C_{m,n}$ depends upon m and n . Then

$$|D_r^m(\tau u)| = \left| \sum_{k=0}^m \binom{m}{k} D_r^{m-k} \tau D_r^k u \right| \leq \sum_{k=0}^m \binom{m}{k} \frac{A_{m-k}}{\rho^{m-k}} |D_r^k u|$$

or

$$|D_r^m(\tau u)|^2 \leq C' \sum_{k=0}^m \rho^{2(k-m)} |D_r^k u|^2$$

and hence

$$|u(y)|^2 \leq C_{m,n} C' \sum_{k=0}^m \rho^{2k-n} \int_{K(y)} |D_r^k u|^2 dx$$

By the chain rule,

$$|D_r^k u|^2 \leq C'' \sum_{|\alpha| \leq k} |D^\alpha u(x)|^2$$

Then

$$\begin{aligned} \sup_{y \in G} |u(y)|^2 &\leq C \sup_{y \in G} \sum_{|\alpha| \leq m} \int_{K(y)} |D^\alpha u(x)|^2 dx \\ &\leq C \sum_{|\alpha| \leq m} \int_G |D^\alpha u(x)|^2 dx = C \|u\|_{H^m(G)}^2 \end{aligned}$$




Sobolev's Lemma and Imbedding

Def: An imbedding $i : H^m(G) \rightarrow C_u^k(G)$ is defined so that for $u \in H^m(G)$, the smooth function $i(u)$ (understood as identified with the Cauchy sequence $(i(u), i(u), \dots)$) satisfies $\|u - i(u)\|_{H^m(G)} = 0$. The continuity of i is represented by $H^m(G) \hookrightarrow C_u^k(G)$.

Theorem: Suppose $G \subset \mathbb{R}^n$ is open and bounded and satisfies a cone condition. Then for $m > k + n/2$ the imbedding $i : H^m(G) \rightarrow C_u^k(G)$ is continuous.

Proof: Applying Theorem  to $D^\alpha u$, $|\alpha| \leq k$, gives

$$\|u\|_{C_u^k(G)} = \|u\|_{C^k(G)} \leq C\|u\|_{H^m(G)}, \quad u \in C^m(\bar{G})$$

Thus, the imbedding is continuous from the dense subset $C^m(\bar{G})$ of $H^m(G)$ into the Banach space $C_u^k(G)$. The claim then follows from Theorem . ■

Study Question: For $G \subset \mathbb{R}^n$ and $x_0 \in G$, define $\delta_{x_0}(\varphi) = \bar{\varphi}(x_0)$, $\varphi \in C^\infty(\bar{G})$, and show that $\delta_{x_0} \in (H^m(G))'$ for $m > n/2$.

Density and Compactness

Def: $\mathcal{H}^m(G) = \{f \in L^2(G) : D^\alpha f \in L^2(G), |\alpha| \leq m\}$ is a Hilbert space equipped with $(\cdot, \cdot)_{H^m(G)}$ and $H^m(G) \leq \mathcal{H}^m(G)$.

Goal: Show $\mathcal{H}^m(G) \leq H^m(G)$.

Lemma: $C_0^\infty(\mathbb{R}^n)$ is dense in $\mathcal{H}^m(\mathbb{R}^n)$.

Proof: See the proof of Theorem . ■

Lemma: $\mathcal{H}^m(\mathbb{R}_+^n) \leq H^m(\mathbb{R}_+^n)$.

Proof: Fix $u \in \mathcal{H}^m(\mathbb{R}_+^n)$. Define $\sigma_k(t) = \sigma(kt)$ where $\sigma = \psi_\epsilon \star \chi_{[\frac{1}{2}, \infty)}$, $\epsilon = \frac{1}{2}$, and set $u^k(x', x_n - 1/k) = \sigma_k(x_n)u(x', x_n)$ for $x' \in \mathbb{R}^{n-1}$, $x_n \geq 0$, and otherwise $u^k = 0$. Then $\|u - u^k\|_{H^m(\mathbb{R}_+^n)} \xrightarrow{k \rightarrow \infty} 0$. (Why? LDCT) Also, $u^k \in \mathcal{H}^m(\mathbb{R}^n)$. By the above Lemma, $\exists \{u_\epsilon^k\}_{\epsilon > 0} \subset C_0^\infty(\mathbb{R}^n)$ such that $\|u^k - u_\epsilon^k\|_{H^m(\mathbb{R}^n)} \xrightarrow{\epsilon \rightarrow 0} 0$, and $\{u_\epsilon^k|_{\mathbb{R}_+^n}\}_{\epsilon > 0} \subset C^\infty(\overline{\mathbb{R}_+^n})$.

Density and Compactness

Since the right side in

$$\|u - u_\epsilon^k\|_{H^m(\mathbb{R}_+^n)} \leq \|u - u^k\|_{H^m(\mathbb{R}_+^n)} + \|u^k - u_\epsilon^k\|_{H^m(\mathbb{R}_+^n)}$$

can be made arbitrarily small for first k large enough and then ϵ small enough, u can be approximated in $\mathcal{H}^m(\mathbb{R}_+^n)$ arbitrarily well with $C^\infty(\mathbb{R}^n)$ and hence $u \in H^m(\mathbb{R}_+^n)$. ■

Lemma: $\exists \mathcal{P} \in \mathcal{L}(\mathcal{H}^m(\mathbb{R}_+^n), \mathcal{H}^m(\mathbb{R}^n)) \ni (\mathcal{P}u)(x) = u(x)$ for a.e. $x \in \mathbb{R}_+^n$.

Proof: By the last Lemma, it suffices to construct \mathcal{P} on $C^m(\overline{\mathbb{R}_+^n})$. Let $\{\lambda_j\} \subset \mathbb{R}^m$ solve the system,

$$\sum_{i=1}^m i^k \lambda_i = (-1)^k, \quad k = 0, \dots, m-1$$

For each $u \in C^m(\overline{\mathbb{R}_+^n})$, define

$$(\mathcal{P}u)(x) = \begin{cases} u(x), & x_n \geq 0 \\ \sum_{i=1}^m \lambda_i u(x', -ix_n), & x_n < 0 \end{cases}$$

By the construction for $\{\lambda_j\}$, $D_n^j(\mathcal{P}u)$ is continuous at $x_n = 0$ for $j = 0, \dots, m-1$. It follows that $\mathcal{P}u \in \mathcal{H}^m(\mathbb{R}^n)$. \mathcal{P} is clearly linear and continuous. ■

Density and Compactness

Theorem: Suppose $G \subset \mathbb{R}^n$ is open and bounded and lies on one side of ∂G which is a C^m manifold. Then

$$\exists \mathcal{P}_G \in \mathcal{L}(\mathcal{H}^m(G), \mathcal{H}^m(\mathbb{R}^n)) \ni \mathcal{P}_G u|_G = u, \forall u \in \mathcal{H}^m(G).$$

Proof: Let $\{(\varphi_k, G_k)\}_{k=1}^N$ be coordinate patches on ∂G and let $\{\beta_k\}_{k=0}^N$ be a partition-of-unity subordinate to G . Then $u \in \mathcal{H}^m(G) \Rightarrow u = \sum_{j=0}^N (\beta_j u)$. Since $\beta_0 u$ is smoothly and compactly supported, its extension by zero lies in $\mathcal{H}^m(\mathbb{R}^n)$. For the extension of $\beta_k u$, $k \geq 1$, note that $v \mapsto v \circ \varphi_k$ is an isomorphism from $\mathcal{H}^m(G_k \cap G)$ onto $\mathcal{H}^m(Q_+)$. (Why?) Since $(\beta_k u) \circ \varphi_k$ is smoothly supported in Q , it can be extended by zero in $\mathbb{R}_+^n \setminus Q$ to obtain an element of $\mathcal{H}^m(\mathbb{R}_+^n)$. By the previous Lemma, and the details of the proof, this can be extended to an element $\mathcal{P}((\beta_k u) \circ \varphi_k)$ of $\mathcal{H}^m(\mathbb{R}^n)$ with support in Q . The desired extension of $\beta_k u$ is given by $\mathcal{P}((\beta_k u) \circ \varphi_k) \circ \varphi_k^{-1}$ extended by zero outside G_k . The following is linear,

$$\mathcal{P}_G u = \beta_0 u + \sum_{k=1}^N \mathcal{P}((\beta_k u) \circ \varphi_k) \circ \varphi_k^{-1}$$

and satisfies $\|\mathcal{P}_G u\|_{\mathcal{H}^m(\mathbb{R}^n)} \leq c(\{\beta_k\}, \{\varphi_k\}) \|u\|_{\mathcal{H}^m(G)}$. ■

Density and Compactness

Theorem: Suppose $G \subset \mathbb{R}^n$ is open and bounded and lies on one side of ∂G which is a C^m manifold. Then $H^m(G) = \mathcal{H}^m(G)$.

Proof: Note that $H^m(G) \leq \mathcal{H}^m(G)$. To be shown is that $\mathcal{H}^m(G) \leq H^m(G)$. Let $u \in \mathcal{H}^m(G)$. Then $\mathcal{P}_G u \in \mathcal{H}^m(\mathbb{R}^n)$ and the density of $C_0^\infty(\mathbb{R}^n)$ in $\mathcal{H}^m(\mathbb{R}^n)$ gives a sequence $\{\varphi_k\} \subset C_0^\infty(\mathbb{R}^n)$ which converges to $\mathcal{P}_G u$. Thus, $\{\varphi_k|_G\}$ converges to u in $\mathcal{H}^m(G)$. ■

Lemma: Let Q be a cube in \mathbb{R}^n with edges of length $d > 0$. If $u \in C^1(\bar{Q})$ and $\bar{u} = \int_Q u / |Q|$, then

$$\|u\|_{L^2(Q)}^2 \leq d^{-n} |Q|^2 \bar{u}^2 + (nd^2/2) \sum_{i=1}^n \|D_i u\|_{L^2(Q)}^2$$

Proof: For $x, y \in Q$,

$$u(y) - u(x) = \sum_{i=1}^n \int_{x_i}^{y_i} D_i u(y_1, \dots, y_{i-1}, s, x_{i+1}, \dots, x_n) ds$$

Squaring and using Cauchy-Schwarz,

$$u^2(x) + u^2(y) - 2u(x)u(y) \leq$$

$$\left[\sum_{i=1}^n \int_{a_i}^{b_i} ds \right] \left[\sum_{i=1}^n \int_{a_i}^{b_i} (D_i u)^2(\dots, y_{i-1}, s, x_{i+1}, \dots) ds \right]$$

Density and Compactness

where $Q = \prod_{i=1}^n [a_i, b_i]$ and $b_k - a_k = d$, $1 \leq k \leq n$. Integrating the preceding w.r.t. $\{x_i\}_{i=1}^n$ and $\{y_i\}_{i=1}^n$ gives

$$2d^n \|u\|_{L^2(Q)}^2 \leq 2|Q|^2 \bar{u}^2 + nd^{n+2} \sum_{i=1}^n \|D_i u\|_{L^2(Q)}^2 \quad \blacksquare$$

Theorem: Suppose $G \subset \mathbb{R}^n$ is open and bounded. Then the imbedding $i : H_0^1(G) \rightarrow L^2(G)$ is compact.

Proof: Fix $\{u_k\} \subset C_0^\infty(G)$ with $M = \sup\{\|u_k\|_{H^1(G)}\} < \infty$. Let Q be a hypercube containing G , where the sides of Q have length $d \geq 1$. Extend each u_k by zero in $Q \setminus G$ so

$\{u_k\} \subset C_0^\infty(Q)$ with $M = \sup\{\|u_k\|_{H^1(Q)}\} < \infty$.

Let $\epsilon > 0$. Choose N so that $2nd^2 M^2 / N^2 < \epsilon$. Divide Q into congruent hypercubes Q_j , $j = 1, \dots, N^n$, with edges of length d/N . Since $\{u_k\}$ is bounded in $L^2(Q)$, it follows with Theorem 1 (polynomials being dense in $L^2(G)$) that there is a subsequence, again denoted for convenience by $\{u_k\}$, converging weakly in $L^2(Q)$. So $\exists K \ni$

$$\left| \int_{Q_j} (u_k - u_l) \right|^2 < \frac{\epsilon}{2N^{2n}}, \quad j = 1, \dots, N^n, \quad k, l \geq K$$

Density and Compactness

According to the previous Lemma, the following is obtained after summing over each Q_j ,

$$\|u_k - u_l\|_{L^2(Q)}^2 \leq N^n \left(\frac{d}{N}\right)^{-n} \frac{\epsilon}{2N^{2n}} + \frac{n}{2} \left(\frac{d}{N}\right)^2 4M^2 < \epsilon.$$

With $\|u_k - u_l\|_{L^2(G)}^2 \leq \|u_k - u_l\|_{L^2(Q)}^2$, $\{u_k\}$ is Cauchy in $L^2(G)$. ■

Corollary: Suppose $G \subset \mathbb{R}^n$ is open and bounded. Then the imbedding $i : H_0^m(G) \rightarrow H_0^{m-1}(G)$ is compact.

Theorem: Suppose $G \subset \mathbb{R}^n$ is open and bounded and lies on one side of ∂G which is a C^m manifold. Then the imbedding $i : H^m(G) \rightarrow H^{m-1}(G)$ is compact.

Proof: Fix $\{u_k\}$ bounded in $H^m(G)$. Then $\mathcal{P}_G(u_k)$ is bounded in $H^m(\mathbb{R}^n)$. For $\epsilon > 0$, set $G_\epsilon = \{x \in \mathbb{R}^n : \text{dist}(x, G) < \epsilon\}$, fix $\sigma = \psi_\epsilon \star \chi_{\overline{G}_\epsilon}$ and $\Omega = \sigma^\circ$. Then $\sigma \mathcal{P}_G(u_k)$ is bounded in $H_0^m(\Omega)$ and, hence, has a subsequence $\sigma \mathcal{P}_G(u'_k)$ converging in $H_0^{m-1}(\Omega)$. Since $\sigma \mathcal{P}_G(u'_k)|_G = u'_k$, $\{u'_k\}$ converges in $H^{m-1}(G)$. ■

Boundary Value Problems


Example: For $G \subset \mathbb{R}^n$ and $f : G \rightarrow \mathbb{K}$, find $u : G \rightarrow \mathbb{K}$ satisfying
$$-\Delta u + u = f \text{ in } G$$

- ▶ with *Dirichlet Boundary Conditions*, $u = 0$ on ∂G , or
- ▶ *Neumann Boundary Conditions* $\partial u / \partial \nu = 0$ on ∂G .

For a weak formulation, recall Green's identity:

$$\int_G [v \Delta u + \nabla u \cdot \nabla v] = \int_{\partial G} v \frac{\partial u}{\partial \nu} = \int_{\partial G} \gamma_0(v) \gamma_1(u)$$

holding for sufficiently smooth G , u and v .

- ▶ For the Dirichlet problem, take $\gamma_0(u) = 0$. By , $K(\gamma_0) = H_0^1(G)$, so seek $u \in H_0^1(G)$ with test functions $v \in H_0^1(G)$. Hence, the Green's identity gives:

$$\int_G [v(u - f) + \nabla u \cdot \nabla v] = 0, \quad \forall v \in H_0^1(G)$$

$$\Leftrightarrow (u, v)_{H^1(G)} = (f, v)_{L^2(G)}, \quad \forall v \in H_0^1(G)$$

- ▶ For the Neumann problem, take $\gamma_1(u) = 0$. Now seek $u \in H^1(G)$ with test functions $v \in H^1(G)$, and the Green's identity gives:

$$\int_G [v(u - f) + \nabla u \cdot \nabla v] = 0, \quad \forall v \in H^1(G)$$

$$\Leftrightarrow (u, v)_{H^1(G)} = (f, v)_{L^2(G)}, \quad \forall v \in H^1(G)$$

Introduction


Conversely, if $u \in H^2(G)$ satisfies

$$(u, v)_{H^1(G)} = (f, v)_{L^2(G)}, \forall v \in H_0^1(G) \text{ or } \forall v \in H^1(G)$$

then $C_0^\infty(G) \subset H_0^1(G) \subset H^1(G)$ means

$$(-\Delta u + u, \phi)_{L^2(G)} = (f, \phi)_{L^2(G)}, \forall \phi \in C_0^\infty(G)$$

so $-\Delta u + u = f$ holds in the sense of distributions.

- ▶ If also $u \in H_0^1(G)$, then by Theorem , $\gamma_0(u) = 0$ holds as a boundary condition.
- ▶ Otherwise with $\overline{\text{Rg}(\gamma_0)} = L^2(\partial G)$ and $-\Delta u + u - f = 0 \in L^2(G)$, $0 = (-\Delta u + u - f, v)_{L^2(G)} = (\gamma_1(u), \gamma_0(v))_{L^2(\partial G)}$, $\forall v \in H^1(G)$ means $\gamma_1(u) = 0$ holds as a boundary condition.

Weak formulations of the above boundary value problems:

- ▶ For the Dirichlet problem, find $u \in H_0^1(G) \ni$
 $(u, v)_{H^1(G)} = (f, v)_{L^2(G)}, \forall v \in H_0^1(G)$
- ▶ For the Neumann problem, find $u \in H^1(G) \ni$
 $(u, v)_{H^1(G)} = (f, v)_{L^2(G)}, \forall v \in H^1(G)$

Here, $\gamma_1(u)$ is not (yet and need not be) defined for $u \in H^1(G)$.

Introduction

Theorem: Suppose V is a Hilbert space equipped with $(\cdot, \cdot)_V$ and suppose $b \in V'$. Then $\exists! u \in V \ni (u, v)_V = b(v), \forall v \in V$, and $\|u\|_V = \|b\|_{V'}$.

Proof: Follows from Theorem . ■

Corollary: If $(u_1, v)_V = b_1(v), \forall v \in V$, and $(u_2, v)_V = b_2(v), \forall v \in V$, then $\|u_2 - u_1\|_V = \|b_2 - b_1\|_{V'}$.

Proof: Define $u = u_2 - u_1$ and $b(v) = b_2(v) - b_1(v), v \in V$. ■

Note: The theorem gives a solution for the Dirichlet or the Neumann problem by taking $V = H_0^1(G)$ or $V = H^1(G)$, respectively, and $b(v) = (f, v)_{L^2(G)}$ with $\|b\|_{V'} \leq \|f\|_{L^2(G)}$.

Forms, Operators and Green's Formula

Def: Given a Hilbert space V , a continuous sesquilinear form $a : V \times V \rightarrow \mathbb{K}$ and $b \in V'$, the *abstract boundary value problem* is to find $u \in V \ni$

$$a(u, v) = b(v), \quad \forall v \in V$$

Theorem: Given a continuous sesquilinear form $a : V \times V \rightarrow \mathbb{K}$ on a Hilbert space V , $\exists \alpha, \beta \in \mathcal{L}(V) \ni$

$$a(u, v) = (\alpha(u), v)_V = (u, \beta(v))_V, \quad \forall u, v \in V.$$

Also, given $b \in V'$, $\exists f \in V$, $f = R_V^{-1} b$, \ni

$$b(v) = (f, v)_V, \quad \forall v \in V.$$

Proof: Follows from Theorem . ■

A condition for invertibility of α in $\alpha(u) = R_V^{-1} b$ is as follows.

Def: The sesquilinear form $a : V \times V \rightarrow \mathbb{K}$ is V -coercive if $\exists a_0 > 0 \ni$


$$a(v, v) \geq a_0 \|v\|_V^2, \quad \forall v \in V.$$

Forms, Operators and Green's Formula



Theorem: (Lax-Milgram) Given a Hilbert space V , let $a : V \times V \rightarrow \mathbb{K}$ be a V -coercive continuous sesquilinear form. Then $\forall b \in V', \exists ! u \in V \ni a(u, v) = b(v), \forall v \in V$, and $a_0 \|u\|_V \leq \|b\|_{V'}$.

Proof: Coercivity of a ,

$$a_0 \|v\|_V^2 \leq a(v, v) \leq \|\alpha(v)\|_V \|v\|_V \text{ or } \|v\|_V \|\beta(v)\|_V$$

implies $\text{Rg}(\alpha)$ is closed and $\beta = \alpha^*$ is injective. By Theorem 

$\text{Rg}(\alpha)^\perp = K(\beta) = \{\theta\}$ and thus $\text{Rg}(\alpha) = V$. Then

$u = \alpha^{-1} R_V^{-1} b$ satisfies $a_0 \|u\|_V \leq \|\alpha(u)\|_V = \|R_V^{-1} b\|_V$, and $\|R_V^{-1} b\|_V = \|b\|_{V'}$ since R_V is an isometry . 

Def: Given a Hilbert space V and a continuous sesquilinear form $a : V \times V \rightarrow \mathbb{K}$, the operator $\mathcal{A} \in \mathcal{L}(V, V')$ is defined by

$$a(u, v) = \mathcal{A}u(v), \quad u, v \in V$$

and u solves the abstract boundary value problem when $\mathcal{A}u = b$ holds in V' .

Note: $C_0^\infty(G)$ is not always dense in V , so how to identify V' with $\mathcal{D}^*(G)$ to get a PDE in a distributional sense?

Forms, Operators and Green's Formula

Strategy 1: Assume there is a Hilbert space H (a *pivot space*) satisfying the identifications and continuous imbeddings,

$$V \hookrightarrow H = H' \hookrightarrow V'$$

where $H = H'$ is obtained through the Riesz map. Also, V (and, in practice, $C_0^\infty(G)$ too) is dense in H . Define

$$\mathcal{D} = \{u \in V : \mathcal{A}u \in H'\}$$

where $u \in \mathcal{D}$ iff $u \in V$ and $\exists K > 0 \ni$

$$|a(u, v)| = |\mathcal{A}u(v)| \leq K \|v\|_H, \quad \forall v \in V$$

Then, (in practice) $\mathcal{A}u = b \in H' \Leftrightarrow \mathcal{A}u(\phi) = b(\phi), \forall \phi \in C_0^\infty(G)$, gives a PDE in the distributional sense.

Example: $V = H^1(G)$, $H = L^2(G)$ and $\mathcal{D} = H^2(G)$ for the Neumann problem.

Theorem: Given a Hilbert space V and a V -coercive continuous sesquilinear form $a : V \times V \rightarrow \mathbb{K}$, \mathcal{D} above is dense in V and hence in H .

Forms, Operators and Green's Formula

Proof: To show that \mathcal{A} maps \mathcal{D} onto H' , let $b \in H' \leq V'$ so that, by Theorem 1, $\exists! u \in V \ni a(u, v) = b(v), \forall v \in V$. Then $|a(u, v)| = |b(v)| \leq \|b\|_{H'} \|v\|_H \Rightarrow u \in \mathcal{D}$. Now suppose $\exists w \in V \ni (u, w)_V = 0, \forall u \in \mathcal{D}$. As in the proof of Theorem 1, $\text{Rg}(\beta) = V$, so $\exists v \in V \ni \beta(v) = w$. Hence, $0 = (u, w)_V = (u, \beta(v))_V = \mathcal{A}u(v)$. For $u \in \mathcal{D}$, $\mathcal{A}u(v) = (R_H^{-1} \mathcal{A}u, v)_H$. Since $\mathcal{A}\mathcal{D} = H'$, choose $u \in \mathcal{D} \ni \mathcal{A}u = R_H v$. Then $0 = \mathcal{A}u(v) = \|v\|_H^2 \Rightarrow w = \beta(v) = 0$. ■

Strategy 2: Assume there is a closed subspace V_0 of V satisfying the identifications and continuous imbeddings,

$$V_0 \leq V \hookrightarrow H = H' \hookrightarrow V' \leq V'_0$$

where V_0 (in practice, the completion of $C_0^\infty(G)$ in V) is dense in H . Define $A \in \mathcal{L}(V, V'_0)$ by

$$a(u, v) = Au(v), \quad u \in V, v \in V_0$$

as the *formal operator* determined by a , V and V_0 , and set

$$D = \{u \in V : Au \in H'\}$$

Forms, Operators and Green's Formula

where $u \in D$ iff $u \in V$ and $\exists K > 0 \ni$

$$|a(u, v)| = |Au(v)| \leq K \|v\|_H, \quad \forall v \in V_0$$

Then, (in practice) $Au = f \in H' \Leftrightarrow Au(\phi) = f(\phi), \forall \phi \in C_0^\infty(G)$, gives a PDE in the distributional sense.

Example: $V_0 = H_0^1(G)$, $V = H^1(G)$, $H = L^2(G)$,

$R_H^{-1}Au = -\Delta u + u$ and $D = \{u \in H^1(G) : \Delta u \in L^2(G)\}$ for the Neumann problem.

To compare \mathcal{A} and A , note that $D \subset \mathcal{D}$, fix $u \in D$ and define

$$\varphi_u(v) = \mathcal{A}u(v) - Au(v), \quad v \in V$$

satisfying $\varphi_u \in V'$ and $\varphi_u|_{V_0} = 0$. Define

$$\hat{\varphi}_u(\hat{v}) = \varphi_u(v), \quad \hat{v} = \{v + v_0 : v_0 \in V_0\} \in (V/V_0).$$

Then

$$|\hat{\varphi}_u(\hat{v})| = \inf_{v_0 \in V_0} |\varphi_u(v + v_0)| \leq 2K \inf_{v_0 \in V_0} \|v + v_0\|_V = 2K \|\hat{v}\|_{V/V_0}$$

so $\hat{\varphi}_u \in (V/V_0)'$.

Forms, Operators and Green's Formula

Assume there is a trace operator γ which is a linear surjection onto a Hilbert space $B = \text{Rg}(\gamma)$ and $V_0 = K(\gamma)$.


Define also on V/V_0 ,

$$\hat{\gamma}(\hat{v}) = \gamma(v), \quad \hat{v} = \{v + v_0 : v_0 \in V_0\} \in (V/V_0).$$

Let B be equipped with the norm

$$\|\hat{\gamma}(\hat{v})\|_B = \|\hat{v}\|_{V/V_0}$$

Then $\hat{\gamma} \in \mathcal{L}(V/V_0, B)$ and $\hat{\gamma}$ is norm preserving.

Hence $\hat{\gamma}' \in \mathcal{L}(B', (V/V_0)')$ and $\hat{\gamma}'$ is injective .

Furthermore, since for $v \in V$

$\|\gamma(v)\|_B = \|\hat{\gamma}(\hat{v})\|_B = \|\hat{v}\|_{V/V_0} = \inf_{v_0 \in V_0} \|v + v_0\|_V \leq \|v\|_V$
it follows that $\gamma \in \mathcal{L}(V, B)$.

To see that $\hat{\gamma}'$ is surjective, let $f \in (V/V_0)'$ and define $d \in B'$ by $d(g) = f(\hat{\gamma}^{-1}(g))$, $g \in B$, so that $\hat{\gamma}'(d) = d \circ \hat{\gamma} = f$.

Forms, Operators and Green's Formula

Since $\hat{\gamma}'$ is bijective, it follows that for $\hat{\varphi}_u \in (V/V_0)'$ given above, $\exists! \partial u \in B' \ni \hat{\gamma}'(\partial u) = \partial u \circ \hat{\gamma} = \hat{\varphi}_u$. Combining these results gives

$$\partial u(\gamma v) = \partial u(\hat{\gamma}(\hat{v})) = \hat{\varphi}_u(\hat{v}) = \varphi_u(v)$$

for $\hat{v} = \{v + v_0 : v_0 \in V_0\} \in (V/V_0)$ with linear dependence upon u . This result is summarized as follows.

Theorem: Under above stated **assumptions**,

$$\exists \partial \in L(D, B') \ni \forall u \in D, \partial u(\gamma v) = Au(v) - Au(v), \forall v \in V$$

This result is called the *abstract Green's identity* where ∂ is the *abstract Green's operator*.

Study Question: Determine a norm on D so that $\partial \in \mathcal{L}(D, B')$.

Example: For the Neumann problem, $V = H^1(G)$, $V_0 = H_0^1(G)$, $\gamma = \gamma_0$, $B = H^{\frac{1}{2}}(\partial G) \leq L^2(\partial G) = L^2(\partial G)' \leq B'$ and ∂ is an extension of γ_1 from $H^2(G)$ to D .

Abstract Boundary Value Problems

As in the Note ,

- ▶ If boundary conditions are explicitly prescribed in V (e.g., Dirichlet boundary conditions with $V = H_0^1(G)$) then these are called *forced* or *stable* boundary conditions.
- ▶ If boundary conditions are not explicitly prescribed with V (e.g., Neumann boundary conditions in $V = H^1(G)$) then these are called *variational (natural)* or *unstable* boundary conditions.

Assumptions (corresponding to Strategy 2):

- ▶ There is a closed subspace V_0 of a Hilbert space V satisfying the identifications and continuous imbeddings,
$$V_0 \leq V \hookrightarrow H = H' \hookrightarrow V' \leq V_0'$$
- ▶ $\exists \gamma \in \mathcal{L}(V, B)$ where $B = \text{Rg}(\gamma)$ is isomorphic to V/V_0 .
- ▶ $V_0 = K(\gamma)$ is dense in H .

(In practice, V_0 is the completion of $C_0^\infty(G)$ in V , and the pivot space is $H = L^2(G)$.)

Abstract Boundary Value Problems

- ▶ $\exists a_1 : V \times V \rightarrow \mathbb{K}$ and $a_2 : B \times B \rightarrow \mathbb{K}$ both continuous such that

$$a(u, v) = a_1(u, v) + a_2(\gamma u, \gamma v), \quad u, v \in V$$

(e.g., $a_1(u, v) = (\nabla u, \nabla v)_{L^2(G)}$,
 $a_2(\varphi, \psi) = (\varphi, \psi)_{L^2(\partial G)}$)

- ▶ $\exists b_1 \in H'$, $b_2 \in B'$ such that

$$b(v) = b_1(v)_H + b_2(\gamma v), \quad v \in V$$

(e.g., $b_1(v) = (f, v)_H$,
 $b_2(\psi) = (g, \psi)_{L^2(\partial G)}$)

Consequences:

- ▶ Define $\mathcal{A}_2 \in \mathcal{L}(B, B')$ by

$$\mathcal{A}_2 \varphi(\psi) = a_2(\varphi, \psi), \quad \varphi, \psi \in B$$

(e.g., $R_B^{-1} \mathcal{A}_2 \varphi = \varphi \in B = H^{\frac{1}{2}}(G)$)

- ▶ Define $A_1 \in \mathcal{L}(V, V'_0)$ by

$$A_1 u(v) = a_1(u, v), \quad u \in V, \quad v \in V_0$$

(e.g., $R_H^{-1} A_1 u = -\Delta u \in H = L^2(G)$)

- ▶ Define $D_1 = \{u \in V : A_1 u \in H'\}$

(e.g., $u \in D_1 \Rightarrow \Delta u \in L^2(G)$)

Abstract Boundary Value Problems

- According to Theorem  define $\partial_1 \in L(D_1, B')$ by

$$a_1(u, v) - A_1 u(v) = \partial_1 u(\gamma v), \quad u \in D_1, \quad v \in V \quad (1)$$

(e.g., ∂_1 extends γ_1)

Theorem: Assume the Hilbert spaces, forms and operators are given as above. Then $u \in V$ solves

$$a_1(u, v) + a_2(\gamma u, \gamma v) = b_1(v) + b_2(\gamma v), \quad \forall v \in V \quad (2)$$

if and only if $u \in D_1$ solves

$$A_1 u = b_1 \quad \text{and} \quad \partial_1 u + \mathcal{A}_2(\gamma u) = b_2. \quad (3)$$

Proof: Let $u \in V$ solve (2). Choosing $v \in V_0 = K(\gamma)$ in (2) gives $|a_1(u, v)| = |b_1(v)| \leq \|b\|_{H'} \|v\|_H, \forall v \in V_0$, so $u \in D_1$. Thus, $(R_H^{-1} A_1 u, v)_H = A_1 u(v) = b_1(v) = (R_H^{-1} b_1, v)_H, \forall v \in V_0$. Since V_0 is dense in H , the first equation in (3) is obtained.

Thus, (1) and (2) may be combined to give

$$\partial_1 u(\gamma v) = g(\gamma v) - a_2(\gamma u, \gamma v) = g(\gamma v) - \mathcal{A}_2(\gamma u)(\gamma v)$$

$\forall v \in V$, and the second equation in (3) follows. Now let $u \in D_1$ solve (3). Combining the previous equation with (1) gives (2) after $A_1 u(v)$ is replaced by $(F, v)_H$. ■

Examples

Suppose $G \subset \mathbb{R}^n$ is open and bounded and that $\mathbf{A} = \{a_{ij}\}_{i,j=1}^n$, $\mathbf{a} = \{a_j\}_{j=1}^n$, a_0 satisfy $a_{i,j}, a_j, a_0 \in L^\infty(G)$. Then define

$$a_1(u, v) = \int_G \{ \nabla u^T \mathbf{A} \nabla v + \mathbf{a}^T \nabla u v + a_0 u v \}$$

Take $H = L^2(G)$. Fix $\Gamma \subset \partial G$ and define the closed subspace

$$V = \{ v \in H^1(G) : (\gamma_0 v)|_\Gamma = 0 \} \leq H^1(G)$$

Take $V_0 = H_0^1(G)$, $\gamma = \gamma_0|_V$ and $B = \text{Rg}(\gamma)$. Then A_1 satisfies

$$R_H^{-1} A_1 u = -\nabla \cdot (\mathbf{A} \nabla u) + \mathbf{a}^T \nabla u + a_0 u, \quad u \in D_1$$

where $D_1 = \{ u \in V : A_1 u \in H' \}$. ($= \emptyset$ when $a_{i,j}, a_j, a_0 \notin C^1(\overline{G})$?)


According to Theorem  define $\partial \in L(D_1, B')$ by

$$a_1(u, v) - A_1 u(v) = \partial u(\gamma v), \quad u \in D_1, \quad v \in V$$

For $a_{i,j}, u, v \in C^\infty(\overline{G})$, the Green's Theorem gives

$$a_1(u, v) - A_1 u(v) = \int_{\partial G \setminus \Gamma} \partial_{\mathbf{A}} u(\gamma_0 v), \quad \partial_{\mathbf{A}} u = \nu^T \mathbf{A} \nabla u$$

so ∂ extends $\partial_{\mathbf{A}}$ from $H^2(G)$ to D_1 . Define $b_1(v) = (f, v)_{L^2(G)}$.

By Theorem  the weak solution, $a_1(u, v) = b_1(v)$, $\forall v \in V$, is a generalized solution to

$$R_H^{-1} A_1 u = f \text{ in } G, \quad u = 0 \text{ on } \Gamma, \quad \partial_{\mathbf{A}} u = 0 \text{ on } \partial G \setminus \Gamma$$

called a *mixed* Dirichlet-Neumann BVP for $|\Gamma|, |\partial G \setminus \Gamma| > 0$.

Examples

It is a purely Dirichlet (type 1) BVP if $|\partial G \setminus \Gamma| = 0$ and a purely Neumann (type 2) BVP if $|\Gamma| = 0$.

Example of a Robin (type 3) BVP: Define $H = L^2(G)$, $V_0 = H_0^1(G)$, $V = H^1(G)$, $\gamma = \gamma_0$, $B = \text{Rg}(\gamma)$ and

$$a_1(u, v) = \int_G \nabla u \cdot \nabla \bar{v}, \quad u, v \in V.$$

Then $R_H^{-1} A_1 u = -\Delta u$ for $u \in D_1 = \{u \in V : A_1 u \in H'\}$.

According to Theorem  define $\partial_1 \in L(D_1, B')$ by


$$a_1(u, v) - A_1 u(v) = \partial_1 u(\gamma v), \quad u \in D_1, \quad v \in V$$

which extends γ_1 from $H^2(G)$ to D_1 .

Suppose $f \in L^2(G)$, $g \in L^2(\partial G)$ and $\alpha \in L^\infty(\partial G)$ and define

$$a_2(\varphi, \psi) = \int_{\partial G} \alpha \varphi \bar{\psi}, \quad \varphi, \psi \in L^2(\partial G)$$

$$b(v) = b_1(v) + b_2(v) = (f, v)_{L^2(G)} + (g, \gamma_0 v)_{L^2(\partial G)}, \quad v \in V.$$

By Theorem  the weak solution u to

$$a(u, v) = b(v), \quad \forall v \in V$$

is a generalized solution to

$$-\Delta u = f \text{ in } G, \quad \partial_\nu u + \alpha u = g \text{ on } \partial G.$$

Examples

For $\alpha = 0$, this is an inhomogeneous Neumann BVP.
How to formulate an inhomogeneous Dirichlet BVP?

For example, consider the Dirichlet BVP,

$$-\Delta u = F \text{ in } G, \quad u = g \text{ on } \partial G$$

where $g = \gamma_0 g_e$ for some $g_e \in H^2(G)$. Thus, g_e is an extension or a *lifting* of g from ∂G to G .


With $w = u - g_e$ and $\tilde{F} = F + \Delta g_e$ the above BVP becomes

$$-\Delta w = \tilde{F} \text{ in } G, \quad w = 0 \text{ on } \partial G$$

For this, take $H = L^2(G)$, $V_0 = V = H_0^1(G)$, $\gamma = \gamma_0$ and

$$a_1(w, v) = A_1 w(v) = \int_G \nabla w \cdot \nabla \bar{v}, \quad w, v \in V,$$

so $A_1 w = -R_H \Delta w$ for $w \in D_1 = \{w \in V : A_1 w \in H'\}$.

By Theorem  the weak solution w to

$$a_1(w, v) = (\tilde{F}, v)_H, \quad \forall v \in V$$

is a generalized solution to

$$R_H^{-1} A_1 w = \tilde{F} \text{ in } G, \quad w = 0 \text{ on } \partial G$$

and $u = w + g_e$ is a generalized solution to the above inhomogeneous Dirichlet BVP.

Examples

Suppose $G = \hat{G} \cup \check{G} \cup \Sigma^\circ$ for open \hat{G}, \check{G} where $\hat{G} \cap \check{G} = \emptyset$ and $\Sigma = \partial\hat{G} \cap \partial\check{G}$. Assume $\partial\hat{G}, \partial\check{G}$ are sufficiently regular.

Let $\hat{\nu}$ and $\check{\nu}$ be the outward unit normals at \hat{G} and \check{G} with $\hat{\nu} + \check{\nu} = 0$ at Σ . With $H = L^2(G)$ and

$$[v] = \gamma_0|_{\hat{G}} v - \gamma_0|_{\check{G}} v$$

set

$$V = \{v \in H : [v] = 0, v \in H^1(\hat{G}), v \in H^1(\check{G})\} \quad (H^1(G)?)$$

and

$$V_0 = \{v \in H : v \in H_0^1(\hat{G}), v \in H_0^1(\check{G})\}.$$

so that $V_0 = K(\gamma)$ for $\gamma = \gamma_0|_V$ on $\partial\hat{G} \cup \partial\check{G}$ taking $B = \text{Rg}(\gamma) \subset L^2(\partial\hat{G} \cup \partial\check{G})$.

For $\hat{a} \in C^1(\overline{\hat{G}_k})$, $\check{a} \in C^1(\overline{\check{G}_k})$, define continuous

$$a_1(u, v) = A_1 u(v) = \int_{\hat{G}} \hat{a} \nabla u \cdot \nabla v + \int_{\check{G}} \check{a} \nabla u \cdot \nabla v, \quad u, v \in V$$

Then taking $v \in V_0$ and $u \in D_1 = \{u \in V : A_1 u \in H'\}$ gives

$$R_H^{-1} A_1 u = -\nabla \cdot (\hat{a} \nabla u) \text{ in } \hat{G} \quad \text{and} \quad -\nabla \cdot (\check{a} \nabla u) \text{ in } \check{G}$$

Examples

So by Theorem $\blacktriangleright \exists \partial_1 \in L(D_1, B)$ satisfying

$$\partial_1 u(v) = a_1(u, v) - A_1 u(v), \quad u \in D_1, \quad v \in V$$

For $u, v \in C^\infty(\overline{G})$, the Green's formula gives

$$\begin{aligned} a_1(u, v) - A_1 u(v) &= \int_{\partial \hat{G}} \hat{a} \bar{v} \partial_{\nu_1} u + \int_{\partial \check{G}} \check{a} \bar{v} \partial_{\nu_2} u \\ &= \int_{\partial G \cap \partial \hat{G}} \bar{v} \hat{a} \partial_{\hat{\nu}} u + \int_{\partial G \cap \partial \check{G}} \check{a} \partial_{\check{\nu}} u + \int_{\Sigma} \gamma_0 \bar{v} [\hat{a} \partial_{\hat{\nu}} + \check{a} \partial_{\check{\nu}}] u \end{aligned}$$

$$\text{so } \partial_1 \text{ extends } \begin{cases} \hat{a} \partial_{\hat{\nu}} & \text{on } \partial \hat{G} \setminus \Sigma \\ \check{a} \partial_{\check{\nu}} & \text{on } \partial \check{G} \setminus \Sigma \\ \hat{a} \partial_{\hat{\nu}} + \check{a} \partial_{\check{\nu}} & \text{on } \Sigma \end{cases}$$

from $H^2(G)$ to D_1 . For $f \in L^2(G)$ and $g \in L^2(\partial G)$ define
 $b(v) = b_1(v) + b_2(v) = (f, v)_H + (g, \gamma_0 v)_{L^2(\partial G)}$.

By Theorem \blacktriangleright the weak solution $u \in V$ to

$$a_1(u, v) = b(v), \quad \forall v \in V,$$

is a generalized solution to

$$\begin{aligned} R_H^{-1} A_1 u &= f \text{ in } G, \quad [u] = 0 \text{ on } \Sigma, \\ \hat{a} \partial_{\hat{\nu}} u &= g \text{ on } \partial G \cap \partial \hat{G}, \quad \check{a} \partial_{\check{\nu}} u = g \text{ on } \partial G \cap \partial \check{G}, \\ \hat{a} \partial_{\hat{\nu}} u + \check{a} \partial_{\check{\nu}} u &= 0 \text{ on } \Sigma \end{aligned}$$

Coercivity and Elliptic Forms

Suppose $G \subset \mathbb{R}^n$ is open and that $\mathbf{A} = \{a_{ij}\}_{i,j=1}^n$, $\mathbf{a} = \{a_j\}_{j=1}^n$, a_0 satisfy $a_{i,j}, a_j, a_0 \in L^\infty(G)$. Then define

$$a(u, v) = \int_G \{ \nabla u^T \mathbf{A} \nabla \bar{v} + \mathbf{a}^T \nabla u \bar{v} + a_0 u \bar{v} \}$$

Def: The form a is *strongly elliptic* if $\exists c_0 > 0 \ni$

$$\Re \xi^T \mathbf{A} \bar{\xi} \geq c_0 |\xi|^2, \quad \forall \xi \in \mathbb{K}^n, \quad \forall x \in G$$

Theorem: If the form a is strongly elliptic $\exists \lambda_0 \in \mathbb{R} \ni \forall \lambda > \lambda_0$ the form $a(u, v) + \lambda(u, v)_{L^2(G)}$ is coercive on $H^1(G)$.

Proof: Set $K_1 = \max\{\|a_j\|_{L^\infty(G)}\}_{j=1}^n$, $K_0 = \text{ess inf}\{\Re a_0(x) : x \in G\}$. Then for $1 \leq j \leq n$ and $\forall \epsilon > 0$,

$$\begin{aligned} |(a_j \partial_j u, u)_{L^2(G)}| &\leq K_1 \|\partial_j u\|_{L^2(G)} \|u\|_{L^2(G)} \\ &\leq \frac{1}{2} K_1 [\epsilon \|\partial_j u\|_{L^2(G)}^2 + \|u\|_{L^2(G)}^2 / \epsilon] \end{aligned}$$

Also, $\Re(a_0 u, u) \geq K_0 \|u\|_{L^2(G)}^2$.

Combining with the ellipticity condition gives $\forall u \in H^2(G)$,

$$\Re a(u, u) \geq (c_0 - \frac{1}{2} \epsilon K_1) \|\nabla u\|_{L^2(G)}^2 + (K_0 - \frac{1}{2} n K_1 / \epsilon) \|u\|_{L^2(G)}^2$$

The result follows with $K_1 \epsilon \leq c_0$ and $\lambda_0 = \frac{1}{2} n K_1^2 / c_0 - K_0$. ■

Coercivity and Elliptic Forms

Corollary: Set $H = L^2(G)$, $V_0 = H_0^1(G)$ and $V = \{v \in H^1(G) : (\gamma_0 v)|_\Gamma = 0\}$ for $\Gamma \subset \partial G$, and define A_λ so that

$$a(u, v) + \lambda(u, v) = A_\lambda u(v), \quad u \in V, v \in V_0$$

Then for $f \in L^2(G)$ and $\lambda > \lambda_0$, the BVP

$$a(u, v) + \lambda(u, v) = (f, v)_{L^2(G)}, \quad \forall v \in V$$

is well-posed and the solution satisfies


$u \in D_\lambda = \{u \in V : A_\lambda u \in H'\}$ and

$$\|(\lambda - \lambda_0)u\|_{L^2(G)} \leq \|f\|_{L^2(G)}$$


Proof: Well-posedness follows from Theorems  and .

Then Theorem  gives $u \in D_\lambda$. Since

$$\|R_H^{-1} A_\lambda u\|_{L^2(G)} = \|F\|_{L^2(G)},$$

the claimed estimate follows with estimates from the proof of Theorem ,

$$\begin{aligned} (\lambda - \lambda_0)\|u\|_{L^2(G)}^2 &\leq a(u, u) + \lambda(u, u)_{L^2(G)} \\ &= A_\lambda u(u) \leq \|R_H^{-1} A_\lambda u\|_{L^2(G)} \|u\|_{L^2(G)} \end{aligned}$$

after factoring $\|u\|_{L^2(G)}$. 

Coercivity and Elliptic Forms

Theorem: Let G be open in \mathbb{R}^n and suppose $0 \leq x_n \leq K$, $\forall x = (x', x_n) \in G$. Let ∂G be a C^1 manifold with G on one side of ∂G . Let $\nu = (\nu_1, \dots, \nu_n)$ be the outward unit normal at ∂G with $\nu_n > 0$ on $\Sigma \subset \partial G$. Then $\forall u \in H^1(G)$,

$$\int_G |u|^2 \leq 2K \int_\Sigma |\gamma_0 u|^2 + 4K^2 \int_G |D_n u|^2$$

Proof: For $u \in C^1(\overline{G})$, the Gauss Theorem gives

$$\int_{\partial G} \nu_n s_n |u|^2 = \int_G D_n(x_n |u|^2) = \int_G |u|^2 + \int_G u(2x_n D_n u)$$

The last term satisfies

$$\begin{aligned} \left| \int_G 2x_n u D_n u \right| &\leq \frac{1}{2} \int_G |u|^2 + \frac{1}{2} \int_G |2x_n D_n u|^2 \\ &\leq \frac{1}{2} \int_G |u|^2 + 2K^2 \int_G |D_n u|^2 \end{aligned}$$

Combining these estimates gives

$$\begin{aligned} \int_G |u|^2 &= \int_{\partial G} \nu_n s_n |u|^2 - \int_G D_n(x_n |u|^2) \\ &\leq \int_\Sigma \nu_n s_n |u|^2 + \frac{1}{2} \int_G |u|^2 + 2K^2 \int_G |D_n u|^2 \end{aligned}$$

since $\nu_n s_n \leq 0$ on $\partial G \setminus \Sigma$. With $0 < \nu_n s_n \leq K$ on Σ , the claimed estimate follows. ■

Coercivity and Elliptic Forms

Corollary: If a_1 of the mixed Dirichlet-Neumann BVP is strongly elliptic and its coefficients additionally satisfy $a_j = 0$, $1 \leq j \leq n$, and $\Re a_0 \geq 0$ in G and $\{x \in \partial G : \nu_n > 0\} = \Sigma \subseteq \Gamma \subseteq \partial G$, then the BVP is well-posed.

Proof: According to Theorem , the form $a(u, v) = a_1(u, v) + a_2(u, v)$ with

$$a_2(u, v) = \int_{\Sigma} \gamma_0 u \gamma_0 v$$

is coercive on all of $H^1(G)$ and so in particular on the subspace $V = \{v \in H^1(G) : (\gamma_0 v)|_{\Gamma} = 0\}$ where $a = a_1$. Also, $a = a_1$ is bounded on $V \times V$, and $b = b_1$ is bounded on V . Thus, well-posedness follows from Theorems and .

Study Question: Show that the weak Robin BVP is well posed for $\alpha > 0$.

Study Question: For data c, f supported on $S \subset G$ with $|S| > 0$, define $a(u, v) = \int_G [\nabla^2 u : \nabla^2 v + cuv]$ and $b(v) = \int_G fv$ for $u, v \in H^2(G)$ and show well-posedness of the problem $a(u, v) = b(v), \forall v \in H^2(G)$. Hint: Adapt the following.

Coercivity and Elliptic Forms

Theorem: If a_1 of the mixed Dirichlet-Neumann BVP is strongly elliptic and its coefficients additionally satisfy $a_j = 0$, $1 \leq j \leq n$, and $\Re a_0 \geq 0$ in a bounded G , then the BVP is well-posed.

Proof: Clearly, the form a_1 is bounded on $V \times V$ and the form b_1 is bounded on V . To show that a_1 is coercive on $H^1(G)$, suppose $\exists \{u_k\} \subset V \ni \|u_k\|_{H^1(G)} = 1$ while $a_1(u_k, u_k) \xrightarrow{k \rightarrow \infty} 0$. Then $H^1(\Omega)$ boundedness implies weak subsequential convergence in V . Let $\{u_k\}$ again denote the subsequence. By compactness of $H^1(\Omega)$ in $L^2(\Omega)$, the sequence converges strongly in $L^2(\Omega)$. Because of $a_1(u_k, u_k) \xrightarrow{k \rightarrow \infty} 0$, the sequence converges strongly in V to some u_0 , which must satisfy $a_1(u_0, u_0) = \lim_{k \rightarrow \infty} a_1(u_k, u_k) = 0$. So $u_0 \in V$ is constant, and $u_0|_\Gamma = 0$ means $u_0 = 0$. This contradiction of $\|u_0\|_{H^1(\Omega)} = \lim_{k \rightarrow \infty} \|u_k\|_{H^1(\Omega)} = 1$ implies coercivity. Thus, well-posedness follows from Theorems and .

Regularity

Theorem: Let G be bounded and open in \mathbb{R}^n and suppose ∂G is a C^2 -manifold of dimension $n - 1$. Set $H = L^2(G)$ and $V = H^1(G)$. Let $a_{i,j}, a_j, a_0 \in C^1(\overline{G})$ and assume that

$$a(u, v) = \int_G \{ \nabla u^T \mathbf{A} \nabla v + \mathbf{a}^T \nabla u v + a_0 u v \}$$

is strongly elliptic. Let $f \in H$ and suppose $u \in V$ satisfies

$$a(u, v) = (f, v)_{L^2(G)}, \quad \forall v \in V$$

Then $u \in H^2(G)$. The same holds for $V = H_0^1(G)$. ■

Def: Let V be a closed subspace of $H^1(G)$ with $H_0^1(G) \leq V$, and let a be a continuous sesquilinear form on V . Then a is called k -regular on V if $\forall f \in H^s(G)$, $0 \leq s \leq k$ and $\forall u \in V$ solving $a(u, v) = (f, v)_{L^2(G)}$, $\forall v \in V$, we have $u \in H^{2+s}(G)$.

Theorem: The form a of Theorem [▶](#) is k -regular over $H^1(G)$ and $H_0^1(G)$ if ∂G is a C^{k+2} -manifold and $a_{i,j}, a_j, a_0 \in C^{k+1}(\overline{G})$. ■

Closed Operators, Adjoins, Eigenfunction Expansions

Def: Given a Hilbert space H and a $D \leq H$ with $A \in L(D, H)$, the *graph* of A is the subspace

$$G(A) = \{[x, Ax] : x \in D\}$$



of the Hilbert space $H \times H$ equipped with

$$([x_1, x_2], [y_1, y_2])_{H \times H} = (x_1, y_1)_H + (x_2, y_2)_H$$

and A is *closed* on H if $G(A)$ is a closed subset of $H \times H$.

Lemma: If A is closed and continuous, then D is closed.

Proof: If $D \ni x_n \rightarrow x \in H$, then $\{x_n\}$ and hence $\{Ax_n\}$ are Cauchy sequences. Since H is complete, $\exists y \in H \ni Ax_n \rightarrow y$. Since $G(A)$ is closed, $[x, y] \in G(A)$, $y = Ax$ and $x \in D$. ■

Def: If $\overline{D} = H$, the adjoint A^* is defined with domain D^* , the subspace of $y \in H$ such that $y \mapsto (Ax, y)_H$ is in H' for every $x \in D$. By , this functional has a unique continuous extension to H , and applying Theorem  to the extension means $\exists! A^*y \in H \ni$


$$(Ax, y)_H = (x, A^*y)_H, \quad x \in D, \quad y \in D^*.$$

Closed Operators, Adjoint, Eigenfunction Expansions

Lemma: A^* is closed.

Proof: Choose $y_n \in D^* \ni y_n \rightarrow y$ and $A^*y_n \rightarrow z$. Then for $x \in D$, $(Ax, y)_H \leftarrow (Ax, y_n)_H = (x, A^*y_n)_H \rightarrow (x, z)$. Since (x, z) is continuous at $x \in D$, so is (Ax, y) . Thus, $y \in D^*$ and $z = A^*y$. ■

Lemma: If $D = H$, then A^* is continuous and hence D^* is closed.


Proof: If A^* is not continuous, $\exists \{y_n\} \subset D^* \ni \|y_n\|_H = 1$ and $\|A^*y_n\|_H \rightarrow \infty$. By $(Ax, y)_H = (x, A^*y)_H$, $x \in D$, $y \in D^*$, it follows that $|(x, A^*y_n)_H| = |(Ax, y_n)_H| \leq \|Ax\|_H$, $\forall x \in H$, so $\{A^*y_n\}$ is weakly bounded. By Theorem , $\{A^*y_n\}$ is bounded, a contradiction. ■

Lemma: If A is closed, then D^* is dense in H .

Closed Operators, Adjoins, Eigenfunction Expansions

Proof: Suppose $0 \neq y \in (D^*)^\perp$. Since $A \in L(D, H)$, $[0, y] \notin G(A)$. Since $G(A)$ is closed, $G(A) \neq H \times H$, so define the projection $P : H \times H \rightarrow G(A)^\perp$. Then define $[u, v] = P[0, y]$ so that $f(x_1, x_2) = (u, x_1)_H + (v, x_2)_H$, $x_1, x_2 \in H$, satisfies $f(0, y) \neq 0$ and $f(G(A)) = 0$. By $0 = (u, x)_H + (v, Ax)_H = f(x, Ax)$, $x \in D$, the continuity of $x \mapsto (v, Ax)_H$ follows from the continuity of $x \mapsto (u, x)_H$. Hence, $v \in D^*$, and $(v, y)_H = f(0, y) \neq 0$. But this contradicts the assumption that $y \in (D^*)^\perp$. Hence, $(D^*)^\perp = \{0\}$, so D^* is dense in H . ■

Theorem: (Closed-Graph) For $D \leq H$, suppose $A \in L(D, H)$. Then A is closed and $D = H$ if and only if $A \in \mathcal{L}(H)$.

Proof: If A is closed and $D = H$, then the last two lemmas imply that $A^* \in \mathcal{L}(H)$. Then Theorem  shows $(A^*)^* \in \mathcal{L}(H)$. But

$$(Ax, y)_H = (x, A^*y)_H, \quad \forall x \in D, \quad \forall y \in D^*$$

with $D = H = D^*$ shows $A = (A^*)^*$, so $A \in \mathcal{L}(H)$. Conversely, $A \in \mathcal{L}(H)$ means that $x_n \xrightarrow{n \rightarrow \infty} x \Rightarrow Ax_n \xrightarrow{n \rightarrow \infty} Ax$, so A is closed. ■

Closed Operators, Adjoins, Eigenfunction Expansions

Example: Take $H = L^2(G)$ and $G = (0, 1)$. Let $A = iD$ and $D(A) = H_0^1(G)$. If $G(A) \ni [u_n, Au_n] \rightarrow [u, v] \in H \times H$ then

$$\begin{aligned} \int_0^1 \underbrace{Au_n}_{\downarrow} \bar{\varphi} &= -i \int_0^1 \underbrace{u_n}_{\downarrow} D\bar{\varphi}, \quad \varphi \in C_0^\infty(G) \\ \int_0^1 v \bar{\varphi} &= -i \int_0^1 u D\bar{\varphi} \end{aligned}$$

so that $v = iDu = Au$ and $u_n \xrightarrow{H^1(G)} u$. Hence, $u \in H_0^1(G)$ and A is closed. To determine the adjoint, note that

$$\int_0^1 Au \bar{v} = \int_0^1 u \bar{f}, \quad \forall u \in H_0^1(G) \quad (= D(A))$$

holds for some $v, f \in L^2(G)$ if and only if $f = iDv$ and $v \in H^1(G)$. (Alternative Definition!) Thus $D(A^*) = H^1(G)$ and $A^* = iD$ is an extension of A .

Closed Operators, Adjoins, Eigenfunction Expansions

Since A^* is an adjoint, it is closed. To determine the adjoint $A^{**} = (A^*)^*$, note that $[u, f] \in G(A^{**})$ if and only if

$$\int_0^1 A^* v \bar{u} = \int_0^1 v \bar{f}, \quad \forall v \in H^1(G) \quad (= D(A^*))$$

Since this holds $\forall v \in C_0^\infty(G)$, it must be that $f = iDu$.
Substituting shows that

$$i \int_0^1 Dv \bar{u} = -i \int_0^1 v D\bar{u}, \quad \forall v \in H^1(G) \quad (= D(A^*))$$

or

$$0 = \int_0^1 D(v\bar{u}) = v(1)\bar{u}(1) - v(0)\bar{u}(0), \quad \forall v \in H^1(G)$$

implying $u(0) = u(1) = 0$, so $u \in H_0^1(G) = D(A^{**})$. Hence,

$$\int_0^1 Au \bar{v} = \int_0^1 u A^* \bar{v} = \int_0^1 A^{**} u \bar{v}, \quad \forall u \in H_0^1(G), \forall v \in H^1(G)$$

it follows that $A^{**} = A$.

Closed Operators, Adjoins, Eigenfunction Expansions

Example: Take $H = L^2(G)$ and $G = (0, 1)$. Let $B = iD$ with
 $D(B) = \{u \in H^1(G) : u(0) = cu(1)\}$
for some $c \in \mathbb{C}$. Then for $v, f \in L^2(G)$, $B^*v = f$ if and only if

$$\int_0^1 iDu\bar{v} = \int_0^1 u\bar{f}, \quad \forall u \in D(B)$$

Taking $u \in C_0^\infty(G) \leq D(B)$ gives $f = iDv (= B^*v)$ and $v \in H^1(G)$. Substituting shows that

$$0 = i \int_0^1 D(u\bar{v}) = iu(1)[\bar{v}(1) - c\bar{v}(0)], \quad \forall u \in D(B)$$

implying $\bar{v}(1) - c\bar{v}(0) = 0$, so

$$D(B^*) = \{v \in H^1(G) : v(1) = \bar{c}v(0)\}.$$

and $B^* = iD$.

Closed Operators, Adjoins, Eigenfunction Expansions

Hilbert spaces V, H are given with V dense in H and $V \hookrightarrow H$.

Suppose a is a sesquilinear form, continuous on V . Recall

$$a(u, v) = \mathcal{A}u(v), \quad u, v \in V \text{ and } \mathcal{D} = \{u \in V : \mathcal{A}u \in H'\}.$$

Now take

$$D(\mathbb{A}) = \mathcal{D} = \{u \in V : \exists K > 0 \ni |a(u, v)| \leq K\|v\|_H, \forall v \in V\}$$

and $\mathbb{A} \in L(D(\mathbb{A}), H)$ defined by

$$a(u, v) = (\mathbb{A}u, v)_H, \quad \forall u \in D(\mathbb{A}), \quad \forall v \in V.$$

Define the adjoint sesquilinear form

$$r(u, v) = \overline{a(v, u)}, \quad u, v \in V$$

with

$$D(\mathbb{R}) = \{u \in V : \exists K > 0 \ni |r(u, v)| \leq K\|v\|_H, \forall v \in V\}$$

and $\mathbb{R} \in L(D(\mathbb{R}), H)$ defined by



$$r(u, v) = (\mathbb{R}u, v)_H, \quad \forall u \in D(\mathbb{R}), \quad \forall v \in V.$$

Theorem: If $\exists \lambda, c > 0 \ni$

$$\Re a(u, u) + \lambda \|u\|_H^2 \geq c \|u\|_V^2, \quad \forall u \in V$$

then $D(\mathbb{A})$ is dense in H , \mathbb{A} is closed, $\mathbb{A}^* = \mathbb{R}$ and $D^*(\mathbb{A}) = D(\mathbb{R})$.

Closed Operators, Adjoint, Eigenfunction Expansions

Proof. Theorem  shows $D(A)$ is dense in H . Since the sesquilinear forms a and b are adjoints of each other, $A^* = R \Leftrightarrow R^* = A$. So if it is shown that $A^* = R$, it follows from Lemma  that $A = (A^*)^*$ is closed.


Fix $v \in D(R)$. Then $\forall u \in D(A)$,

$$(Au, v)_H = a(u, v) = r(v, u) = \overline{(Rv, u)_H} = (u, Rv)_H$$

so $D(R) \leq D^*(A)$ and $A^*|_{D(R)} = R$. To show $D^*(A) \leq D(R)$, let $u \in D^*(A)$.

To show that $(A + \lambda)$ is surjective, let $f \in H$ and define

$$b(v) = (f, v)_H, \quad v \in H.$$

Then b and a are continuous on V , and because of the assumed coercivity, it follows from Theorem , $\exists! w \in V \ni$

$$a(w, v) + \lambda(w, v)_H = b(v), \quad \forall v \in V$$

and

$$|a(w, v) + \lambda(w, v)| = |b(v)| \leq \|f\|_H \|v\|_H$$

Thus $w \in D(A + \lambda) = D(A)$ and

$$((A + \lambda)w, v)_H = a(w, v) + \lambda(w, v)_H = (f, v)_H, \quad \forall v \in V$$

Closed Operators, Adjoins, Eigenfunction Expansions

Since V is dense in H , $(A + \lambda)w = f$. Therefore, $A + \lambda$ and similarly $R + \lambda$ are surjective. In particular, there is a $u_0 \in D(R)$ such that

$$(R + \lambda)u_0 = (A^* + \lambda)u.$$

Then $\forall v \in D(A)$,

$$\begin{aligned} ((A + \lambda)v, u)_H &= (v, (A^* + \lambda)u)_H = (v, (R + \lambda)u_0)_H \\ &= a(v, u_0) + \lambda(v, u_0)_H = ((A + \lambda)v, u_0)_H. \end{aligned}$$

Since $A + \lambda$ is surjective, $u = u_0 \in D(R)$. Hence, $D^*(A) = D(R)$. ■

Theorem: Let V and H be Hilbert spaces with V dense in H and assume the injection $V \hookrightarrow H$ is compact. Let a be a sesquilinear form continuous, elliptic and symmetric on V ,

$$a(u, v) = a(v, u) = \overline{a(u, v)}, \quad u, v \in V$$

Let $A \in L(D(A), H)$ be defined by

$$a(u, v) = (Au, v)_H, \quad \forall u \in D(A), \quad \forall v \in V$$

where

$$D(A) = \{u \in V : \exists K > 0 \ni |a(u, v)| \leq K\|v\|_H, \forall v \in V\}.$$

Closed Operators, Adjoint, Eigenfunction Expansions

Then $\exists \{v_j\}$ eigenfunctions of A with

$$Av_j = \lambda_j v_j, \quad (v_i, v_j)_H = \delta_{ij}, \quad 0 < \lambda_1 \leq \dots \leq \lambda_n \xrightarrow{n \rightarrow \infty} +\infty$$

and $\{v_j\}$ is a basis for H .

Proof. By Theorem [▶](#), $D(A)$ is dense in H . To define A^{-1} let $f \in H$ and define $b(v) = (f, v)_H$ for $v \in V$. Then b is continuous on V and a is continuous and coercive on $V \times V$. Thus, by Theorem [▶](#), $\exists! u \in V \ni$

$$a(u, v) = b(v), \quad \forall v \in V$$

and

$$|a(u, v)| = |b(v)| \leq \|f\|_H \|v\|_H$$

Thus $u \in D(A)$ and

$$(Au, v)_H = a(u, v) = (f, v)_H, \quad \forall v \in V$$

Since V is dense in H , $Au = f$. Therefore, A is surjective. For $f \in H$, set $u = A^{-1}f \in D(A)$. By V ellipticity and $V \hookrightarrow H$,

$$\|f\|_H \|A^{-1}f\|_H \geq (Au, u)_H = a(u, u) \geq c_0 \|u\|_V^2 \geq c_1 \|A^{-1}f\|_H^2$$

Since $f \in H$ is arbitrary, $A^{-1} \in \mathcal{L}(H)$.

Closed Operators, Adjoins, Eigenfunction Expansions

From the symmetry of a and Theorem 10.1 it follows that $\mathbb{A} = \mathbb{A}^*$.

For $x, y \in H$, $u = \mathbb{A}^{-1}x$, $v = \mathbb{A}^{-1}y$ satisfy $u, v \in D(\mathbb{A})$ and so

$$(\mathbb{A}^{-1}x, y)_H = (u, \mathbb{A}v)_H = (\mathbb{A}u, v)_H = (x, \mathbb{A}^{-1}y)_H, \quad \forall x, y \in H$$

and thus \mathbb{A}^{-1} is self-adjoint. For $\{f_n\} \subset H$, $u_n = \mathbb{A}^{-1}f_n \in D(\mathbb{A})$, $c\|f_n\|_H\|u_n\|_V \geq \|f_n\|_H\|u_n\|_H \geq (\mathbb{A}u_n, u_n)_H = a(u_n, u_n) \geq c_0\|u_n\|_V^2$ where $V \hookrightarrow H$ has been used. So if $\{f_n\}$ is bounded in H , $\{u_n\}$ is bounded in V . Since the injection $V \hookrightarrow H$ is compact, $\{u_n\}$ has a convergent subsequence in H , and thus \mathbb{A}^{-1} is compact.

Applying Theorem 10.1 to \mathbb{A}^{-1} gives a sequence $\{v_j\}$ of eigenfunctions orthonormal in H such that

$\text{Rg}(\mathbb{A}^{-1}) = D(\mathbb{A}) \subset \overline{\langle v_j \rangle}$ or $H = \overline{D(\mathbb{A})} \subset \overline{\langle v_j \rangle} \subset H$, i.e., $\overline{\langle v_j \rangle} = H$.

Also, the corresponding eigenvalues $\{\mu_j\}$ satisfy

$|\mu_j| \geq |\mu_{j+1}| \xrightarrow{j \rightarrow \infty} 0$. Since a is symmetric,

$$\begin{aligned} \frac{\|v_j\|_H^2}{\mu_j} &= (\mathbb{A}v_j, v_j)_H = a(v_j, v_j) \\ &= \overline{a(v_j, v_j)} = \overline{(\mathbb{A}v_j, v_j)_H} = (v_j, \mathbb{A}v_j)_H = \|v_j\|_H^2 / \bar{\mu}_j \end{aligned}$$

it follows that $\mu_j = \bar{\mu}_j$.

Closed Operators, Adjoint, Eigenfunction Expansions

Since a is V elliptic,

$$\|v_j\|_H^2 / \mu_j = (\mathbb{A}v_j, v_j)_H = a(v_j, v_j) \geq c \|v_j\|_V^2 > 0$$

it follows that $\mu_j > 0$. Then the eigenvalues $\lambda_j = 1/\mu_j$ for \mathbb{A} satisfy $0 < \lambda_1 \leq \dots \leq \lambda_n \xrightarrow{n \rightarrow \infty} +\infty$. ■

Corollary: Let the assumption in the previous theorem that a is V elliptic be replaced by the condition that

$$a(v, v) + \lambda \|v\|_H^2 \geq c \|v\|_V, \quad \forall v \in V$$

for some $\lambda \in \mathbb{R}$ and $c > 0$. Then there is an orthonormal sequence of eigenfunctions of \mathbb{A} which is a basis for H and the corresponding eigenvalues satisfy $-\lambda < \lambda_1 \leq \dots \leq \lambda_n \xrightarrow{n \rightarrow \infty} \infty$.

Example: Take $H = L^2(G)$ and $G = (0, 1)$. Let $V = H_0^1(G)$ and

$$a(u, v) = \int_0^1 DuD\bar{v}$$

The compactness of $V \hookrightarrow H$ follows from Theorem [▶](#). Then Theorem [▶](#) shows a is $H_0^1(G)$ elliptic. Thus Theorem [▶](#) applies. The eigenfunctions and corresponding eigenvalues for

Closed Operators, Adjoint, Eigenfunction Expansions

$\mathbb{A} = -D^2$ with domain $D(\mathbb{A}) = H_0^1(G) \cap H^2(G)$ are:

$$\lambda = (j\pi)^2, \quad v_j(x) = 2 \sin(j\pi x), \quad j = 1, 2, \dots$$

Since $\{v_j\}$ is a basis for $L^2(G)$, each $f \in L^2(G)$ has a Fourier sine-series expansion. Similarly for $G = (0, 1)^n \subset \mathbb{R}^n$.

Example: As above but now let $V = H^1(G)$. The compactness of $V \hookrightarrow H$ follows from Theorem [▶](#). So Corollary [▶](#) applies for any $\lambda > 0$. The eigenfunctions and corresponding eigenvalues for $\mathbb{A} = -D^2$ with $D(\mathbb{A}) = \{v \in H^2(G) : v'(0) = v'(1) = 0\}$ are:

$$v_0(x) = 1, \quad v_j(x) = 2 \cos(j\pi x), \quad j \geq 1, \quad \lambda_j = (j\pi)^2, \quad j \geq 0.$$

Similarly for $G = (0, 1)^n \subset \mathbb{R}^n$.

Example: As above but now let

$V = \{v \in H^1(G) : v(0) = v(1)\}$. The compactness of $V \hookrightarrow H$ follows from Theorem [▶](#). So Corollary [▶](#) applies for some $\lambda > 0$. The eigenfunction expansion for $\mathbb{A} = -D^2$ with $D(\mathbb{A}) = \{v \in H^2(G) : v(0) = v(1), v'(0) = v'(1)\}$ is just the standard Fourier series.

Introduction to Evolution Equations

Consider the model problem for $G = (0, \pi)$,

$$\begin{cases} u_t = u_{xx} = \mathbb{A}u, & x \in G, & t > 0 \\ u = 0, & x \in \partial G, & t > 0 \\ u = u_0, & x \in G, & t = 0 \end{cases}$$

The solution is

$$u(x, t) = S(t)u_0(x) = \sum_{k=0}^{\infty} (u_0, v_k) v_k e^{\lambda_k t}$$

where $\{v_k\}$ are orthonormal eigenfunctions in $H = L^2(G)$ and $\{\lambda_k\}$ are the corresponding eigenvalues of \mathbb{A} .

Also the solution operator $S(t)$ is roughly $\exp(\mathbb{A}t)$ and satisfies the following properties:

Def: A contraction semigroup on H is a family of operators $\{S(t)\}_{t \geq 0} \subset \mathcal{L}(H)$ satisfying:

- $\|S(t)\|_H \leq 1$,
- $S(t + \tau) = S(t)S(\tau)$, $t, \tau \geq 0$,
- $S(0) = I$,
- $S(t)u \in C([0, \infty), H)$, $\forall u \in H$.

Introduction to Evolution Equations

Differentiation gives formally $D_t S(t) = D_t \exp(\mathbb{A}t) \xrightarrow{t \rightarrow 0} \mathbb{A}$:

Def: The *generator* of a contraction semigroup is an operator \mathbb{A} with domain

$$D(\mathbb{A}) = \{u \in H : D_t S(t)u|_{t=0} \text{ exists in } H\}$$

and value $\mathbb{A}u = D_t S(t)u|_{t=0}$. ■

For $S(t) = \exp(\mathbb{A}t)$ to exist for $t > 0$, the generator \mathbb{A} should be a negative operator:

Def: An operator $\mathbb{A} \in L(D(\mathbb{A}), H)$ is *dissipative* if $\Re(\mathbb{A}u, u)_H \leq 0$, $\forall u \in D(\mathbb{A})$.

Theorem: (Lumer-Phillips) Given a Hilbert space H and $D(\mathbb{A}) \leq H$, an operator $\mathbb{A} \in L(D(\mathbb{A}), H)$ generates a contraction semigroup if and only if

- ▶ \mathbb{A} is dissipative,
- ▶ $\mathbb{A} - \lambda I$ is surjective for $\forall \lambda > 0$.

Introduction to Evolution Equations

Def: A *classical* solution to the Cauchy problem is a function $u \in C([0, \infty), H) \cap C^1((0, \infty), H)$ satisfying

$$u'(t) = \mathbb{A}u(t), \quad u(0) = u_0$$

as well as $u(t) \in D(\mathbb{A}), \forall t > 0$.

Def: A function $u \in C([0, \infty), H)$ is a *mild* solution to the Cauchy problem if

$$\int_0^t u(s) ds \in D(\mathbb{A}) \quad \text{and} \quad \mathbb{A} \int_0^t u(s) ds = u(t) - u_0.$$

Theorem: Given a Hilbert space H and $D(\mathbb{A}) \leq H$, an operator $\mathbb{A} \in L(D(\mathbb{A}), H)$ generates a contraction semigroup $S(t)$ if and only if for all $\forall u_0 \in H$ there exists a unique mild solution $u(t)$ to the Cauchy problem, and $u(t) = S(t)u_0$.

Example: For the model problem define $H = L^2(G)$, $V = H_0^1(G)$ and

$$a(u, v) = \int_G Du D\bar{v}, \quad u, v \in V$$

Then take

$$D(\mathbb{A}) = \{u \in V : \exists K > 0 \ni |a(u, v)| \leq K \|v\|_H, \forall v \in V\}$$

and $\mathbb{A} \in L(D(\mathbb{A}), H)$ defined by

Introduction to Evolution Equations

$$-a(u, v) = (\mathbb{A}u, v)_H, \quad \forall u \in D(\mathbb{A}), \quad \forall v \in V.$$

It follows immediately that \mathbb{A} is dissipative.

To show the surjectivity condition above with $\lambda > 0$, let $f \in H$ and define

$$b(v) = \int_G f \bar{v}, \quad v \in V$$

By Theorem [▶](#), $\exists! u \in V \ni$

$$a(u, v) + \lambda(u, v)_H = b(v), \quad \forall v \in V$$

By Theorem [▶](#), $u \in D(\mathbb{A})$ and hence

$$-(\mathbb{A}u, v)_H + \lambda(u, v)_H = a(u, v) + \lambda(u, v)_H = b(v), \quad \forall v \in V$$

Since V is dense in H , it follows that $-\mathbb{A}u + \lambda u = f$ in H .

By Theorem [▶](#), \mathbb{A} generates a contraction semigroup $S(t)$.

By Theorem [▶](#), $u(t) = S(t)u_0$ is a mild solution to the Cauchy problem, $u'(t) = \mathbb{A}u(t)$, $u(0) = u_0$.

Study Question: Obtain a mild solution for the wave equation,

$$\begin{cases} u_{tt} = u_{xx}, & x \in G, & t > 0 \\ u = 0, & x \in \partial G, & t > 0 \\ u = u_0, & x \in G, & t = 0 \\ u_t = u_1, & x \in G, & t = 0 \end{cases}$$