

# WHEN DOES A QUOTIENT RING OF A PID HAVE THE CANCELLATION PROPERTY?

GYU WHAN CHANG AND JUN SEOK OH

ABSTRACT. An ideal  $I$  of a commutative ring is called a cancellation ideal if  $IB = IC$  implies  $B = C$  for all ideals  $B$  and  $C$ . Let  $D$  be a principal ideal domain (PID),  $a, b \in D$  be nonzero elements with  $a \nmid b$ ,  $(a, b)D = dD$  for some  $d \in D$ ,  $D_a = D/aD$  be the quotient ring of  $D$  modulo  $aD$ , and  $bD_a = (a, b)D/aD$ ; so  $bD_a$  is a nonzero commutative ring. In this paper, we show that the following three properties are equivalent: (i)  $\frac{a}{d}$  is a prime element and  $a \nmid d^2$ , (ii) every nonzero ideal of  $bD_a$  is a cancellation ideal, and (iii)  $bD_a$  is a field.

## 1. INTRODUCTION

Let  $S$  be a commutative semigroup under multiplication. The zero element of  $S$  (if it exists) is an element  $0 \in S$  such that  $a \cdot 0 = 0 \cdot a = 0$  for all  $a \in S$ . An element  $a \in S$  is said to be *cancellative* if  $ab = ac$  implies  $b = c$  for all  $b, c \in S$ . Clearly if  $S$  has an identity, then every invertible element of  $S$  is cancellative, so the cancellation property is a natural generalization of invertibility. We say that  $S$  is *cancellative* if every nonzero element of  $S$  is cancellative. Let  $S^\bullet = S \setminus \{0\}$ . Then  $S^\bullet$  is not a semigroup in general, while  $S$  is cancellative if and only if  $S^\bullet$  is a cancellative semigroup. The cancellation property plays an important role for the study on algebra. For example, assume that  $S^\bullet$  is a semigroup. Then (i)  $S$  is cancellative if and only if  $S^\bullet$  can be embedded in a group (i.e.,  $S^\bullet$  has a quotient group) [5, Theorem 1.2], (ii)  $S$  is torsion-free and cancellative if and only if  $S^\bullet$  admits a total order compatible with its semigroup operation [5, Corollary 3.4], and (iii) several kinds of factorization properties of a semigroup (e.g., atomic, factorial, half-factorial, bounded factorization) have been studied under the assumption that it is cancellative (see [3] for a survey).

Let  $R$  be a commutative ring (not necessarily having an identity),  $\mathcal{I}(R)$  be the set of ideals of  $R$ , and  $\mathcal{P}(R)$  be the set of principal ideals of  $R$ . Then  $\mathcal{I}(R)$  becomes a commutative semigroup with zero element under the usual ideal multiplication,  $\mathcal{P}(R)$  is a subsemigroup of  $\mathcal{I}(R)$ , and if  $R$  has an identity, then  $\mathcal{I}(R)$  has an identity. We say that an ideal  $I$  of  $R$  is a *cancellation ideal* if  $I$  is cancellative as an element of  $\mathcal{I}(R)$ . It is easy to see that a principal ideal  $(a)$  of  $R$  generated by  $a \in R$  is a cancellation ideal if and only if  $a$  is a regular element of  $R$  (i.e.,  $a$  is not a zero-divisor). Furthermore, if  $R$  has an identity, then a nonzero ideal  $I$  of  $R$  is a cancellation ideal if and only if  $IR_M$  is a regular principal ideal for all maximal ideals  $M$  of  $R$  [1, Theorem, p. 2853], and  $\mathcal{P}(R)$  is cancellative if and only if  $R$  is an integral domain, if and only if  $\mathcal{I}(R)^\bullet$  is a semigroup. It is well known that if  $R$  is an integral domain, then (i)  $\mathcal{P}(R)$  is factorial if and only if  $R$  is a unique factorization

---

*Date:* April 16, 2021.

2010 Mathematics Subject Classification: 13A15.

Key words and Phrases: principal ideal domain, cancellation ideals.

domain, (ii)  $\mathcal{I}(R)$  is factorial if and only if  $R$  is a Dedekind domain [4, Theorem 37.8], (iii)  $\mathcal{I}(R)$  is cancellative if and only if  $R$  is an almost Dedekind domain (i.e.,  $R_M$  is a principal ideal domain (PID) for all maximal ideals  $M$  of  $R$ ) [4, Theorem 36.5], and (iv) every nonzero finitely generated ideal of  $R$  is a cancellation ideal if and only if  $R$  is a Prüfer domain, (i.e., every nonzero finitely generated ideal of  $R$  is invertible) [4, Theorem 24.3].

Now let  $\mathbb{Z}$  be the ring of integers,  $m$  and  $n$  be positive integers,  $\gcd(m, n)$  denote the greatest common divisor of  $m$  and  $n$ ,  $\mathbb{Z}_n$  be the ring of integers modulo  $n$ , and  $m\mathbb{Z}_n$  be the ideal of  $\mathbb{Z}_n$  generated by  $m$ ; so  $m\mathbb{Z}_n$  is a commutative ring. Then  $\mathcal{I}(\mathbb{Z}_n) = \mathcal{P}(\mathbb{Z}_n)$ , and hence  $\mathcal{I}(\mathbb{Z}_n)$  is cancellative if and only if  $\mathbb{Z}_n$  is an integral domain, if and only if either  $n = 1$  or  $n$  is a prime number. Moreover, in [2, Theorem 2.5], the authors showed that if  $n \nmid m$ , then every nonzero ideal in  $m\mathbb{Z}_n$  is a cancellation ideal, i.e.,  $\mathcal{I}(m\mathbb{Z}_n)$  is a cancellative semigroup, if and only if  $\frac{n}{\gcd(n, m)}$  is a prime number and  $n \nmid \gcd(n, m)^2$ .

Let  $D$  be a PID,  $a$  and  $b$  be nonzero elements of  $D$ , and  $d \in D$  be such that  $(a, b)D = dD$ .

1.  $D_a = D/aD$  is the quotient ring of  $D$  modulo  $aD$ .
2.  $[a, b] = d$  is the greatest common divisor of  $a$  and  $b$ .
3.  $bD_a = (a, b)D/aD$ .

Then  $bD_a$  is a commutative ring,  $[a, b]$  is determined only up to units, and if  $[a, b] = d$ , then  $[a/d, b/d] = 1$  and  $bD_a = dD_a$ . In this paper, we show that every nonzero ideal of  $bD_a$  is a cancellation ideal if and only if  $\frac{a}{[a, b]}$  is a prime element and  $a \nmid [a, b]^2$ . This result is applied in two special cases, i.e., the ring of integers and the polynomial ring over a field, and the former case recovers the result of [2, Theorem 2.5].

## 2. RESULTS

Let  $R$  be a commutative ring with identity. Then two ideals  $I, J$  of  $R$  are said to be *comaximal* if  $I + J = R$ , and we say that two elements  $a, b$  of  $R$  are *comaximal* if the principal ideals  $aR$  and  $bR$  are comaximal. Clearly,  $a, b$  are comaximal if and only if  $ar + bs = 1$  for some  $r, s \in R$ .

**Lemma 1.** *Let  $D$  be a PID and  $a, b \in D$  be nonzero elements. Then  $bD_a$  has an identity if and only if  $b$  and  $\frac{a}{[a, b]}$  are comaximal.*

*Proof.* Let  $d = [a, b]$ ,  $a_1 = \frac{a}{d}$  and  $b_1 = \frac{b}{d}$ .

( $\Rightarrow$ ) Let  $bx + aD$  be the identity of  $bD_a$  for some  $x \in D$ . Then

$$(bx + aD)(b + aD) = b + aD,$$

whence  $a \mid b(bx - 1)$ . Also,  $[a_1, b_1] = 1$  implies  $a_1 \mid bx - 1$ , and hence  $bx + a_1y = 1$  for some  $y \in D$ . Thus,  $b$  and  $a_1$  are comaximal.

( $\Leftarrow$ ) By assumption,  $bx + a_1y = 1$  for some  $x, y \in D$ , and hence  $bx = 1 - a_1y$ . So, for every  $z \in D$ , we have

$$\begin{aligned} (bx + aD)(bz + aD) &= (1 - a_1y)(bz) + aD = (bz - a_1byz) + aD \\ &= (bz + aD) - (a_1byz + aD) \\ &= bz + aD. \end{aligned}$$

Thus,  $bx + aD$  is the identity of  $bD_a$ . □

We now give the main result of this paper.

**Theorem 2.** *Let  $D$  be a PID and  $a, b \in D$  be nonzero elements with  $a \nmid b$ . Then the following statements are equivalent:*

1.  $\frac{a}{[a,b]}$  is a prime element and  $a \nmid [a,b]^2$ .
2.  $bD_a$  is a field.
3. Every nonzero ideal of  $bD_a$  is a cancellation ideal.

*Proof.* Let  $d = [a, b]$ ,  $a_1 = \frac{a}{d}$ , and  $b_1 = \frac{b}{d}$ . Clearly,  $bD_a = dD_a$ ,  $(a_1, b_1)D = D$ ,  $a_1$  is a nonunit, and  $bD_a \neq (0)$  because  $a \nmid b$ .

(1)  $\Rightarrow$  (2) Note that  $bD_a$  is a commutative ring; so  $bD_a$  is a field if and only if  $bD_a$  has an identity and  $bD_a$  does not have a proper nonzero ideal.

We first show that  $bD_a$  has an identity. Note that  $a = da_1$  and  $b = db_1$ ; so  $a \nmid d^2$  implies  $a_1 \nmid d$ . Hence,  $a_1 \nmid b$  because  $[a_1, b_1] = 1$  and  $b = db_1$ . Since  $a_1 = \frac{a}{d}$  is a prime element by assumption, it follows that  $a_1$  and  $b$  are comaximal. Hence, by Lemma 1,  $bD_a$  has an identity. Next, let  $A$  be an ideal of  $bD_a$ . Then  $D_a A = D_a(bD_a A) = bD_a A = A$  because  $bD_a$  has an identity. Thus,  $A$  is an ideal of  $D_a$ , so there exists  $e \in D$  such that

$$aD \subseteq eD \subseteq dD \quad \text{and} \quad A = eD/aD \subseteq bD_a.$$

Hence,  $e = dx$  and  $a = ey$  for some  $x, y \in D$ , and thus  $a = dxy$  or  $a_1 = xy$ . By assumption,  $a_1 = xy$  is a prime element of  $D$ , whence either  $x$  or  $y$  is a unit of  $D$ . If  $x$  is a unit, then  $eD = dD$ , and hence  $A = eD/aD = dD_a = bD_a$ . If  $y$  is a unit, then  $eD = aD$ , whence  $A = eD/aD = aD/aD$  is the zero ideal of  $bD_a$ . Therefore,  $bD_a$  does not have a proper nonzero ideal.

(2)  $\Rightarrow$  (3) Clear.

(3)  $\Rightarrow$  (1) If  $a \mid d^2$ , then  $(dD_a)^2 = d^2D_a = (0)$ , and since  $dD_a$  is a cancellation ideal in  $bD_a$ ,  $dD_a = (0)$ , a contradiction. Thus,  $a \nmid d^2$ .

Next, assume to the contrary that  $a_1 = pq$  for some nonunit elements  $p, q$  of  $D$ . Let  $I = pdD_a$  and  $J = qdD_a$ . Then  $I$  and  $J$  are ideals of  $dD_a$ . If  $I = (0)$ , then  $pd + aD = aD$ , and hence  $a \mid pd$ . Note that  $a = a_1d$ ; so  $a_1 \mid p$ , and thus  $q$  is a unit, a contradiction. Similarly, we have  $J \neq (0)$ . However,

$$IJ = pdqdD_a = a_1d^2D_a = adD_a = (0).$$

Thus,  $I$  and  $J$  are not cancellation ideals, a contradiction.  $\square$

As a corollary of Theorem 2, we have.

**Corollary 3.** *Let  $D$  be a PID and  $a \in D$  be a nonzero element. Then every nonzero ideal in  $D_a$  is a cancellation ideal if and only if  $a$  is a unit or  $a$  is a prime element.*

We have two applications of Theorem 2: one is to the ring of integers and the other is to the polynomial ring over a field.

**Corollary 4.** *Let  $n, m \in \mathbb{Z}$  be positive integers with  $n \nmid m$ .*

1. ([2, Theorem 2.5]) *Every nonzero ideal in  $m\mathbb{Z}_n$  is a cancellation ideal if and only if  $\frac{n}{\gcd(n,m)}$  is a prime number and  $n \nmid \gcd(n,m)^2$ .*
2. ([2, Corollary 2.6]) *Every nonzero ideal in  $\mathbb{Z}_n$  is a cancellation ideal if and only if either  $n = 1$  or  $n$  is a prime number.*

**Corollary 5.** *Let  $F$  be a field,  $X$  be an indeterminate over  $F$ ,  $F[X]$  be the polynomial ring over  $F$ ,  $f, g \in F[X]$  be nonzero polynomials with  $f \nmid g$  in  $F[X]$ , and  $(f, g)F[X] = hF[X]$  for some  $h \in F[X]$ .*

1. Every nonzero ideal in  $(f, g)F[X]/fF[X]$  is a cancellation ideal if and only if  $\frac{f}{g}$  is irreducible over  $F$  and  $f \nmid h^2$  in  $F[X]$ .
2. Every nonzero ideal in  $F[X]/fF[X]$  is a cancellation ideal if and only if either  $f \in F$  or  $f$  is irreducible over  $F$ .

*Proof.* This follows directly from Theorem 2 and Corollary 3, because an irreducible polynomial of  $F[X]$  is a prime element.  $\square$

#### REFERENCES

- [1] D. D. Anderson and M. Roitman, *A characterization of cancellation ideals*, Proc. Amer. Math. Soc. **125** (1997), 2853–2854.
- [2] S. Chaopraknoi, K. Savettaseranee and P. Lertwichitsilp, *Some cancellation ideal rings*, General Mathematics **13** (2005), 39–46.
- [3] A. Geroldinger and Q. Zhong, *Factorization theory in commutative monoids*, Semigroup Forum **100** (2020), 22–51.
- [4] R. Gilmer, *Multiplicative Ideal Theory*, Marcel Dekker Inc., New York, 1972.
- [5] R. Gilmer, *Commutative Semigroup Rings*, Univ. Chicago Press, Chicago, 1984.

DEPARTMENT OF MATHEMATICS EDUCATION, INCHEON NATIONAL UNIVERSITY, INCHEON 22012, REPUBLIC OF KOREA

*Email address:* whan@inu.ac.kr

DEPARTMENT OF MATHEMATICS EDUCATION, JEJU NATIONAL UNIVERSITY, JEJU 63243, REPUBLIC OF KOREA

*Email address:* junseok.oh@jejunu.ac.kr