

THE MONOTONE CATENARY DEGREE OF MONOIDS OF IDEALS

ALFRED GEROLDINGER AND ANDREAS REINHART

ABSTRACT. Factoring ideals in integral domains is a central topic in multiplicative ideal theory. In the present paper we study monoids of ideals and consider factorizations of ideals into multiplicatively irreducible ideals. The focus is on the monoid of nonzero divisorial ideals and on the monoid of v -invertible divisorial ideals in weakly Krull Mori domains. Under suitable algebraic finiteness conditions we establish arithmetical finiteness results, in particular for the monotone catenary degree and for the structure of sets of lengths and of their unions.

1. INTRODUCTION

Factoring ideals in integral domains is a central topic in multiplicative ideal theory (for a monograph reflecting recent developments we refer to [12]). In the present paper we study monoids of ideals, consider factorizations of ideals into multiplicatively irreducible ideals, and prove finiteness results on the monotone catenary degree and structural results on sets of lengths. First we recall the concept of the monotone catenary degree and then we discuss the monoids of ideals under consideration.

Let H be an atomic monoid which is not factorial. Then every non-unit can be written as a finite product of atoms and there is an element $a \in H$ having at least two distinct factorizations, say $a = u_1 \cdots u_\ell v_1 \cdots v_m = u_1 \cdots u_\ell w_1 \cdots w_n$ where all u_i, v_j, w_k are atoms and the v_j and w_k are pairwise not associated. Then $\ell + m$ and $\ell + n$ are the lengths of the two factorizations and $\max\{m, n\} > 0$ is their distance. Then for every $M \in \mathbb{N}$, the element

$$a^M = (u_1 \cdots u_\ell v_1 \cdots v_m)^\nu (u_1 \cdots u_\ell w_1 \cdots w_n)^{M-\nu} \quad \text{for all } \nu \in [0, M]$$

has factorizations with distance greater than M . However, at least those factorizations, which are powers of factorizations of a , can be concatenated, step by step, by factorizations whose distance is small and does not depend on M . This phenomenon is formalized by the catenary degree which is defined as follows. The catenary degree $\mathfrak{c}(H)$ of H is the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ such that for each $a \in H$ and each two factorizations z, z' of a there is a concatenating chain of factorizations $z = z_0, z_1, \dots, z_{k+1} = z'$ of a such that the distance $\mathfrak{d}(z_{i-1}, z_i)$ between two successive factorizations is bounded by N . It is well-known that the catenary degree is finite for Krull monoids with finite class group and for C-monoids (these include Mori domains R with nonzero conductor $\mathfrak{f} = (R : \widehat{R})$ for which the residue class ring \widehat{R}/\mathfrak{f} and the class group $\mathcal{C}(\widehat{R})$ are finite).

In order to study further structural properties of concatenating chains, Foroutan introduced the monotone catenary degree ([13]). The monotone catenary degree $\mathfrak{c}_{\text{mon}}(H)$ is the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ such that for each $a \in H$ and each two factorizations z, z' of a there is a concatenating chain of factorizations $z = z_0, z_1, \dots, z_{k+1} = z'$ of a such that the distance between two successive factorizations is bounded by N and in addition the sequence of lengths $|z_0|, \dots, |z_{k+1}|$ of the factorizations is monotone (thus either $|z_0| \leq \dots \leq |z_{k+1}|$ or $|z_0| \geq \dots \geq |z_{k+1}|$). Therefore, by definition, we have $\mathfrak{c}(H) \leq \mathfrak{c}_{\text{mon}}(H)$ and

2010 *Mathematics Subject Classification.* 13A15, 13F15, 20M12, 20M13.

Key words and phrases. ideal systems, Cohen-Kaplansky domains, generalized Cohen-Kaplansky domains, weakly Krull domains, monotone catenary degree, sets of lengths.

This work was supported by the Austrian Science Fund FWF, Project Number P26036-N26.

Foroutan showed that the monotone catenary degree of Krull monoids with finite class group is finite again. Subsequently the monotone catenary degree, or more generally, monotonic properties of concatenating chains were studied in a variety of papers (e.g., [7, 15, 18, 23, 28, 29]). We mention one result in detail, namely that C-monoids have the following property ([14, Theorem 1.1]):

There exists some constant $M \in \mathbb{N}$ such that for every $a \in H$ and for each two factorizations z, z' of a there exist factorizations $z = z_0, z_1, \dots, z_{k+1} = z'$ of a such that, for every $i \in [1, k+1]$,

$$d(z_{i-1}, z_i) \leq M \quad \text{and} \quad (\text{either } |z_1| \leq \dots \leq |z_k| \text{ or } |z_1| \geq \dots \geq |z_k|).$$

Thus the sequence of lengths of factorizations is monotone apart from the first and the last step. However, there are C-monoids having infinite monotone catenary degree and this may happen even for one-dimensional local Noetherian domains (see Remark 5.3).

In Section 4 we study integral domains R with complete integral closure \widehat{R} , nonzero conductor $(R: \widehat{R})$, and with an r -Noetherian ideal system r such that $r\text{-max}(R) = \mathfrak{X}(R)$. Under suitable algebraic finiteness conditions the monoid of nonzero r -ideals $\mathcal{I}_r(R)$ is finitely generated up to a free abelian factor. However, since the monoid need not be cancellative, its study needs some semigroup theoretical preparations (done in Section 3) valid for unit-cancellative monoids. The main arithmetical result on $\mathcal{I}_r(R)$ is given in Theorem 4.15, and for its assumptions see Theorems 4.8 and 4.13.

In Section 5 we study the monoid $\mathcal{I}_v^*(H)$ of v -invertible v -ideals of v -Noetherian weakly Krull monoids with nontrivial conductor. This is a C-monoid and isomorphic to a finite direct product of finitely primary monoids and free abelian factor. However, shifting the finiteness of the monotone catenary degree to finite direct products might not work as it should (see Example 3.3). In order to overcome these difficulties we introduce a new arithmetical invariant, the weak successive distance (Definition 3.4), and establish a result allowing to shift the finiteness of the monotone catenary degree to finite direct products (Theorem 3.8). This is done under the additional assumption that the Structure Theorem for Sets of Lengths holds (which is the case for $\mathcal{I}_v^*(H)$). Along our way we prove that the Structure Theorem for Unions holds, both for the monoid $\mathcal{I}_r(R)$ (studied in Section 4) as well as for $\mathcal{I}_v^*(H)$. This is based on a recent characterization result for the validity of the Structure Theorem for Unions obtained in [10]. The main results on $\mathcal{I}_v^*(H)$ are given in Theorem 5.13 and Corollary 5.14.

2. BACKGROUND ON THE ARITHMETIC OF MONOIDS

We denote by \mathbb{N} the set of positive integers, and we put $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For $s \in \mathbb{N}$, we will consider the product order on \mathbb{N}_0^s which is induced by the usual order of \mathbb{N}_0 . For real numbers $a, b \in \mathbb{R}$, we set $[a, b] = \{x \in \mathbb{Z} \mid a \leq x \leq b\}$. Let $L, L' \subset \mathbb{Z}$. We denote by $L + L' = \{a + b \mid a \in L, b \in L'\}$ their *sumset*. Two distinct elements $k, \ell \in L$ are called *adjacent* if $L \cap [\min\{k, \ell\}, \max\{k, \ell\}] = \{k, \ell\}$. A positive integer $d \in \mathbb{N}$ is called a *distance* of L if there exist adjacent elements $k, \ell \in L$ with $d = |\ell - k|$, and we denote by $\Delta(L)$ the *set of distances* of L . If $L \subset \mathbb{N}$, we denote by $\rho(L) = \sup L / \min L \in \mathbb{Q}_{\geq 1} \cup \{\infty\}$ the *elasticity* of L . We set $\rho(\{0\}) = 1$ and $\max \emptyset = \min \emptyset = \sup \emptyset = 0$.

Monoids and factorizations. All rings and semigroups are commutative and have an identity element. Let H be a semigroup. If not stated otherwise, we use multiplicative notation and $1 = 1_H \in H$ means the identity element of H . We denote by H^\times the group of invertible elements of H and we say that H is reduced if $H^\times = \{1\}$. Then $H_{\text{red}} = \{aH^\times \mid a \in H\}$ denotes the associated reduced semigroup of H . Furthermore, H is called

- *unit-cancellative* if $a, u \in H$ and $a = au$, implies that $u \in H^\times$,
- *cancellative* if $a, b, u \in H$ and $au = bu$ implies that $a = b$.

Obviously, every cancellative semigroup is unit-cancellative. The property of being unit-cancellative is a frequently studied property both for rings and semigroups. Indeed, a commutative ring R is called *présimplifiable* if $a, u \in R$ and $a = au$ implies that either $a = 0$ or $u \in R^\times$. This concept was introduced by Bouvier and further studied by D.D. Anderson et al. ([1, 6, 8, 16]).

If H is cancellative, then $\mathfrak{q}(H)$ denotes the quotient group of H and

$$\widehat{H} = \{x \in \mathfrak{q}(H) \mid \text{there is a } c \in H \text{ such that } cx^n \in H \text{ for every } n \in \mathbb{N}\} \subset \mathfrak{q}(H)$$

is the *complete integral closure* of H . Let R be a domain with quotient field K . For every subset $X \subset K$ we set $X^\bullet = X \setminus \{0\}$. Then R^\bullet is a cancellative semigroup, \overline{R} is the integral closure of R , and $\widehat{R} = \widehat{R^\bullet} \cup \{0\}$ is the complete integral closure of R .

Throughout this paper, a monoid will always mean a commutative unit-cancellative semigroup with identity element.

For a set \mathcal{P} , we denote by $\mathcal{F}(\mathcal{P})$ the *free abelian monoid* with basis \mathcal{P} . Then every $a \in \mathcal{F}(\mathcal{P})$ has a unique representation in the form

$$a = \prod_{p \in \mathcal{P}} p^{\nu_p(a)} \quad \text{with } \nu_p(a) \in \mathbb{N}_0 \text{ and } \nu_p(a) = 0 \text{ for almost all } p \in \mathcal{P}.$$

We call $|a| = \sum_{p \in \mathcal{P}} \nu_p(a)$ the *length* of a and $\text{supp}(a) = \{p \in \mathcal{P} \mid \nu_p(a) > 0\} \subset \mathcal{P}$ the *support* of a .

Let H be a monoid. A submonoid $S \subset H$ is said to be

- *divisor-closed* if $a \in S$ and $b \in H$ with $b \mid a$ implies that $b \in S$,
- *saturated* if $a, c \in S$ and $b \in H$ with $a = bc$ implies that $b \in S$.

Clearly, every divisor-closed submonoid is saturated. An element $u \in H$ is said to be *irreducible* (or an *atom*) if $u \notin H^\times$ and an equation $u = ab$ with $a, b \in H$ implies that $a \in H^\times$ or $b \in H^\times$. Then $\mathcal{A}(H)$ denotes the set of atoms of H , and H is said to be *atomic* if every non-unit can be written as a finite product of atoms of H . A simple argument shows that H is atomic whenever the ACCP (ascending chain condition on principal ideals) holds ([10, Lemma 3.1]). From now on we suppose that H is atomic.

The free abelian monoid $\mathbf{Z}(H) = \mathcal{F}(\mathcal{A}(H_{\text{red}}))$ is called the *factorization monoid* of H , and the homomorphism

$$\pi: \mathbf{Z}(H) \rightarrow H_{\text{red}} \quad \text{satisfying } \pi(u) = u \text{ for each } u \in \mathcal{A}(H_{\text{red}})$$

is called the *factorization homomorphism* of H . For $a \in H$ and $k \in \mathbb{N}$,

$$\mathbf{Z}_H(a) = \mathbf{Z}(a) = \pi^{-1}(aH^\times) \subset \mathbf{Z}(H) \quad \text{is the set of factorizations of } a,$$

$$\mathbf{Z}_{H,k}(a) = \mathbf{Z}_k(a) = \{z \in \mathbf{Z}(a) \mid |z| = k\} \quad \text{is the set of factorizations of } a \text{ of length } k, \quad \text{and}$$

$$\mathbf{L}_H(a) = \mathbf{L}(a) = \{|z| \mid z \in \mathbf{Z}(a)\} \subset \mathbb{N}_0 \quad \text{is the set of lengths of } a.$$

If $S \subset H$ is a divisor-closed submonoid and $a \in S$, then $\mathbf{Z}_S(a) = \mathbf{Z}_H(a)$ whence $\mathbf{L}_S(a) = \mathbf{L}_H(a)$. Then

- $\mathcal{L}(H) = \{\mathbf{L}(a) \mid a \in H\}$ is the *system of sets of lengths* of H ,
- $\Delta(H) = \bigcup_{L \in \mathcal{L}(H)} \Delta(L)$ is the *set of distances* of H , and
- $\rho(H) = \sup\{\rho(L) \mid L \in \mathcal{L}(H)\}$ is the *elasticity* of H .

The monoid H is said to be

- *half-factorial* if $\Delta(H) = \emptyset$ (equivalently, $|\mathbf{L}(a)| = 1$ for all $a \in H$),
- an *FF-monoid* if $\mathbf{Z}(a)$ is finite for all $a \in H$,
- a *BF-monoid* if $\mathbf{L}(a)$ is finite for all $a \in H$.

Let $z, z' \in \mathbf{Z}(H)$. Then we can write

$$z = u_1 \cdots u_\ell v_1 \cdots v_m \quad \text{and} \quad z' = u_1 \cdots u_\ell w_1 \cdots w_n,$$

where $\ell, m, n \in \mathbb{N}_0$ and $u_1, \dots, u_\ell, v_1, \dots, v_m, w_1, \dots, w_n \in \mathcal{A}(H_{\text{red}})$ are such that

$$\{v_1, \dots, v_m\} \cap \{w_1, \dots, w_n\} = \emptyset.$$

Then $\text{gcd}(z, z') = u_1 \cdots u_\ell$, and we call

$$\mathbf{d}(z, z') = \max\{m, n\} = \max\{|z \text{gcd}(z, z')^{-1}|, |z' \text{gcd}(z, z')^{-1}|\} \in \mathbb{N}_0$$

the *distance* between z and z' . If $\pi(z) = \pi(z')$ and $z \neq z'$, then

$$(2.1) \quad 1 + ||z| - |z'|| \leq \mathbf{d}(z, z') \quad \text{resp.} \quad 2 + ||z| - |z'|| \leq \mathbf{d}(z, z') \quad \text{if } H \text{ is cancellative}$$

(see [10, Proposition 3.2] and [19, Lemma 1.6.2]). For subsets $X, Y \subset \mathbf{Z}(H)$, we set

$$\begin{aligned} \mathbf{d}(X, Y) &= \min\{\mathbf{d}(x, y) \mid x \in X, y \in Y\} \text{ and} \\ \text{Dist}(X, Y) &= \sup\{\mathbf{d}(\{x\}, Y), \mathbf{d}(X, \{y\}) \mid x \in X, y \in Y\}. \end{aligned}$$

Note that $\mathbf{d}(X, Y) = 0$ if and only if ($X \cap Y \neq \emptyset$ or $X = \emptyset$ or $Y = \emptyset$). For a factorization $z \in \mathbf{Z}(H)$, we denote by $\delta(z)$ the smallest $N \in \mathbb{N}_0$ with the following property:

If $k \in \mathbb{N}$ is such that k and $|z|$ are adjacent lengths of $\pi(z)$, then there exists some $y \in \mathbf{Z}(H)$ such that $\pi(y) = \pi(z)$, $|y| = k$ and $\mathbf{d}(z, y) \leq N$.

We call

$$\delta(H) = \sup\{\delta(z) \mid z \in \mathbf{Z}(H)\} \in \mathbb{N}_0 \cup \{\infty\}$$

the (*strong*) *successive distance* of H . Note that

$$\delta(H) = \sup\{\text{Dist}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \mid a \in H, k, \ell \in \mathbf{L}(a) \text{ are adjacent}\}.$$

Chains of factorizations. Let $a \in H$ and $N \in \mathbb{N}_0 \cup \{\infty\}$. A finite sequence $z_0, \dots, z_k \in \mathbf{Z}(a)$ is called a (*monotone*) *N-chain of factorizations of a* if $\mathbf{d}(z_{i-1}, z_i) \leq N$ for all $i \in [1, k]$ (and $|z_0| \leq \dots \leq |z_k|$ or $|z_0| \geq \dots \geq |z_k|$). We denote by $\mathbf{c}(a)$ (or by $\mathbf{c}_{\text{mon}}(a)$ resp.) the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ such that any two factorizations $z, z' \in \mathbf{Z}(a)$ can be concatenated by an N -chain (or by a monotone N -chain resp.). Then

$$\mathbf{c}(H) = \sup\{\mathbf{c}(b) \mid b \in H\} \in \mathbb{N}_0 \cup \{\infty\} \quad \text{and} \quad \mathbf{c}_{\text{mon}}(H) = \sup\{\mathbf{c}_{\text{mon}}(b) \mid b \in H\} \in \mathbb{N}_0 \cup \{\infty\}$$

denote the *catenary degree* and the *monotone catenary degree* of H . The monotone catenary degree is studied by using the two auxiliary notions of the equal and the adjacent catenary degrees. Let $\mathbf{c}_{\text{eq}}(a)$ denote the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ such that any two factorizations $z, z' \in \mathbf{Z}(a)$ with $|z| = |z'|$ can be concatenated by a monotone N -chain. We call

$$\mathbf{c}_{\text{eq}}(H) = \sup\{\mathbf{c}_{\text{eq}}(b) \mid b \in H\} \in \mathbb{N}_0 \cup \{\infty\}$$

the *equal catenary degree* of H . We set

$$\mathbf{c}_{\text{adj}}(a) = \sup\{\mathbf{d}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \mid k, \ell \in \mathbf{L}(a) \text{ are adjacent}\},$$

and the *adjacent catenary degree* of H is defined as

$$\mathbf{c}_{\text{adj}}(H) = \sup\{\mathbf{c}_{\text{adj}}(b) \mid b \in H\} \in \mathbb{N}_0 \cup \{\infty\}.$$

Obviously, we have $\mathbf{c}_{\text{adj}}(H) \leq \delta(H)$,

$$\mathbf{c}(a) \leq \mathbf{c}_{\text{mon}}(a) = \sup\{\mathbf{c}_{\text{eq}}(a), \mathbf{c}_{\text{adj}}(a)\} \leq \sup \mathbf{L}(a) \quad \text{for all } a \in H,$$

and hence

$$(2.2) \quad \mathbf{c}(H) \leq \mathbf{c}_{\text{mon}}(H) = \sup\{\mathbf{c}_{\text{eq}}(H), \mathbf{c}_{\text{adj}}(H)\}.$$

Note that $\mathbf{c}_{\text{adj}}(H) = 0$ if and only if H is half-factorial and if this holds, then $\mathbf{c}_{\text{eq}}(H) = \mathbf{c}(H)$. If H is not half-factorial, then (2.1) shows that $1 + \sup \Delta(H) \leq \mathbf{c}(H)$. Moreover, $\mathbf{c}_{\text{eq}}(H) = 0$ if and only if for all $a \in H$ and all $k \in \mathbf{L}(a)$ we have $|\mathbf{Z}_k(a)| = 1$. A result by Coykendall and Smith implies that for the multiplicative monoid H of non-zero elements from a domain we have $\mathbf{c}_{\text{eq}}(H) = 0$ if and only if H is factorial ([9, Corollary 2.12]).

Structure of sets of lengths. Let $d \in \mathbb{N}$, $M \in \mathbb{N}_0$ and $\{0, d\} \subset \mathcal{D} \subset [0, d]$. Then L is called an *almost arithmetical multiprogression* (AAMP for short) with *difference* d , *period* \mathcal{D} , and *bound* M , if

$$L = y + (L' \cup L^* \cup L'') \subset y + \mathcal{D} + d\mathbb{Z}$$

where

- L^* is finite and nonempty with $\min L^* = 0$ and $L^* = (\mathcal{D} + d\mathbb{Z}) \cap [0, \max L^*]$

- $L' \subset [-M, -1]$ and $L'' \subset \max L^* + [1, M]$
- $y \in \mathbb{Z}$.

Note that an AAMP is finite and nonempty. It is straightforward to prove that if $M \in \mathbb{N}_0$, $d \in \mathbb{N}$, L is an AAMP with bound M and difference d , $x \in L$ and $y \in d\mathbb{Z}$ are such that $\min L + M \leq x + y \leq \max L - M$, then $x + y \in L$.

We say that the *Structure Theorem for Sets of Lengths* holds (for the monoid H) if H is atomic and there exist some $M \in \mathbb{N}_0$ and a finite nonempty set $\Delta \subset \mathbb{N}$ such that for every $a \in H$, the set of lengths $L(a)$ is an AAMP with some difference $d \in \Delta$ and bound M . Suppose that the Structure Theorem for Sets of Lengths holds for the monoid H . Then H is a BF-monoid with finite set of distances. The monoids of ideals (studied in Section 5) satisfy the Structure Theorem for Sets of Lengths and we use this property to show that their monotone catenary degree is finite (see Theorem 3.8).

Unions of sets of lengths. For every $k \in \mathbb{N}$,

$$\mathcal{U}_k(H) = \bigcup_{L \in \mathcal{L}(H) \text{ with } k \in L} L$$

denotes the *union of all sets of lengths containing k* , provided that $H \neq H^\times$. In the extremal case where $H = H^\times$, it is convenient to set $\mathcal{U}_k(H) = \{k\}$ for all $k \in \mathbb{N}$. Furthermore,

$$\rho_k(H) = \sup \mathcal{U}_k(H) \in \mathbb{N}_0 \cup \{\infty\}$$

denotes the k -th *elasticity* of H . Unions of sets of lengths are a classic invariant in factorization theory, and the last decade has seen a renewed interest in the structure of unions (e.g., [10, 17, 33]).

We say that the *Structure Theorem for Unions* holds (for the monoid H) if there are $d \in \mathbb{N}$ and $M \in \mathbb{N}_0$ such that, for all sufficiently large $k \in \mathbb{N}$, $\mathcal{U}_k(H)$ has the form

$$\mathcal{U}_k(H) = y_k + (L'_k \cup L_k^* \cup L''_k) \subset y_k + d\mathbb{Z}$$

where L_k^* is a nonempty arithmetical progression with difference d such that $\min L_k^* = 0$, $L'_k \subset [-M, -1]$, $L''_k \subset \sup L_k^* + [1, M]$ (with the convention that $L''_k = \emptyset$ if L_k^* is infinite), and $y_k \in \mathbb{Z}$. Note, if $\mathcal{U}_k(H)$ is finite, then $\mathcal{U}_k(H)$ is an AAMP with period $\{1, d\}$ and bound M .

Suppose that $\Delta(H)$ is finite and the Structure Theorem for Unions holds. Then, by [10, Corollary 2.3 and Lemma 2.12], the parameter d in the above definition satisfies $d = \min \Delta(H)$ and we have

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{U}_k(H)|}{k} = \frac{1}{d} \left(\rho(H) - \frac{1}{\rho(H)} \right).$$

3. FINITELY GENERATED MONOIDS AND FINITE DIRECT PRODUCTS

Let H be a monoid and $\pi: Z(H) \rightarrow H_{\text{red}}$ the canonical epimorphism. Its monoid of relations, defined as

$$\sim_H = \{(x, y) \in Z(H) \times Z(H) \mid \pi(x) = \pi(y)\},$$

is a crucial tool for studying the arithmetic of H . Suppose that H is reduced, cancellative, and atomic. Then $\sim_H \subset Z(H) \times Z(H)$ is a saturated submonoid and hence a Krull monoid ([30, Lemma 11]). If, moreover, H is finitely generated, then \sim_H is finitely generated. However, this need not be true without the assumption that H is cancellative ([10, Remarks 3.11]) (although, by Redei's Theorem, $\ker(\pi)$ is finitely generated as a congruence and H is finitely presented). A further striking difference between the cancellative case and the non-cancellative case is, that cancellative finitely generated monoids have accepted elasticity which need not be the case in the noncancellative setting (for an example we refer again to [10, Remarks 3.11]).

In spite of all these differences, we show in our next result that finitely generated monoids have finite successive distance and finite monotone catenary degree (both results are known in the cancellative setting, [19, Theorem 3.1.4] and [13, Theorem 3.9]).

Theorem 3.1. *Let H be a monoid such that H_{red} is finitely generated. Then $\delta(H) < \infty$ and $\mathfrak{c}_{\text{mon}}(H) < \infty$.*

Proof. By [10, Proposition 3.4], H is a BF-monoid with finite set of distances. Without restriction we may suppose that H is reduced and that $\mathcal{A}(H) = \{u_1, \dots, u_s\}$ is nonempty. Let $\pi: Z(H) \rightarrow H$ be the factorization homomorphism. The homomorphism

$$f: Z(H) \times Z(H) \rightarrow (\mathbb{N}_0^s \times \mathbb{N}_0^s, +), \left(\prod_{i=1}^s u_i^{m_i}, \prod_{i=1}^s u_i^{n_i} \right) \mapsto ((m_i)_{i=1}^s, (n_i)_{i=1}^s).$$

is an isomorphism. By Dickson's Theorem ([19, Theorem 1.5.3]), every subset $A \subset \mathbb{N}_0^{2s}$ has only finitely many minimal points.

1. For every distance $d \in \Delta(H)$, we define subsets $R_d^+, R_d^- \subset Z(H) \times Z(H)$ as follows: let R_d^\pm consist of all $(z, y) \in Z(H) \times Z(H)$ such that

$$\pi(z) = \pi(y), \quad |z| \text{ and } |y| \text{ are adjacent lengths of } \pi(z), \text{ and } |y| = |z| \pm d.$$

The sets $M_d^\pm = \text{Min}(f(R_d^\pm))$ of minimal points of $f(R_d^\pm)$ are finite. We set

$$\delta^* = \max\{|z'|, |y'| \mid (z', y') \in f^{-1}(M_d^+ \cup M_d^-), d \in \Delta(H)\},$$

and assert that $\delta(z) \leq \delta^*$ for all $z \in Z(H)$.

Let $z \in Z(H)$ and $k \in \mathbb{N}$ be such that k and $|z|$ are adjacent lengths of $\pi(z)$, and let $y_0 \in Z(H)$ be any factorization with $\pi(y_0) = \pi(z)$ and $|y_0| = k$. Then we have $k = |z| \pm d$ for some $d \in \Delta(H)$ and $(z, y_0) \in R_d^\pm$. Let $(z', y') \in f^{-1}(M_d^\pm)$ be such that $f(z', y') \leq f(z, y_0)$. Then $z = z'z_1$ and $y_0 = y'y_1$ for some $z_1, y_1 \in Z(H)$, and we set $y = y'z_1 \in Z(H)$. Then $\pi(y) = \pi(y')\pi(z_1) = \pi(z')\pi(z_1) = \pi(z)$,

$$|y| = |y'| + |z_1| = |z| + |y'| - |z'| = |z| \pm d = k, \quad \text{and} \quad \mathfrak{d}(z, y) \leq \max\{|z'|, |y'|\} \leq \delta^*.$$

2. Since $\mathfrak{c}_{\text{adj}}(H) \leq \delta(H) < \infty$, it remains to show that $\mathfrak{c}_{\text{eq}}(H) < \infty$. Consider the monoid

$$S = \{(x, y) \in Z(H) \times Z(H) \mid |x| = |y| \text{ and } \pi(xz) = \pi(yz) \text{ for some } z \in Z(H)\},$$

and set $S^* = S \setminus \{(1, 1)\}$. First we show that S is a saturated submonoid of $Z(H) \times Z(H)$. Clearly, S is a submonoid of $Z(H) \times Z(H)$. Let $(x, y), (x', y') \in S$ and $(x'', y'') \in Z(H) \times Z(H)$ be such that $(x, y) = (x'x'', y'y'')$. There are some $z, z' \in Z(H)$ such that $\pi(xz) = \pi(yz)$ and $\pi(x'z') = \pi(y'z')$. Set $z'' = x'y'z'z$. Then $\pi(y''z'') = \pi(yz'x'z') = \pi(yz)\pi(x'z') = \pi(xz)\pi(y'z') = \pi(xzy'z') = \pi(x''z'')$. Moreover, $|x''| = |x| - |x'| = |y| - |y'| = |y''|$, and thus $(x'', y'') \in S$.

Recall that a subset $X \subset Z(H)$ is an s -ideal of $Z(H)$ if $xy \in X$ for all $x \in X$ and $y \in Z(H)$. Note that $Z(H)$ and $Z(H) \times Z(H)$ are finitely generated reduced cancellative monoids. Consequently, $Z(H)$ satisfies the ascending chain condition on s -ideals by [19, Proposition 2.7.4] and S is a finitely generated monoid by [19, Proposition 2.7.5.1]. For $(x, y) \in S$ set

$$A(x, y) = \{z \in Z(H) \mid \pi(xz) = \pi(yz)\}, \text{ and}$$

$$B(x, y) = \{z \in Z(H) \mid (x', y') \mid_S (x, y) \text{ and } \pi(x'z) = \pi(y'z) \text{ for some } (x', y') \in S^* \setminus \{(x, y)\}\}.$$

Observe that if $(x, y), (v, w) \in S$ are such that $(v, w) \mid_S (x, y)$, then $A(x, y)$ and $B(x, y)$ are s -ideals of $Z(H)$, $B(v, w) \subset B(x, y)$ and if $(1, 1) \neq (v, w) \neq (x, y)$, then $A(v, w) \subset B(x, y)$.

A1. For all $t \in \mathbb{N}$, $a \in S$ and $(b_i)_{i=1}^t \in S^t$ there is some $N \in \mathbb{N}$ such that for all $(k_i)_{i=1}^t \in \mathbb{N}_0^t$ and $j \in [1, t]$ with $k_j \geq N$ it follows that $B(a \prod_{i=1}^t b_i^{k_i}) = B(ab_j^N \prod_{i=1, i \neq j}^t b_i^{k_i})$.

Proof of A1. We proceed by induction on t . Let $t \in \mathbb{N}$, $a \in S$ and $(b_i)_{i=1}^t \in S^t$. Note that $(B(a \prod_{i=1}^t b_i^{k_i}))_{k \in \mathbb{N}_0}$ is an ascending chain of s -ideals of $Z(H)$. Therefore, there is some $M \in \mathbb{N}$ such that for all $k \in \mathbb{N}_0$ with $k \geq M$ it follows that $B(a \prod_{i=1}^t b_i^k) = B(a \prod_{i=1}^t b_i^M)$.

By the induction hypothesis there is some $(N_{(r,g)})_{(r,g) \in [1,t] \times [0, M-1]} \in \mathbb{N}^{[1,t] \times [0, M-1]}$ such that for all $r \in [1, t]$, $g \in [0, M-1]$, $(k_i)_{i=1}^t \in \mathbb{N}_0^t$ and $j \in [1, t] \setminus \{r\}$ such that $k_r = g$ and $k_j \geq N_{(r,g)}$ it follows that

$B(a \prod_{i=1}^t b_i^{k_i}) = B(ab_j^{N(r,g)} \prod_{i=1, i \neq j}^t b_i^{k_i})$. Set $N = \max\{N_{(r,g)} \mid (r,g) \in [1, t] \times [0, M-1]\} \cup \{M\}$. Let $(k_i)_{i=1}^t \in \mathbb{N}_0^t$ and $j \in [1, t]$ be such that $k_j \geq N$.

CASE 1: $k_r \geq M$ for all $r \in [1, t]$.

Set $k = \max\{k_i \mid i \in [1, t]\}$. It follows that $B(a \prod_{i=1}^t b_i^M) \subset B(ab_j^N \prod_{i=1, i \neq j}^t b_i^{k_i}) \subset B(a \prod_{i=1}^t b_i^{k_i}) \subset B(a \prod_{i=1}^t b_i^k) = B(a \prod_{i=1}^t b_i^M)$, and thus $B(a \prod_{i=1}^t b_i^{k_i}) = B(ab_j^N \prod_{i=1, i \neq j}^t b_i^{k_i})$.

CASE 2: $k_r < M$ for some $r \in [1, t]$.

Note that $k_j \geq N \geq M > k_r$, hence $j \neq r$ and $k_j \geq N \geq N_{(r, k_r)}$. We infer that $B(a \prod_{i=1}^t b_i^{k_i}) = B(ab_j^{N(r, k_r)} \prod_{i=1, i \neq j}^t b_i^{k_i})$. Since $B(ab_j^{N(r, k_r)} \prod_{i=1, i \neq j}^t b_i^{k_i}) \subset B(ab_j^N \prod_{i=1, i \neq j}^t b_i^{k_i}) \subset B(a \prod_{i=1}^t b_i^{k_i})$ we have $B(a \prod_{i=1}^t b_i^{k_i}) = B(ab_j^N \prod_{i=1, i \neq j}^t b_i^{k_i})$. $\square(\mathbf{A1})$

Since S is finitely generated, there are some $t \in \mathbb{N}$ and $b_1, \dots, b_t \in S^*$ such that S is generated by b_1, \dots, b_t . By **A1** there is some $N \in \mathbb{N}$ such that if $(k_i)_{i=1}^t \in \mathbb{N}_0^t$ and $j \in [1, t]$ are such that $k_j \geq N$, then $B(\prod_{i=1}^t b_i^{k_i}) = B(b_j^N \prod_{i=1, i \neq j}^t b_i^{k_i})$, and thus $A(\prod_{i=1}^t b_i^{k_i}) \subset B(b_j \prod_{i=1}^t b_i^{k_i}) = B(\prod_{i=1}^t b_i^{k_i})$.

This implies that $\{a \in S \mid A(a) \not\subset B(a)\} \subset \{\prod_{i=1}^t b_i^{k_i} \mid (k_i)_{i=1}^t \in \mathbb{N}_0^t, k_j < N \text{ for all } j \in [1, t]\}$, hence $\{a \in S \mid A(a) \not\subset B(a)\}$ is finite. Set $K = \max\{d(x, y) \mid (x, y) \in S, A(x, y) \not\subset B(x, y)\}$.

A2. For all $r \in \mathbb{N}_0$, $(x, y) \in S$ and $z \in A(x, y)$ such that $|x| = r$ it follows that xz and yz can be concatenated by a monotone K -chain of factorizations of $\pi(xz)$.

Proof of A2. We proceed by induction on r . Let $r \in \mathbb{N}_0$, $(x, y) \in S$ and $z \in A(x, y)$ be such that $|x| = r$.

CASE 1: $A(x, y) \not\subset B(x, y)$.

We have $d(xz, yz) = d(x, y) \leq K$, and thus the assertion is trivially satisfied.

CASE 2: $A(x, y) \subset B(x, y)$.

There are some $(x', y'), (x'', y'') \in S^* \setminus \{(x, y)\}$ such that $x = x'x''$, $y = y'y''$ and $\pi(x'z) = \pi(y'z)$. Clearly, $|x'| < r$ and $|x''| < r$. Observe that $\pi(x'x''z) = \pi(x'z)\pi(x'') = \pi(y'z)\pi(x'') = \pi(y'x''z)$ and $\pi(x''y'z) = \pi(x''x'z) = \pi(y''y'z)$, where the last equality holds because $z \in A(x, y)$. We have $x''z \in A(x', y')$ and $y'z \in A(x'', y'')$. Consequently, $xz = x'x''z$ and $y'x''z$ can be concatenated by a monotone K -chain of factorizations of $\pi(xz)$ by the induction hypothesis. It follows by analogy that $x''y'z$ and $y''y'z = yz$ can be concatenated by a monotone K -chain of factorizations of $\pi(x''y'z) = \pi(xz)$. Therefore, xz and yz can be concatenated by a monotone K -chain of factorizations of $\pi(xz)$. $\square(\mathbf{A2})$

It is sufficient to show that $c_{\text{eq}}(H) \leq K$. Let $a \in H$ and $x, y \in Z(a)$ be such that $|x| = |y|$. Then $(x, y) \in S$ and $1 \in A(x, y)$. We infer by **A2** that x and y can be concatenated by a monotone K -chain of factorizations of a . Therefore, $c_{\text{eq}}(b) \leq K$ for each $b \in H$, hence $c_{\text{eq}}(H) \leq K$. \square

We will apply Theorem 3.1 to finitely generated monoids of ideals in Section 4. We refer to [10, Section 3.3] and to [24] for a discussion of finitely generated (not necessarily cancellative) semigroups of modules.

We continue with an example (due to S. Tringali) of a finitely generated monoid whose monoid of equal-length relations is not finitely generated, although the monoid S , defined in the proof of Theorem 3.1.2, is finitely generated and its equal catenary degree is finite.

Example 3.2. For an atomic monoid H ,

$$\sim_{H, \text{eq}} = \{(x, y) \in Z(H) \times Z(H) \mid \pi(x) = \pi(y) \text{ and } |x| = |y|\}$$

is the monoid of equal-length relations of H . If H is cancellative, then $\sim_{H, \text{eq}} \subset \sim_H$ is a saturated submonoid and hence a Krull monoid and it is finitely generated whenever H_{red} is finitely generated ([7, Proposition 4.4]). We provide an example showing that $\sim_{H, \text{eq}}$ need not be finitely generated if H is not cancellative. Let

$$\mathcal{P}_{\text{fin}, 0}(\mathbb{N}_0) = \{A \subset \mathbb{N}_0 \mid 0 \in A, A \subset \mathbb{N}_0 \text{ is finite}\}$$

be the monoid of finite subsets of \mathbb{N}_0 containing 0, with set addition as operation. Clearly, $\{0\}$ is the zero-element of the monoid, and for every $A \subset \mathbb{N}_0$ and every $k \in \mathbb{N}$, $kA = A + \dots + A$ means the k -fold sumset of A . Consider the submonoid H generated by

$$\{0, 1\}, \quad A = \{0, 1, 3\}, \quad \text{and} \quad B = \{0, 2, 3\}.$$

Of course, H is a reduced, finitely generated, commutative semigroup with identity element, and it is unit-cancellative by [11, Theorem 2.22(ii) and Proposition 3.3]. Moreover, we have (by induction) that $\{0, 1\} + kA = \{0, 1\} + kB = [0, 3k + 1]$ for all $k \in \mathbb{N}_0$. Since the monoid H is written additively, $Z(H) \times Z(H)$ and $\sim_{H, \text{eq}}$ will be written additively too. For every $k \in \mathbb{N}_0$, we define

$$\mathbf{a}_k = (\{0, 1\} + kA, \{0, 1\} + kB) \in \sim_{H, \text{eq}},$$

and assert that \mathbf{a}_k is an atom in $\sim_{H, \text{eq}}$.

Assume to the contrary that there are $k \in \mathbb{N}$ and $\mathbf{b}, \mathbf{c} \in \sim_{H, \text{eq}}$, distinct from the zero-element $(\{0\}, \{0\})$ of $\sim_{H, \text{eq}}$, such that $\mathbf{a}_k = \mathbf{b} + \mathbf{c}$. Since $\{0, 1\} + \ell A, \{0, 1\} + \ell B$ are intervals for all $\ell \in \mathbb{N}$ but ℓA and ℓB are not intervals for any $\ell \in \mathbb{N}$, it follows that $(\{0, 1\}, \{0, 1\})$ must divide (in $\sim_{H, \text{eq}}$) either \mathbf{b} or \mathbf{c} , say it divides \mathbf{b} . Thus we obtain that for some $q \in [0, k - 1]$,

$$\mathbf{b} = (\{0, 1\} + qA, \{0, 1\} + qB) \text{ which implies that } \mathbf{c} = ((k - q)A, (k - q)B).$$

However, since $k - q > 0$, we obtain that $1 \in (k - q)A$ but $1 \notin (k - q)B$, a contradiction to $\mathbf{c} \in \sim_{H, \text{eq}}$.

The monoid of v -invertible v -ideals studied in Section 5 is isomorphic to a finite direct product of finitely primary monoids and a free abelian factor. Our next goal in this section is to shift the finiteness of the monotone catenary degree of given monoids H_1 and H_2 to their direct product $H_1 \times H_2$. However, this may not be true in general. We start with an example highlighting the problem.

Example 3.3. Let $k, d \in \mathbb{N}$ with $d \geq 10$, $y_{k,1}, y_{k,2} \in \mathbb{N}_{\geq 2}$ and let

$$L_{k,1} = y_{k,1} + (\{\nu d \mid \nu \in [0, k]\} \cup \{kd + 2\}) \quad \text{and} \quad L_{k,2} = y_{k,2} + (\{\nu d \mid \nu \in [0, k]\} \cup \{kd + 1, kd + 3\}).$$

We claim that both, $y_k + kd + 1$ and $y_k + kd + 2$, have a unique representation in the sumset, where $y_k = y_{k,1} + y_{k,2}$. Indeed, if $y_k + kd + 1 = a_1 + a_2$ with $a_1 \in L_{k,1}$ and $a_2 \in L_{k,2}$, then $a_1 = y_{k,1} + 0$ and $a_2 = y_{k,2} + kd + 1$. If $y_k + kd + 2 = b_1 + b_2$ with $b_1 \in L_{k,1}$ and $b_2 \in L_{k,2}$, then $b_1 = y_{k,1} + kd + 2$ and $b_2 = y_{k,2} + 0$. Thus $|a_1 - b_1| \geq kd$ and $|a_2 - b_2| \geq kd$.

Now suppose that $L_{k,1}$ and $L_{k,2}$ are realized as sets of lengths, say $L_{k,i} = \mathbf{L}(a_{k,i})$ with $a_{k,i} \in H_i$ for some atomic monoids H_i and $i \in [1, 2]$, and set $a_k = a_{k,1}a_{k,2}$ (there is a variety of realization results for sets of lengths guaranteeing the existence of H_1 and H_2 ; see, for example, [19, Proposition 4.8.3], [20, Theorem 4.2]). Then

$$\mathbf{d}(Z_{y_k + kd + 1}(a_k), Z_{y_k + kd + 2}(a_k)) \geq 2kd.$$

For a further analysis of the problem, consider an atomic monoid H . If $a \in H$ and $k, \ell \in \mathbf{L}(a)$ are adjacent lengths with $k < \ell$, then by definition we have $\mathbf{d}(Z_k(a), Z_\ell(a)) \leq \mathbf{c}_{\text{adj}}(a) \leq \mathbf{c}_{\text{adj}}(H)$. However, if $\ell, m \in \mathbf{L}(a)$ are adjacent with $\ell < m$, then we cannot conclude that

$$\mathbf{d}(Z_k(a), Z_m(a)) \leq \mathbf{d}(Z_k(a), Z_\ell(a)) + \mathbf{d}(Z_\ell(a), Z_m(a)).$$

In order to be able to obtain an inductive upper bound for $\mathbf{d}(Z_{k_0}(a), Z_{k_r}(a))$ for pairwise adjacent lengths $k_0 < \dots < k_r$ in $\mathbf{L}(a)$, we introduce a new invariant.

Definition 3.4. Let H be an atomic monoid. For an element $a \in H$, the (*weak*) *successive distance* $\delta_w(a)$ is the smallest $N \in \mathbb{N}_0 \cup \{\infty\}$ such that for all $k, \ell \in \mathbf{L}(a)$ we have $\mathbf{d}(Z_k(a), Z_\ell(a)) \leq N|\ell - k|$. Then

$$\delta_w(H) = \sup\{\delta_w(a) \mid a \in H\} \in \mathbb{N}_0 \cup \{\infty\}$$

is called the (*weak*) *successive distance* of H .

Note that $\delta_w(H) = 0$ if and only if H is half-factorial. The invariant $\delta_w(H)$ is bounded above by $\delta(H)$ (as we outline in the next lemma). However, in Section 5 we will meet monoids having finite weak successive distance but infinite strong successive distance (see Corollary 5.7, Proposition 5.12, and Remark 5.3.4).

Lemma 3.5. *Let H be an atomic monoid.*

1. $\delta_w(H) \leq \delta(H)$.
2. Suppose that $\Delta(H)$ is finite. Then $c_{\text{adj}}(H) \leq \delta_w(H) \max \Delta(H)$. In particular, if $\delta_w(H) < \infty$ and $c_{\text{eq}}(H) < \infty$, then $c_{\text{mon}}(H) < \infty$.

Proof. If H is half-factorial, then $\Delta(H) = \emptyset$ and $c_{\text{adj}}(H) = \max \Delta(H) = \delta_w(H) = 0$ whence all assertions hold. Now we suppose that H is not half-factorial.

1. Clearly, it is sufficient to show that $\delta_w(a) \leq \delta(H)$ for every $a \in H$. Let $a \in H$ and $k, \ell \in \mathbf{L}(a)$ be given, say $k < \ell$. Then there are pairwise adjacent lengths $k = k_0 < k_1 < \dots < k_r = \ell$ of $\mathbf{L}(a)$. By definition of $\delta(H)$, there are factorizations z_0, \dots, z_r such that $|z_i| = k_i$ for $i \in [0, r]$ and $d(z_{i-1}, z_i) \leq \delta(H)$ for all $i \in [1, r]$. Therefore, it follows that

$$d(\mathbf{Z}_{k_0}(a), \mathbf{Z}_{k_r}(a)) \leq d(z_0, z_r) \leq \sum_{i=1}^r d(z_{i-1}, z_i) \leq r\delta(H) \leq |\ell - k|\delta(H).$$

2. If $a \in H$ and $k, \ell \in \mathbf{L}(a)$ are adjacent, then $d(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq \delta_w(a) \max \Delta(\mathbf{L}(a))$ whence $c_{\text{adj}}(a) \leq \delta_w(a) \max \Delta(\mathbf{L}(a))$. Thus it follows that $c_{\text{adj}}(H) \leq \delta_w(H) \max \Delta(H)$, and the in particular statement follows immediately. \square

Lemma 3.6. *Let H be an atomic monoid satisfying the Structure Theorem for Sets of Lengths and suppose that the following conditions hold:*

- (a) *There are some $M_1, C_1 \in \mathbb{N}$ such that for each $a \in H$ and all adjacent $k, \ell \in \mathbf{L}(a)$ such that $\max\{k, \ell\} + M_1 \leq \max \mathbf{L}(a)$ it follows that $\text{Dist}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq C_1$.*
- (b) *For every $N \in \mathbb{N}$ there is some $C_2 \in \mathbb{N}$ such that for all $a \in H$ and $k, \ell \in \mathbf{L}(a)$ for which $\min\{k, \ell\} + N \geq \max \mathbf{L}(a)$ it follows that $d(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq C_2|\ell - k|$.*

Then $\delta_w(H) < \infty$.

Proof. Without restriction we may suppose that H be reduced. We start with the following assertion.

- A.** There is some $M_0 \in \mathbb{N}$ such that for all $a \in H$ and $k \in \mathbf{L}(a)$ with $k + 2M_0 \leq \max \mathbf{L}(a)$ we have $k + M_0 \in \mathbf{L}(a)$.

Proof of A. Since H satisfies the Structure Theorem for Sets of Lengths there is some $N \in \mathbb{N}$ and some finite nonempty $\Delta \subset \mathbb{N}$ such that for all $a \in H$, $\mathbf{L}(a)$ is an AAMP with bound N and difference in Δ . Set $M_0 = N \prod_{d \in \Delta} d$. Let $a \in H$ and $k \in \mathbf{L}(a)$ be such that $k + 2M_0 \leq \max \mathbf{L}(a)$. There is some $d \in \Delta$ such that $\mathbf{L}(a)$ is an AAMP with bound N and difference d . Note that $\min \mathbf{L}(a) + M_0 \leq k + M_0 \leq \max \mathbf{L}(a) - M_0$. Since $k \in \mathbf{L}(a)$ and $M_0 \in d\mathbb{Z}$ we infer that $k + M_0 \in \mathbf{L}(a)$. $\square(\mathbf{A})$

Set $M = M_0 M_1$. It follows easily by induction that for all $a \in H$ and $k \in \mathbf{L}(a)$ such that $k + 2M \leq \max \mathbf{L}(a)$ it follows that $k + M \in \mathbf{L}(a)$. There is some $C_2 \in \mathbb{N}$ such that for all $k, \ell \in \mathbf{L}(a)$ for which $\min\{k, \ell\} + 2M \geq \max \mathbf{L}(a)$ it follows that $d(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq C_2|\ell - k|$.

We assert that $\delta_w(H) \leq \max\{C_1, C_2\}$. Let $a \in H$ and $k, \ell \in \mathbf{L}(a)$, say $k < \ell$. Let $\mathbf{L}(a) \cap [k, \ell] = \{k_0, \dots, k_r\}$ with $k = k_0 < \dots < k_r = \ell$.

CASE 1: $\ell + M \leq \max \mathbf{L}(a)$.

Note that $\text{Dist}(\mathbf{Z}_{k_i}(a), \mathbf{Z}_{k_{i+1}}(a)) \leq C_1$ for all $i \in [0, r-1]$. By definition, there are factorizations $z_i \in \mathbf{Z}_{k_i}(a)$ for every $i \in [0, r]$ such that $d(z_j, z_{j+1}) = d(\{z_j\}, \mathbf{Z}_{k_{j+1}}(a))$ for all $j \in [0, r-1]$ whence we

infer that

$$d(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq d(z_0, z_r) \leq \sum_{j=0}^{r-1} d(z_j, z_{j+1}) \leq C_1 r \leq \max\{C_1, C_2\}|\ell - k|.$$

CASE 2: $k + 2M \geq \max \mathbf{L}(a)$.

It is clear that $d(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq C_2|\ell - k| \leq \max\{C_1, C_2\}|\ell - k|$.

CASE 3: $\ell + M > \max \mathbf{L}(a)$ and $k + 2M < \max \mathbf{L}(a)$.

There is some maximal $n \in [0, r]$ such that $k_n + 2M \leq \max \mathbf{L}(a)$. Observe that $k_n + M \in \mathbf{L}(a)$. Therefore, there is some $m \in [1, r-1]$ such that $k_m = k_n + M$. There are factorizations z_0, \dots, z_{m+1} such that $z_0 \in \mathbf{Z}_\ell(a)$, $z_i \in \mathbf{Z}_{k_{m+1-i}}(a)$ for every $i \in [1, m+1]$, $d(z_0, z_1) = d(\mathbf{Z}_{k_m}(a), \mathbf{Z}_\ell(a))$ and $d(z_j, z_{j+1}) = d(\{z_j\}, \mathbf{Z}_{k_{m-j}}(a))$ for every $j \in [1, m]$. If $j \in [1, m]$, then $k_{m-j+1} + M \leq \max \mathbf{L}(a)$, and thus $d(z_j, z_{j+1}) \leq C_1$. Moreover, since $k_m + 2M \geq \max \mathbf{L}(a)$ we have $d(z_0, z_1) \leq C_2(\ell - k_m)$. This implies that

$$d(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq d(z_0, z_{m+1}) \leq \sum_{i=0}^m d(z_i, z_{i+1}) \leq C_2(\ell - k_m) + C_1 m \leq \max\{C_1, C_2\}|\ell - k|.$$

□

Lemma 3.7. *Let $H = H_1 \times \dots \times H_n$ where $n \in \mathbb{N}$ and H_1, \dots, H_n are atomic monoids.*

1. $\Delta(H)$ is finite if and only if $\Delta(H_i)$ is finite for every $i \in [1, n]$.
2. The Structure Theorem for Sets of Lengths holds for H if and only if it holds for H_1, \dots, H_n .

Proof. Without restriction we may suppose that H is reduced. Then H_1, \dots, H_n are reduced and they are divisor-closed submonoids of H . Thus, if $\Delta(H)$ is finite (or if the Structure Theorem for Sets of Lengths holds for H), then the same is true for each divisor-closed submonoid.

If $\Delta(H_1), \dots, \Delta(H_n)$ are finite, then $\Delta(H)$ is finite (a proof in the cancellative setting can be found in [19, Proposition 1.4.5], and the proof in the general setting runs along the same lines). Clearly, we have

$$\mathcal{L}(H) = \{L_1 + \dots + L_n \mid L_i \in \mathcal{L}(H_i) \text{ for all } i \in [1, n]\}$$

whence sets of lengths in H are sumsets of sets of lengths in H_1, \dots, H_n . Thus if the Structure Theorem for Sets of Lengths holds for H_1, \dots, H_n , then it holds for H by [19, Theorem 4.2.16]. □

Theorem 3.8. *Let $H = H_1 \times \dots \times H_n$ where $n \in \mathbb{N}$ and H_1, \dots, H_n are atomic monoids. Then the following statements are equivalent:*

- (a) H satisfies the Structure Theorem for Sets of Lengths, $c_{\text{eq}}(H) < \infty$, and $\delta_w(H) < \infty$.
- (b) For every $i \in [1, n]$, H_i satisfies the Structure Theorem for Sets of Lengths, $c_{\text{eq}}(H_i) < \infty$, and $\delta_w(H_i) < \infty$.

Proof. We may suppose that H is reduced. Then H_1, \dots, H_n are reduced divisor-closed submonoids of H and hence (a) implies (b). In order to show that (b) implies (a) it suffices to handle the case $n = 2$.

Therefore, we suppose that H_1 and H_2 satisfy the Structure Theorem for Sets of Lengths and that $c_{\text{eq}}(H_1)$, $c_{\text{eq}}(H_2)$, $\delta_w(H_1)$, and $\delta_w(H_2)$ are all finite. Lemma 3.7 implies that the Structure Theorem for Sets of Lengths holds for H . Now we proceed in two steps.

1. We show that $\delta_w(H_1 \times H_2) < \infty$. For every $i \in [1, 2]$, there is a finite nonempty set Δ_i and a bound $M_i \in \mathbb{N}$ such that for every $a_i \in H_i$, $\mathbf{L}(a_i)$ is an AAMP with bound M_i and difference in Δ_i and for all $k, \ell \in \mathbf{L}(a_i)$, we have $d(\mathbf{Z}_k(a_i), \mathbf{Z}_\ell(a_i)) \leq \delta_w(H_i)|\ell - k|$.

We set $e = \max\{M_1, M_2\} \prod_{d \in \Delta_1 \cup \Delta_2} d$, and assert that

$$\delta_w(H_1 \times H_2) \leq \max\{\delta_w(H_1), \delta_w(H_2)\}(4e + 1).$$

Let $a \in H$ and $k, \ell \in \mathbf{L}(a)$ be distinct. Then there are some $a_1 \in H_1$ and $a_2 \in H_2$ such that $a = a_1 a_2$. Obviously, there are some $k_1, \ell_1 \in \mathbf{L}(a_1)$ and $k_2, \ell_2 \in \mathbf{L}(a_2)$ such that $k = k_1 + k_2$, $\ell = \ell_1 + \ell_2$ and for all

$v_1, w_1 \in L(a_1)$ and $v_2, w_2 \in L(a_2)$ with $k = v_1 + v_2$ and $\ell = w_1 + w_2$ it follows that $|\ell_1 - k_1| + |\ell_2 - k_2| \leq |v_1 - w_1| + |v_2 - w_2|$. We proceed by proving the following assertion.

A. $|\ell_1 - k_1| + |\ell_2 - k_2| \leq (4e + 1)|\ell - k|$.

Proof of A. We distinguish two cases.

CASE 1: $|\ell_1 - k_1| \leq 2e$ or $|\ell_2 - k_2| \leq 2e$.

Without restriction we may suppose that $|\ell_1 - k_1| \leq 2e$. Since $k_2 - \ell_2 = k - \ell - (k_1 - \ell_1)$ we infer that $|\ell_2 - k_2| \leq |\ell - k| + |\ell_1 - k_1| \leq |\ell - k| + 2e \leq (2e + 1)|\ell - k|$. It follows that $|\ell_1 - k_1| + |\ell_2 - k_2| \leq 2e + (2e + 1)|\ell - k| \leq (4e + 1)|\ell - k|$.

CASE 2: $|\ell_1 - k_1| > 2e$ and $|\ell_2 - k_2| > 2e$.

First, suppose that $k_1 \leq \ell_1$ and $k_2 \geq \ell_2$. We set

$$v_1 = k_1 + e, \quad w_1 = \ell_1 - e, \quad v_2 = k_2 - e, \quad \text{and} \quad w_2 = \ell_2 + e,$$

and observe that

$$\min L(a_1) + M_1 \leq v_1 \leq w_1 \leq \max L(a_1) - M_1 \quad \text{and} \quad \min L(a_2) + M_2 \leq w_2 \leq v_2 \leq \max L(a_2) - M_2.$$

Since e is a multiple of the differences of the AAMPs $L(a_1)$ and $L(a_2)$, it follows that $v_1, w_1 \in L(a_1)$ and $v_2, w_2 \in L(a_2)$. Therefore, since $k = v_1 + v_2$ and $\ell = w_1 + w_2$, our minimality choice of k_1, k_2, ℓ_1 , and ℓ_2 implies that

$$\begin{aligned} |\ell_1 - k_1| + |\ell_2 - k_2| &\leq |v_1 - w_1| + |v_2 - w_2| = |k_1 - \ell_1 + 2e| + |k_2 - \ell_2 - 2e| \\ &= \ell_1 - k_1 - 2e + k_2 - \ell_2 - 2e = |\ell_1 - k_1| + |\ell_2 - k_2| - 4e, \end{aligned}$$

a contradiction.

Second, if $k_1 \geq \ell_1$ and $k_2 \leq \ell_2$, then we again obtain a contradiction. Thus we infer that $(k_1 \leq \ell_1$ and $k_2 \leq \ell_2)$ or $(k_1 \geq \ell_1$ and $k_2 \geq \ell_2)$. In both cases it is obvious that

$$|\ell_1 - k_1| + |\ell_2 - k_2| = |\ell - k| \leq (4e + 1)|\ell - k|. \quad \square(\mathbf{A})$$

Clearly, there are some $x_1 \in Z_{k_1}(a_1)$, $y_1 \in Z_{\ell_1}(a_1)$, $x_2 \in Z_{k_2}(a_2)$, and $y_2 \in Z_{\ell_2}(a_2)$ such that $d(x_1, y_1) = d(Z_{k_1}(a_1), Z_{\ell_1}(a_1))$ and $d(x_2, y_2) = d(Z_{k_2}(a_2), Z_{\ell_2}(a_2))$. Therefore, **A** implies that

$$\begin{aligned} d(Z_k(a), Z_\ell(a)) &\leq d(x_1 x_2, y_1 y_2) \leq d(x_1, y_1) + d(x_2, y_2) \\ &\leq \delta_w(H_1)|\ell_1 - k_1| + \delta_w(H_2)|\ell_2 - k_2| \\ &\leq \max\{\delta_w(H_1), \delta_w(H_2)\}(|\ell_1 - k_1| + |\ell_2 - k_2|) \\ &\leq \max\{\delta_w(H_1), \delta_w(H_2)\}(4e + 1)|\ell - k|. \end{aligned}$$

2. We assert that $c_{\text{eq}}(H_1 \times H_2) \leq N$, where

$$N = \max\{2e(\delta_w(H_1) + \delta_w(H_2)), c_{\text{eq}}(H_1), c_{\text{eq}}(H_2)\}.$$

To prove that $c_{\text{eq}}(H_1 \times H_2) \leq N$ it is sufficient to show that for all $a_1 \in H_1$, $a_2 \in H_2$, $r \in \mathbb{N}_0$, $z_1, v_1 \in Z(a_1)$, $z_2, v_2 \in Z(a_2)$ such that $|v_1| - |z_1| = r$ and $|z_1| + |z_2| = |v_1| + |v_2|$ it follows that $z_1 z_2$ and $v_1 v_2$ can be concatenated by a monotone N -chain. Let $a_1 \in H_1$ and $a_2 \in H_2$. We prove the assertion by induction on r . Let $r \in \mathbb{N}_0$ be such that for every $s \in \mathbb{N}_0$ with $s < r$ and all $z_1, v_1 \in Z(a_1)$, $z_2, v_2 \in Z(a_2)$ such that $|v_1| - |z_1| = s$ and $|z_1| + |z_2| = |v_1| + |v_2|$ it follows that $z_1 z_2$ and $v_1 v_2$ can be concatenated by a monotone N -chain. Let $z_1, v_1 \in Z(a_1)$, $z_2, v_2 \in Z(a_2)$ be such that $|v_1| - |z_1| = r$ and $|z_1| + |z_2| = |v_1| + |v_2|$. Obviously, $|z_2| - |v_2| = |v_1| - |z_1|$. We distinguish two cases.

CASE 1: $|z_2| - |v_2| \leq 2e$.

Observe that $d(Z_{|z_1|}(a_1), Z_{|v_1|}(a_1)) \leq 2e\delta_w(H_1)$ and $d(Z_{|z_2|}(a_2), Z_{|v_2|}(a_2)) \leq 2e\delta_w(H_2)$. Note that there are some $x_1 \in Z_{|z_1|}(a_1)$, $y_1 \in Z_{|v_1|}(a_1)$, $x_2 \in Z_{|z_2|}(a_2)$ and $y_2 \in Z_{|v_2|}(a_2)$ such that $d(x_1, y_1) = d(Z_{|z_1|}(a_1), Z_{|v_1|}(a_1))$ and $d(x_2, y_2) = d(Z_{|z_2|}(a_2), Z_{|v_2|}(a_2))$. There is some monotone $c_{\text{eq}}(H_1)$ -chain which concatenates z_1 and x_1 , and thus there is some monotone $c_{\text{eq}}(H_1)$ -chain which concatenates $z_1 z_2$ and

x_1z_2 . Moreover, there is some monotone $c_{\text{eq}}(H_2)$ -chain which concatenates z_2 and x_2 , and thus there is some monotone $c_{\text{eq}}(H_2)$ -chain which concatenates x_1z_2 and x_1x_2 . Therefore, there is some monotone N -chain which concatenates z_1z_2 and x_1x_2 . Along the same lines one can show that there is some monotone N -chain which concatenates y_1y_2 and v_1v_2 . Observe that $d(x_1x_2, y_1y_2) \leq d(x_1, y_1) + d(x_2, y_2) \leq 2e(\delta_w(H_1) + \delta_w(H_2)) \leq N$. Therefore, there is some monotone N -chain which concatenates z_1z_2 and v_1v_2 .

CASE 2: $|z_2| - |v_2| > 2e$.

Note that $|v_1| - |z_1| > 2e$, $\min L(a_1) + M_1 \leq |z_1| + e \leq \max L(a_1) - M_1$ and $\min L(a_2) + M_2 \leq |z_2| - e \leq \max L(a_2) - M_2$. Since e is a multiple of the differences of the AAMPs $L(a_1)$ and $L(a_2)$ we have $|z_1| + e \in L(a_1)$ and $|z_2| - e \in L(a_2)$. There are some $w_1 \in Z(a_1)$ and $w_2 \in Z(a_2)$ such that $|w_1| = |z_1| + e$ and $|w_2| = |z_2| - e$. It follows that $0 \leq |v_1| - |w_1| = r - e < r$ and $|w_1| + |w_2| = |z_1| + |z_2| = |v_1| + |v_2|$. By the induction hypothesis, w_1w_2 and v_1v_2 can be concatenated by a monotone N -chain. Note that there are some $x_1 \in Z_{|z_1|}(a_1)$, $y_1 \in Z_{|w_1|}(a_1)$, $x_2 \in Z_{|z_2|}(a_2)$, $y_2 \in Z_{|w_2|}(a_2)$ such that $d(x_1, y_1) = d(Z_{|z_1|}(a_1), Z_{|w_1|}(a_1))$ and $d(x_2, y_2) = d(Z_{|z_2|}(a_2), Z_{|w_2|}(a_2))$. There is some monotone $c_{\text{eq}}(H_1)$ -chain which concatenates z_1 and x_1 , and thus there is some monotone $c_{\text{eq}}(H_1)$ -chain which concatenates z_1z_2 and x_1z_2 . Moreover, there is some monotone $c_{\text{eq}}(H_2)$ -chain which concatenates z_2 and x_2 , and thus there is some monotone $c_{\text{eq}}(H_2)$ -chain which concatenates x_1z_2 and x_1x_2 . Therefore, there is some monotone N -chain which concatenates z_1z_2 and x_1x_2 . Along the same lines one can show that there is some monotone N -chain which concatenates y_1y_2 and w_1w_2 . Observe that $d(x_1x_2, y_1y_2) \leq d(x_1, y_1) + d(x_2, y_2) \leq e(\delta_w(H_1) + \delta_w(H_2)) \leq N$. Consequently, there is some monotone N -chain which concatenates z_1z_2 and w_1w_2 , and thus there is some monotone N -chain which concatenates z_1z_2 and v_1v_2 . \square

4. FINITELY GENERATED MONOIDS OF r -IDEALS

Our notation of ideal theory follows [26] with the modifications stemming from the fact that the monoids in this paper do not contain a zero element. We recall some basics in the setting of commutative monoids and will use all notations also for integral domains.

Let H be a cancellative monoid. An *ideal system* on H is a map $r: \mathcal{P}(H) \rightarrow \mathcal{P}(H)$ such that the following conditions are satisfied for all subsets $X, Y \subset H$ and all $c \in H$:

- $X \subset X_r$.
- $X \subset Y_r$ implies $X_r \subset Y_r$.
- $cH \subset \{c\}_r$.
- $cX_r = (cX)_r$.

Let r be an ideal system on H . A subset $I \subset H$ is called an r -ideal if $I_r = I$. We denote by $\mathcal{I}_r(H)$ the set of all nonempty r -ideals, and we define r -multiplication by setting $I \cdot_r J = (IJ)_r$ for all $I, J \in \mathcal{I}_r(H)$. Then $\mathcal{I}_r(H)$ together with r -multiplication is a reduced semigroup with identity element H . Let $\mathcal{F}_r(H)$ denote the semigroup of fractional r -ideals, $\mathcal{F}_r(H)^\times$ the group of r -invertible fractional r -ideals, and $\mathcal{I}_r^*(H) = \mathcal{F}_r^\times(H) \cap \mathcal{I}_r(H)$ the cancellative monoid of r -invertible r -ideals of H with r -multiplication. We say that H satisfies the *r -Krull Intersection Theorem* if

$$\bigcap_{n \in \mathbb{N}_0} (I^n)_r = \emptyset \quad \text{for all } I \in \mathcal{I}_r(H) \setminus \{H\}.$$

For subsets $A, B \subset \mathfrak{q}(H)$, we denote by $(A : B) = \{x \in \mathfrak{q}(H) \mid xB \subset A\}$, by $A^{-1} = (H : A)$, by $A_v = (A^{-1})^{-1}$ and by $A_t = \bigcup_{E \subset A, |E| < \infty} E_v$. We will use the s -system, the v -system, and the t -system. By $\mathfrak{X}(H)$, we denote the set of minimal nonempty prime s -ideals of H . If r and q are ideal systems on H , then we say that r is finer than q (resp. q is coarser than r) if $\mathcal{I}_q(H) \subset \mathcal{I}_r(H)$ (equivalently, $X_r \subset X_q$ for all $X \subset H$). Recall that $X_r \subset X_v$, and if r is finitary, then $X_r \subset X_t$ for all $X \subset H$.

We need the following classes of monoids (see [19, Section 2.10], also for the correspondence to ring theory). Let H be a monoid and $\mathfrak{m} = H \setminus H^\times$. Then H is said to be

- *archimedean* if $\bigcap_{n \geq 0} a^n H = \emptyset$ for every $a \in \mathfrak{m}$,
- *primary* if $H \neq H^\times$ and $s\text{-spec}(H) = \{\emptyset, \mathfrak{m}\}$,
- *strongly primary* if for every $a \in \mathfrak{m}$ there is an $n \in \mathbb{N}$ such that $\mathfrak{m}^n \subset aH$,
- a *G-monoid* if the intersection of all nonempty prime s -ideals is nonempty.

To recall some well-known properties, note that every monoid satisfying the ACCP is archimedean, that every v -Noetherian primary monoid is strongly primary, that every primary monoid is a G-monoid, and that overmonoids of G-monoids are G-monoids.

In order to give the definition of C-monoids we need to recall the concept of class semigroups which are a refinement of ordinary class groups in commutative algebra (a detailed presentation can be found in [19, Chapter 2]). Let F be a cancellative monoid and $H \subset F$ a submonoid. Two elements $y, y' \in F$ are called H -equivalent, if $y^{-1}H \cap F = y'^{-1}H \cap F$. H -equivalence is a congruence relation on F . For $y \in F$, let $[y]_H^F$ denote the congruence class of y , and let

$$\mathcal{C}(H, F) = \{[y]_H^F \mid y \in F\} \quad \text{and} \quad \mathcal{C}^*(H, F) = \{[y]_H^F \mid y \in (F \setminus F^\times) \cup \{1\}\}.$$

Then $\mathcal{C}(H, F)$ is a semigroup with unit element $[1]_H^F$ (called the *class semigroup* of H in F) and $\mathcal{C}^*(H, F) \subset \mathcal{C}(H, F)$ is a subsemigroup (called the *reduced class semigroup* of H in F). If H is a submonoid of a factorial monoid F such that $H \cap F^\times = H^\times$ and $\mathcal{C}^*(H, F)$ is finite, then H is called a *C-monoid*. A C-monoid H is v -Noetherian with $(H : \widehat{H}) \neq \emptyset$ and its complete integral closure \widehat{H} is Krull with finite class group ([19, Theorems 2.9.11 and 2.9.13]). If H is a Krull monoid with finite class group, then H is a C-monoid and the class semigroup coincides with the usual class group of a Krull monoid.

Let R be a domain with quotient field K and r an ideal system on R (clearly, R^\bullet is a monoid and r restricts to an ideal system r' on R^\bullet whence for every subset $I \subset R$ we have $I_r = (I^\bullet)_{r'} \cup \{0\}$). We denote by $\mathcal{I}_r(R)$ the semigroup of nonzero r -ideals of R and $\mathcal{I}_r^*(R) \subset \mathcal{I}_r(R)$ is the subsemigroup of r -invertible r -ideals of R . The usual ring ideals form an ideal system, called the d -system, and for these ideals we omit all suffices (i.e., $\mathcal{I}(R) = \mathcal{I}_d(R)$, and so on).

*Throughout the rest of the paper, every ideal system r on a domain R
has the property that $\mathcal{I}_r(R) \subset \mathcal{I}(R)$.*

A domain R is said to be archimedean (primary, strongly primary, a G-domain, a C-domain) if its multiplicative monoid R^\bullet has the respective property. Recall that a domain is primary if and only if it is one-dimensional local. To give an example for a C-domain, let R be a Mori domain with nonzero conductor $\mathfrak{f} = (R : \widehat{R})$. If the class group $\mathcal{C}(\widehat{R})$ and the factor ring \widehat{R}/\mathfrak{f} are both finite, then R is a C-domain by [19, Theorem 2.11.9] (for more on C-domains see [22, 32]).

In the first part of this section we establish sufficient criteria for the r -ideal semigroup $\mathcal{I}_r(R)$ to be unit-cancellative.

Lemma 4.1. *Let R be a domain and r an ideal system on R .*

1. *If R satisfies the r -Krull Intersection Theorem, then $\mathcal{I}_r(R)$ is unit-cancellative.*
2. *$\mathcal{I}_r(R)$ is unit-cancellative if and only if $\mathcal{I}_r(R)$ has no nontrivial idempotents and for every $I \in \mathcal{I}_r(R)$, $\{J \in \mathcal{I}_r(R) \mid (IJ)_r = I\}$ has a minimal element with respect to inclusion.*

Proof. 1. Let $I, J \in \mathcal{I}_r(R)$ be such that $(IJ)_r = I$. We infer that $(IJ^n)_r = I$ for all $n \in \mathbb{N}_0$, hence $\{0\} \neq I \subset \bigcap_{n \in \mathbb{N}_0} (J^n)_r$. Since R satisfies the r -Krull Intersection Theorem we have $J = R$.

2. First let $\mathcal{I}_r(R)$ be unit-cancellative. It is clear that R is the only idempotent element of $\mathcal{I}_r(R)$ and for every $I \in \mathcal{I}_r(R)$, $\{J \in \mathcal{I}_r(R) \mid (IJ)_r = I\} = \{R\}$.

To prove the converse, suppose that $\mathcal{I}_r(R)$ has no nontrivial idempotents and let for every $I \in \mathcal{I}_r(R)$, $\{B \in \mathcal{I}_r(R) \mid (IB)_r = I\}$ have a minimal element with respect to inclusion. Let $I, J \in \mathcal{I}_r(R)$ be such that $(IJ)_r = I$. We have to show that $J = R$. Let $A \in \mathcal{I}_r(R)$ be minimal such that $(IA)_r = I$. Since

$(IA^2)_r = I$ and $(A^2)_r \subset A$, we infer that A is an idempotent of $\mathcal{I}_r(R)$, and thus $A = R$. Note that $(IJA)_r = I$ and $(JA)_r \subset A$. Therefore, $J = (JA)_r = A = R$. \square

Proposition 4.2. *Let R be a domain and r an ideal system on R .*

1. *If R is strongly primary, then R satisfies the r -Krull Intersection Theorem.*
2. *Let R be archimedean such that \widehat{R} is a semilocal principal ideal domain. If $(R : \widehat{R}) \neq \{0\}$ or r is finitary, then $\mathcal{I}_r(R)$ is unit-cancellative.*
3. *If R is an r -Noetherian G-domain, then $\mathcal{I}_r(R)$ is unit-cancellative.*

Proof. 1. Set $\mathfrak{m} = R \setminus R^\times$. Let $I \in \mathcal{I}_r(R)$ be such that $\bigcap_{n \in \mathbb{N}_0} (I^n)_r \neq \{0\}$. There is some nonzero $x \in \bigcap_{n \in \mathbb{N}_0} (I^n)_r$. We infer that $\mathfrak{m}^k \subset xR$ for some $k \in \mathbb{N}_0$, and thus $(\mathfrak{m}^k)_r \subset xR$. Assume that $I \neq R$. Then $I \subset \mathfrak{m}$, hence $(\mathfrak{m}^k)_r \subset xR \subset \bigcap_{n \in \mathbb{N}_0} (I^n)_r \subset \bigcap_{n \in \mathbb{N}_0} (\mathfrak{m}^n)_r \subset (\mathfrak{m}^k)_r$. This implies that $(\mathfrak{m}^k)_r = xR$. It follows that $(\mathfrak{m}^{2k})_r = (\mathfrak{m}^k)_r$, hence $x^2R = xR$. Consequently, $x \in I \cap R^\times$, and thus $I = R$, a contradiction.

2. Let $(R : \widehat{R}) \neq \{0\}$ or let r be finitary. We start with the following assertion.

A. $(\widehat{R} : A) = (\widehat{R} : A_r)$ for every ideal A of R .

Proof of A. Let A be an ideal of R . It is sufficient to show that $(\widehat{R} : A) \subset (\widehat{R} : A_r)$, since the other inclusion is trivially satisfied. Let $x \in (\widehat{R} : A)$. Then $xA \subset \widehat{R}$. Recall that the v system is the coarsest ideal system on R and the t system is the coarsest finitary ideal system on R .

CASE 1: $(R : \widehat{R}) \neq \{0\}$.

We have \widehat{R} is a fractional divisorial ideal of R , and thus $xA_r \subset xA_v = (xA)_v \subset (\widehat{R})_v = \widehat{R}$.

CASE 2: r is finitary.

It follows that $xA_r \subset xA_t = (xA)_t \subset (\widehat{R})_t = \widehat{R}$. $\square(\mathbf{A})$

Note that $A_{v_{\widehat{R}}} = A\widehat{R}$ for every ideal A of R , since \widehat{R} is a principal ideal domain. Now let $I, J \in \mathcal{I}_r(R)$ be such that $(IJ)_r = I$. We infer by **A** that $I\widehat{R}J\widehat{R} = (IJ)_{v_{\widehat{R}}} = ((IJ)_r)_{v_{\widehat{R}}} = I_{v_{\widehat{R}}} = I\widehat{R}$. Consequently, $J\widehat{R} = \widehat{R}$, since \widehat{R} is a principal ideal domain. Note that J is additively and multiplicatively closed and $J \not\subset \mathfrak{m}$ for every $\mathfrak{m} \in \max(\widehat{R})$. Therefore, $J \not\subset \bigcup_{\mathfrak{m} \in \max(\widehat{R})} \mathfrak{m} = \widehat{R} \setminus \widehat{R}^\times$ by prime avoidance, since \widehat{R} is semilocal. Since R is archimedean, we have $\emptyset \neq J \cap \widehat{R}^\times = J \cap R \cap \widehat{R}^\times = J \cap R^\times$, and thus $J = R$.

3. Let $I, J \in \mathcal{I}_r(R)$ be such that $(IJ)_r = I$. Set $S = (I : I)$. It follows by [31, Theorem 4.1(b)] and its proof that S is a Mori domain and $JS = S$. Observe that S is also a G-domain, and thus $\text{spec}(S)$ is finite. Since J is an ideal of R and $J \not\subset \mathfrak{m}$ for all $\mathfrak{m} \in \max(S)$ we have $J \not\subset \bigcup_{\mathfrak{m} \in \max(S)} \mathfrak{m} = S \setminus S^\times$ by prime avoidance. Therefore, $J \cap S^\times \neq \emptyset$. Clearly, R is archimedean (since R is a Mori domain). Since $S \subset \widehat{R}$, we have $R^\times \subset S^\times \cap R \subset \widehat{R}^\times \cap R = R^\times$, hence $S^\times \cap R = R^\times$. We infer that $\emptyset \neq J \cap S^\times = J \cap R \cap S^\times = J \cap R^\times$, and thus $J = R$. \square

Lemma 4.3. *Let R be a domain and r a finitary ideal system on R .*

1. *If $R_{\mathfrak{m}}$ satisfies the $r_{\mathfrak{m}}$ -Krull Intersection Theorem for every $\mathfrak{m} \in r\text{-max}(R)$, then R satisfies the r -Krull Intersection Theorem.*
2. *If $\mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}})$ is unit-cancellative for every $\mathfrak{m} \in r\text{-max}(R)$, then $\mathcal{I}_r(R)$ is unit-cancellative.*

Proof. 1. Let $I \in \mathcal{I}_r(R)$ be such that $I \neq R$. Since r is finitary, there is some $\mathfrak{m} \in r\text{-max}(R)$ such that $I \subset \mathfrak{m}$. We have $I_{\mathfrak{m}} \in \mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}})$ and $I_{\mathfrak{m}} \subset \mathfrak{m}_{\mathfrak{m}} \subsetneq R_{\mathfrak{m}}$. Therefore, $\bigcap_{n \in \mathbb{N}_0} (I^n)_r \subset \bigcap_{n \in \mathbb{N}_0} (I^n)_{r_{\mathfrak{m}}} = \{0\}$.

2. Let $I, J \in \mathcal{I}_r(R)$ be such that $(IJ)_r = I$. Then $(I_{\mathfrak{m}}J_{\mathfrak{m}})_{r_{\mathfrak{m}}} = ((IJ)_r)_{\mathfrak{m}} = I_{\mathfrak{m}}$ for all $\mathfrak{m} \in r\text{-max}(R)$, and thus $J_{\mathfrak{m}} = R_{\mathfrak{m}}$ for all $\mathfrak{m} \in r\text{-max}(R)$. Consequently, $J = R$. \square

Corollary 4.4. *Let R be a domain and r a finitary ideal system on R .*

1. If R is a Mori domain, then $\mathcal{I}_v(R)$ is unit-cancellative if and only if $\mathcal{I}_v(R)$ has no nontrivial idempotents.
2. Let R be a Mori domain or let \widehat{R} be a Krull domain. If $r\text{-max}(R) = \mathfrak{X}(R)$, then $\mathcal{I}_r(R)$ is unit-cancellative.

Proof. 1. Let R be a Mori domain and $I \in \mathcal{I}_v(R)$. Note that $R \in \{J \in \mathcal{I}_v(R) \mid (IJ)_v = I\} \subset \{J \in \mathcal{I}_v(R) \mid I \subset J\}$. Since R is a Mori domain we infer that $\{J \in \mathcal{I}_v(R) \mid (IJ)_v = I\}$ has a minimal element with respect to inclusion. Now the assertion follows from Lemma 4.1.2.

2. Let $r\text{-max}(R) = \mathfrak{X}(R)$. By Lemma 4.3.2 it is sufficient to show that $\mathcal{I}_{r_m}(R_m)$ is unit-cancellative for all $\mathfrak{m} \in \mathfrak{X}(R)$. Let $\mathfrak{m} \in \mathfrak{X}(R)$.

CASE 1: R is a Mori domain. Then R_m is a one-dimensional local Mori domain.

It follows that R_m is a strongly primary domain, hence $\mathcal{I}_{r_m}(R_m)$ is unit-cancellative by Lemma 4.1.1 and Proposition 4.2.1.

CASE 2: \widehat{R} is a Krull domain.

Clearly, R_m is a one-dimensional local domain, and thus R_m is an archimedean G-domain. Moreover, we have $\widehat{R}_m = \widehat{R}_m$ is a Krull domain and a G-domain. Therefore, \widehat{R}_m is a semilocal principal ideal domain. (A Krull domain that is a G-domain has finite prime spectrum. Therefore, it is semilocal and it is at most one-dimensional by prime avoidance (since a Krull domain satisfies Krull's principal ideal theorem). A Krull domain that is at most one-dimensional is a Dedekind domain and a semilocal Dedekind domain is a principal ideal domain.) It is clear that r_m is a finitary ideal system on R_m . We infer by Proposition 4.2.2 that $\mathcal{I}_{r_m}(R_m)$ is unit-cancellative. \square

A domain is called a *Cohen-Kaplansky domain* if it is atomic and has only finitely many atoms up to associates. Our next result is well-known for usual ring ideals (see [5, Theorem 4.3] and [3]).

Proposition 4.5. *Let R be a domain and r an ideal system on R . Then the following statements are equivalent:*

- (a) R is a Cohen-Kaplansky domain.
- (b) $\mathcal{I}_r(R)$ is a finitely generated semigroup.
- (c) $\mathcal{I}_r^*(R)$ is a finitely generated semigroup.

Proof. First let R be a Cohen-Kaplansky domain. We know that $\mathcal{I}(R)$ is a finitely generated semigroup by [5, Theorem 4.3], and thus there is some finite $E \subset \mathcal{I}(R)$ such that $\mathcal{I}(R) = [E]$. Set $F = \{J_r \mid J \in E\}$. Then F is finite and it suffices to show that $\mathcal{I}_r(R) = [F]$. It is clear that $[F] \subset \mathcal{I}_r(R)$. Let $I \in \mathcal{I}_r(R)$. Then $I \in \mathcal{I}(R)$, and thus there is some $(\alpha_J)_{J \in E} \in \mathbb{N}_0^E$ such that $I = \prod_{J \in E} J^{\alpha_J}$. We infer that $I = I_r = (\prod_{J \in E} (J_r)^{\alpha_J})_r \in [F]$.

Next let $\mathcal{I}_r(R)$ be a finitely generated semigroup. Then there is some finite $E \subset \mathcal{I}_r(R)$ such that $\mathcal{I}_r(R) = [E]$. Set $F = E \cap \mathcal{I}_r^*(R)$. Then F is finite and it suffices to show that $\mathcal{I}_r^*(R) = [F]$. Obviously, $[F] \subset \mathcal{I}_r^*(R)$. Let $I \in \mathcal{I}_r^*(R)$. Then $I \in \mathcal{I}_r(R)$ and hence there is some $(\alpha_J)_{J \in E} \in \mathbb{N}_0^E$ such that $I = (\prod_{J \in E} J^{\alpha_J})_r$. Recall that if $A \in \mathcal{I}_r^*(R)$ and $B, C \in \mathcal{I}_r(R)$ are such that $A = (BC)_r$, then $B, C \in \mathcal{I}_r^*(R)$. This implies that $\alpha_J = 0$ or $J \in F$ for all $J \in E$. Therefore, $I = (\prod_{J \in F} J^{\alpha_J})_r \in [F]$.

Finally let $\mathcal{I}_r^*(R)$ be a finitely generated semigroup. Note that $\mathcal{I}_r^*(R)$ is a cancellative monoid and $\{xR \mid x \in R^\bullet\}$ is a saturated submonoid of $\mathcal{I}_r^*(R)$. We infer that $\{xR \mid x \in R^\bullet\}$ is a finitely generated monoid, and thus the monoid of principal ideals of R is finitely generated. Consequently, R is a Cohen-Kaplansky domain by [5, Theorem 4.3]. \square

Let R be a domain, r an ideal system on R and $I, J \in \mathcal{I}_r(R)$. We say that I and J are *r -coprime* if R is the only r -ideal containing I and J .

Lemma 4.6. *Let R be a domain and r a finitary ideal system on R . Let $I, J, L \in \mathcal{I}_r(R)$, $n \in \mathbb{N}_0$ and $(I_i)_{i=1}^n$ a finite sequence of elements of $\mathcal{I}_r(R)$.*

1. I and J are r -coprime if and only if \sqrt{I} and \sqrt{J} are r -coprime. If these equivalent condition are satisfied, then $(IJ)_r = I \cap J$.
2. If I and J are r -coprime and I and L are r -coprime, then I and $J \cap L$ are r -coprime.
3. If for all distinct $i, j \in [1, n]$, I_i and I_j are r -coprime, then $(\prod_{i=1}^n I_i)_r = \bigcap_{i=1}^n I_i$.

Proof. 1. Since r is finitary, every proper r -ideal of R is contained in an r -maximal r -ideal. Moreover, the radical of every r -ideal is again an r -ideal. Therefore, I and J are not r -coprime if and only if there is some $\mathfrak{m} \in r\text{-max}(R)$ such that $I, J \subset \mathfrak{m}$ if and only if there is some $\mathfrak{m} \in r\text{-max}(R)$ such that $\sqrt{I}, \sqrt{J} \subset \mathfrak{m}$ if and only if \sqrt{I} and \sqrt{J} are not r -coprime.

Now let I and J be r -coprime. Then $(I \cup J)_r = R$. Clearly, $(IJ)_r \subset I \cap J$. Let $x \in I \cap J$. Then $x \in xR = x(I \cup J)_r = (xI \cup xJ)_r \subset (IJ)_r$.

2. Let I and $J \cap L$ be not r -coprime. Then there is some $\mathfrak{m} \in r\text{-max}(R)$ such that $I, J \cap L \subset \mathfrak{m}$. Therefore, $J \subset \mathfrak{m}$ or $L \subset \mathfrak{m}$. Without restriction let $J \subset \mathfrak{m}$. Then I and J are not r -coprime.

3. This follows by induction from 1 and 2. \square

Proposition 4.7. *Let R be a domain and r an r -Noetherian ideal system on R with $r\text{-max}(R) = \mathfrak{X}(R)$. Then $\mathcal{I}_r(R)$ is an atomic monoid and $\sqrt{J} \in \mathfrak{X}(R)$ for every $J \in \mathcal{A}(\mathcal{I}_r(R))$.*

Proof. First we show that $\mathcal{I}_r(R)$ is an atomic monoid. Since R is r -Noetherian, we infer that R is a Mori domain and r is a finitary ideal system. Therefore, $\mathcal{I}_r(R)$ is unit-cancellative by Corollary 4.4.2.

Next we show that $\mathcal{I}_r(R)$ satisfies the ACCP. Let $(J_i)_{i \in \mathbb{N}}$ be an ascending chain of principal ideals of $\mathcal{I}_r(R)$. There is some sequence $(I_i)_{i \in \mathbb{N}}$ of elements of $\mathcal{I}_r(R)$ such that $J_i = \{(I_i A)_r \mid A \in \mathcal{I}_r(R)\}$ for each $i \in \mathbb{N}$. Let $j \in \mathbb{N}$. Then $I_j = (I_j R)_r \in J_j \subset J_{j+1}$, and thus there is some $B \in \mathcal{I}_r(R)$ such that $I_j = (I_{j+1} B)_r$. We infer that $I_i \subset I_{i+1}$ for every $i \in \mathbb{N}$. Since R is r -Noetherian, there is some $m \in \mathbb{N}$ such that $I_n = I_m$ for all $n \in \mathbb{N}_{\geq m}$. This clearly implies that $J_n = J_m$ for all $n \in \mathbb{N}_{\geq m}$. Therefore, $\mathcal{I}_r(R)$ satisfies the ACCP. It follows from [10, Lemma 3.1(1)] that $\mathcal{I}_r(R)$ is atomic.

Let $I \in \mathcal{I}_r(R)$. We show that $I = (\prod_{\mathfrak{p} \in \mathfrak{X}(R), I \subset \mathfrak{p}} (I_{\mathfrak{p}} \cap R))_r$ and $\sqrt{I_{\mathfrak{q}} \cap R} = \mathfrak{q}$ for every $\mathfrak{q} \in \mathfrak{X}(R)$ with $I \subset \mathfrak{q}$. Clearly, R is a weakly Krull domain, and thus $\{\mathfrak{q} \in \mathfrak{X}(R) \mid I \subset \mathfrak{q}\}$ is finite. Moreover, $I = \bigcap_{\mathfrak{p} \in \mathfrak{X}(R)} I_{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in \mathfrak{X}(R), I \subset \mathfrak{p}} I_{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in \mathfrak{X}(R), I \subset \mathfrak{p}} (I_{\mathfrak{p}} \cap R)$. Let $\mathfrak{q} \in \mathfrak{X}(R)$ with $I \subset \mathfrak{q}$. Since r is finitary it is straightforward to prove that $I_{\mathfrak{q}} \cap R$ is an r -ideal of R . Moreover, $\sqrt{I_{\mathfrak{q}}} = \mathfrak{q}_{\mathfrak{q}}$ (since $R_{\mathfrak{q}}$ is a one-dimensional local domain), and hence $\sqrt{I_{\mathfrak{q}} \cap R} = \mathfrak{q}$. Therefore, $\bigcap_{\mathfrak{p} \in \mathfrak{X}(R), I \subset \mathfrak{p}} (I_{\mathfrak{p}} \cap R)$ is a finite intersection of pairwise r -coprime r -ideals of R by Lemma 4.6.1 (since their radicals are pairwise r -coprime r -ideals of R). Consequently, $I = (\prod_{\mathfrak{p} \in \mathfrak{X}(R), I \subset \mathfrak{p}} (I_{\mathfrak{p}} \cap R))_r$ by Lemma 4.6.3.

Let $J \in \mathcal{A}(\mathcal{I}_r(R))$. Then $J = (\prod_{\mathfrak{p} \in \mathfrak{X}(R), J \subset \mathfrak{p}} (J_{\mathfrak{p}} \cap R))_r$, hence $J = J_{\mathfrak{p}} \cap R$ for some $\mathfrak{p} \in \mathfrak{X}(R)$ such that $J \subset \mathfrak{p}$. We infer that $\sqrt{J} = \sqrt{J_{\mathfrak{p}} \cap R} = \mathfrak{p} \in \mathfrak{X}(R)$. \square

Theorem 4.8. *Let R be a domain with $(R : \widehat{R}) \neq \{0\}$ and r an r -Noetherian ideal system on R with $r\text{-max}(R) = \mathfrak{X}(R)$. We set $\mathcal{P} = \{\mathfrak{p} \in \mathfrak{X}(R) \mid \mathfrak{p} \not\subset (R : \widehat{R})\}$, $\mathcal{P}^* = \mathfrak{X}(R) \setminus \mathcal{P}$, and let T be the subsemigroup of $\mathcal{I}_r(R)$ generated by $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$.*

1. T is a monoid and $\mathcal{I}_r(R) \cong \mathcal{F}(\mathcal{P}) \times T$.
2. T is finitely generated if and only if $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$ is finite.

Proof. We start with the following assertion.

A. For every $\mathfrak{p} \in \mathcal{P}$, \mathfrak{p} is r -invertible and $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} = \mathfrak{p}\} = \{\mathfrak{p}\}$.

Proof of A. Let $\mathfrak{p} \in \mathcal{P}$. It follows from [19, Theorem 2.10.9] that $R_{\mathfrak{p}}$ is a discrete valuation domain. This implies that $\mathfrak{p}_{\mathfrak{q}}$ is a principal ideal of $R_{\mathfrak{q}}$ for every $\mathfrak{q} \in r\text{-max}(R)$, and thus \mathfrak{p} is an r -invertible r -ideal of R by [26, 12.3 Theorem] (since R is r -Noetherian). Clearly, $\mathfrak{p} \in \{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} = \mathfrak{p}\}$. Let $I \in \mathcal{A}(\mathcal{I}_r(R))$ be such that $\sqrt{I} = \mathfrak{p}$. It is sufficient to show that $I = (\mathfrak{p}^k)_r$ for some $k \in \mathbb{N}$ (then $I = \mathfrak{p}$, since $I \in \mathcal{A}(\mathcal{I}_r(R))$). There is some $\ell \in \mathbb{N}$ such that $(\mathfrak{p}^{\ell})_r \subset I$ (note that $\mathfrak{p} = J_r$ for some

finite $J \subset \mathfrak{p}$). Since \mathfrak{p} is r -invertible there is some greatest $k \in \mathbb{N}$ such that $I \subset (\mathfrak{p}^k)_r$ (otherwise we have $(\mathfrak{p}^\ell)_r \subset I \subset (\mathfrak{p}^{\ell+1})_r$, which contradicts the fact that \mathfrak{p} is a proper r -invertible r -ideal). Suppose that $I \subsetneq (\mathfrak{p}^k)_r$. Then $(I\mathfrak{p}^{-k})_r \subsetneq R$, and thus there is some $\mathfrak{q} \in \mathfrak{X}(R)$ such that $(I\mathfrak{p}^{-k})_r \subset \mathfrak{q}$. We have $I \subset (\mathfrak{p}^k\mathfrak{q})_r \subset \mathfrak{q}$, hence $\mathfrak{p} = \mathfrak{q}$ and $I \subset (\mathfrak{p}^{k+1})_r$, a contradiction. \square (A)

1. By definition, T is a semigroup with identity R and since $\mathcal{I}_r(R)$ is a reduced monoid by Proposition 4.7, it follows that T is unit-cancellative, and thus T is a monoid.

Let $f : \mathcal{F}(\mathcal{P}) \times T \rightarrow \mathcal{I}_r(R)$ be defined by $f((\prod_{\mathfrak{p} \in \mathcal{P}} \mathfrak{p}^{\alpha_{\mathfrak{p}}}, A)) = (\prod_{\mathfrak{p} \in \mathcal{P}} \mathfrak{p}^{\alpha_{\mathfrak{p}}} A)_r$ where $(\alpha_{\mathfrak{p}})_{\mathfrak{p} \in \mathcal{P}}$ is a formally infinite sequence of nonnegative integers. Clearly, f is a well-defined monoid homomorphism. It remains to show that f is bijective.

First we show that f is injective. Let $(\alpha_{\mathfrak{p}})_{\mathfrak{p} \in \mathcal{P}}$ and $(\beta_{\mathfrak{p}})_{\mathfrak{p} \in \mathcal{P}}$ be formally infinite sequences of nonnegative integers and $A, B \in T$ be such that $(\prod_{\mathfrak{p} \in \mathcal{P}} \mathfrak{p}^{\alpha_{\mathfrak{p}}} A)_r = (\prod_{\mathfrak{p} \in \mathcal{P}} \mathfrak{p}^{\beta_{\mathfrak{p}}} B)_r$. Since every $\mathfrak{q} \in \mathcal{P}$ is r -invertible it is sufficient to show that $\alpha_{\mathfrak{q}} = \beta_{\mathfrak{q}}$ for all $\mathfrak{q} \in \mathcal{P}$ (then it clearly follows that $A = B$). Let $\mathfrak{q} \in \mathcal{P}$. Suppose that $\alpha_{\mathfrak{q}} \neq \beta_{\mathfrak{q}}$. Without restriction let $\alpha_{\mathfrak{q}} < \beta_{\mathfrak{q}}$. Since \mathfrak{q} is r -invertible we have $(\prod_{\mathfrak{p} \in \mathcal{P}, \mathfrak{p} \neq \mathfrak{q}} \mathfrak{p}^{\alpha_{\mathfrak{p}}} A)_r = (\prod_{\mathfrak{p} \in \mathcal{P}, \mathfrak{p} \neq \mathfrak{q}} \mathfrak{p}^{\beta_{\mathfrak{p}}} \mathfrak{q}^{\beta_{\mathfrak{q}} - \alpha_{\mathfrak{q}}} B)_r \subset \mathfrak{q}$. Consequently, $A \subset \mathfrak{q}$. Since A is a finite r -product of elements of $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$, there is some $B \in \mathcal{A}(\mathcal{I}_r(R))$ such that $\sqrt{B} \in \mathcal{P}^*$ and $B \subset \mathfrak{q}$. We infer that $\mathfrak{q} = \sqrt{B} \in \mathcal{P}^*$, a contradiction.

Next we show that f is surjective. Let $I \in \mathcal{I}_r(R)$. Since $\mathcal{I}_r(R)$ is atomic (by Proposition 4.7), we have $I = (\prod_{i=1}^n I_i)_r$ for some $n \in \mathbb{N}_0$ and some $I_i \in \mathcal{A}(\mathcal{I}_r(R))$. Set $A = (\prod_{i \in [1, n], \sqrt{I_i} \in \mathcal{P}^*} I_i)_r$ and $B = (\prod_{i \in [1, n], \sqrt{I_i} \in \mathcal{P}^*} I_i)_r$. Then $B \in T$. It follows by Proposition 4.7 that $I = (AB)_r$. It follows by (A) that $A = (\prod_{\mathfrak{p} \in \mathcal{P}} \mathfrak{p}^{\alpha_{\mathfrak{p}}})_r$ for some formally infinite sequence of nonnegative integers $(\alpha_{\mathfrak{p}})_{\mathfrak{p} \in \mathcal{P}}$. Consequently, $f((\prod_{\mathfrak{p} \in \mathcal{P}} \mathfrak{p}^{\alpha_{\mathfrak{p}}}, B)) = I$.

2. It follows by 1 that $\mathcal{A}(\mathcal{I}_r(R)) \cap T = \mathcal{A}(T)$ and $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\} = \mathcal{A}(T)$. This immediately implies that T is finitely generated if and only if $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$ is finite. \square

If R is an order in a quadratic number field and the r -ideals are the usual ring ideals, then there is a very explicit number theoretic characterization of when the set $\{I \in \mathcal{A}(\mathcal{I}(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$ is finite ([10, Corollary 3.8]). Here we continue with a discussion of generalized Cohen-Kaplansky domains which provide a further setting where the set $\{I \in \mathcal{A}(\mathcal{I}(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$ is finite (see Theorem 4.13).

A domain is called a *generalized Cohen-Kaplansky domain* if it is atomic and has only finitely many atoms up to associates that are not prime elements. Generalized Cohen-Kaplansky domains were introduced by D.D. Anderson, D.F. Anderson, and M. Zafrullah who proved the following characterization ([2, Corollary 5 and Theorem 6]).

Lemma 4.9. *For an integral domain R the following statements are equivalent:*

- (a) R is a generalized Cohen-Kaplansky domain.
- (b) \overline{R} is factorial, $R \subset \overline{R}$ is a root extension (i.e., for every $x \in \overline{R}$ there is some $k \in \mathbb{N}$ such that $x^k \in R$), $(R : \overline{R})$ is a principal ideal of \overline{R} , and $\overline{R}/(R : \overline{R})$ is finite.

If these conditions hold, then R is weakly factorial whence in particular a weakly Krull domain.

Lemma 4.10. *Let R be an archimedean domain such that \widehat{R} is factorial, and $\widehat{R}/(R : \widehat{R})$ is finite. Then R is a C-domain.*

Proof. Observe that $\{x^{-1}R \cap \widehat{R} \mid x \in \widehat{R}^\bullet\} \subset \{I \mid I \text{ is an } R\text{-submodule of } \widehat{R} \text{ with } (R : \widehat{R}) \subset I\}$. Therefore, $\{x^{-1}R^\bullet \cap \widehat{R}^\bullet \mid x \in \widehat{R}^\bullet\}$ is finite (since $\widehat{R}/(R : \widehat{R})$ is finite), and thus $\mathcal{C}(R^\bullet, \widehat{R}^\bullet)$ is finite by [19, Lemma 2.8.2.2]. Since $\widehat{R}^\times \cap R = R^\times$, we infer that R^\bullet is a C-monoid. \square

Proposition 4.11. *Let R be a generalized Cohen-Kaplansky domain and r a finitary ideal system on R .*

- 1. R is a C-domain and a Mori domain, and $v\text{-max}(R) = \mathfrak{X}(R)$.
- 2. If $T \subset R^\bullet$ a multiplicatively closed subset, then $T^{-1}R$ is a generalized Cohen-Kaplansky domain.

3. R is a Cohen-Kaplansky domain if and only if R is a G-domain. In particular, if $\mathfrak{p} \in \mathfrak{X}(R)$, then $R_{\mathfrak{p}}$ is a Cohen-Kaplansky domain.
4. If $r\text{-max}(R) = \mathfrak{X}(R)$, then R is r -Noetherian.

Proof. 1. Clearly, \overline{R} is factorial and $\overline{R}/(R : \overline{R})$ is finite by Lemma 4.9. Consequently, $\overline{R} = \widehat{R}$, $(R : \widehat{R}) \neq \{0\}$, and $\widehat{R}^{\times} \cap R = \overline{R}^{\times} \cap R = R^{\times}$. We infer by Lemma 4.10 that R is a C-domain, and hence R is a Mori domain by [19, Theorem 2.9.13]. It follows from Lemma 4.9 that R is a weakly Krull domain. Since R is a Mori domain, we infer that $v\text{-max}(R) = \mathfrak{X}(R)$ (note that $v\text{-max}(R) = \emptyset = \mathfrak{X}(R)$ if R is a field).

2. By Lemma 4.9, we know that \overline{R} is factorial, $R \subset \overline{R}$ is a root extension, $(R : \overline{R})$ is a principal ideal of \overline{R} and $\overline{R}/(R : \overline{R})$ is finite. Again by Lemma 4.9, it is sufficient to show that $T^{-1}\overline{R}$ is factorial, $T^{-1}R \subset T^{-1}\overline{R}$ is a root extension, $(T^{-1}R : T^{-1}\overline{R})$ is a principal ideal of $T^{-1}\overline{R}$ and $T^{-1}\overline{R}/(T^{-1}R : T^{-1}\overline{R})$ is finite.

It is clear that $T^{-1}\overline{R} = T^{-1}\overline{R}$ is factorial. Let $x \in T^{-1}\overline{R} = T^{-1}\overline{R}$. Then $tx \in \overline{R}$ for some $t \in T$, and hence $(tx)^k \in R$ for some $k \in \mathbb{N}$. This implies that $x^k \in T^{-1}R$. Therefore, $T^{-1}R \subset T^{-1}\overline{R}$ is a root extension. We have R is a Mori domain by 1, $(R : \overline{R}) \neq \{0\}$ and $\overline{R} = \widehat{R}$ is a fractional divisorial ideal of R . Therefore, $T^{-1}(R : \overline{R}) = (T^{-1}R : T^{-1}\overline{R}) = (T^{-1}R : T^{-1}\overline{R})$ by [19, Proposition 2.2.8.1]. Because of this it is clear that $(T^{-1}R : T^{-1}\overline{R})$ is a principal ideal of $T^{-1}\overline{R}$.

Finally, we have $T^{-1}\overline{R}/(T^{-1}R : T^{-1}\overline{R}) = T^{-1}\overline{R}/T^{-1}(R : \overline{R}) \cong (T + (R : \overline{R})/(R : \overline{R}))^{-1}(\overline{R}/(R : \overline{R}))$ is finite (since both $T + (R : \overline{R})/(R : \overline{R})$ and $\overline{R}/(R : \overline{R})$ are finite).

3. If R is a Cohen-Kaplansky domain, then R is semilocal and $\dim(R) \leq 1$ by [5, Theorem 4.3], and thus R is a G-domain. Now let R be a G-domain. Since R is also a Mori domain by 1, R has only finitely many prime ideals. Therefore, R has only finitely many prime elements up to associates, hence R has only finitely many atoms up to associates. We infer that R is a Cohen-Kaplansky domain.

Now let $\mathfrak{p} \in \mathfrak{X}(R)$. Then $R_{\mathfrak{p}}$ is a G-domain and it is a generalized Cohen-Kaplansky domain by 2. Therefore, $R_{\mathfrak{p}}$ is a Cohen-Kaplansky domain.

4. Let $r\text{-max}(R) = \mathfrak{X}(R)$. If $\mathfrak{m} \in \mathfrak{X}(R)$, then it follows by 3 that $R_{\mathfrak{m}}$ is a Cohen-Kaplansky domain, hence $R_{\mathfrak{m}}$ is Noetherian by [5, Theorem 4.3]. Let $(I_i)_{i \in \mathbb{N}}$ be an ascending chain of r -ideals of R . Without restriction let $I_1 \neq \{0\}$. Set $\mathcal{Q} = \{\mathfrak{m} \in \mathfrak{X}(R) \mid I_1 \subset \mathfrak{m}\}$. Then \mathcal{Q} is finite, and for all $\mathfrak{m} \in \mathfrak{X}(R)$ with $I_1 \not\subset \mathfrak{m}$ and all $\ell \in \mathbb{N}$ we have $(I_{\ell})_{\mathfrak{m}} = (I_1)_{\mathfrak{m}}$. Clearly, there is some $(n_{\mathfrak{m}})_{\mathfrak{m} \in \mathcal{Q}} \in \mathbb{N}^{\mathcal{Q}}$ such that for all $\mathfrak{m} \in \mathcal{Q}$ and $\ell \in \mathbb{N}_{\geq n_{\mathfrak{m}}}$ it follows that $(I_{\ell})_{\mathfrak{m}} = (I_{n_{\mathfrak{m}}})_{\mathfrak{m}}$. Set $n = \max(\{n_{\mathfrak{m}} \mid \mathfrak{m} \in \mathcal{Q}\} \cup \{1\})$. Obviously, $(I_{\ell})_{\mathfrak{m}} = (I_n)_{\mathfrak{m}}$ for all $\mathfrak{m} \in \mathfrak{X}(R)$ and $\ell \in \mathbb{N}_{\geq n}$. We infer that $I_{\ell} = I_n$ for all $\ell \in \mathbb{N}_{\geq n}$. \square

If R is an integral domain, r is an ideal system on R , and $\mathfrak{m} \in r\text{-max}(R)$, then let $\mathfrak{A}_{r,\mathfrak{m}}(R)$ be the set of all $I \in \mathcal{I}_r(R)$ with $\sqrt{I} = \mathfrak{m}$ such that for all $J, L \in \mathcal{I}_r(R)$ with $I = (JL)_r$ it follows that $J = R$ or $L = R$. Note that if $\mathcal{I}_r(R)$ is unit-cancellative, then $\mathfrak{A}_{r,\mathfrak{m}}(R)$ is the set of atoms of $\mathcal{I}_r(R)$ whose radical is equal to \mathfrak{m} .

Lemma 4.12. *Let R be a domain, r a finitary ideal system on R , and $\mathfrak{m} \in r\text{-max}(R)$. Let $\varphi: \mathfrak{A}_{r,\mathfrak{m}}(R) \rightarrow \mathfrak{A}_{r,\mathfrak{m},\mathfrak{m}}(R_{\mathfrak{m}})$ be defined by $\varphi(I) = I_{\mathfrak{m}}$ for all $I \in \mathfrak{A}_{r,\mathfrak{m}}(R)$ and let $\psi: \mathfrak{A}_{r,\mathfrak{m},\mathfrak{m}}(R_{\mathfrak{m}}) \rightarrow \mathfrak{A}_{r,\mathfrak{m}}(R)$ be defined by $\psi(J) = J \cap R$ for all $J \in \mathfrak{A}_{r,\mathfrak{m},\mathfrak{m}}(R_{\mathfrak{m}})$. Then φ and ψ are mutually inverse bijections.*

Proof. Let $f: \{I \in \mathcal{I}_r(R) \mid \sqrt{I} = \mathfrak{m}\} \rightarrow \{J \in \mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}}) \mid \sqrt{J} = \mathfrak{m}_{\mathfrak{m}}\}$ be defined by $f(I) = I_{\mathfrak{m}}$ for each $I \in \mathcal{I}_r(R)$ such that $\sqrt{I} = \mathfrak{m}$. Let $g: \{J \in \mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}}) \mid \sqrt{J} = \mathfrak{m}_{\mathfrak{m}}\} \rightarrow \{I \in \mathcal{I}_r(R) \mid \sqrt{I} = \mathfrak{m}\}$ be defined by $f(J) = J \cap R$ for each $J \in \mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}})$ such that $\sqrt{J} = \mathfrak{m}_{\mathfrak{m}}$. It is straightforward to prove that f and g are mutually inverse bijections. For instance, note that if $I \in \mathcal{I}_r(R)$ with $\sqrt{I} = \mathfrak{m}$, then I is \mathfrak{m} -primary, and hence $I_{\mathfrak{m}} \cap R = I$. To prove the assertion it is sufficient to show that $f(\mathfrak{A}_{r,\mathfrak{m}}(R)) = \mathfrak{A}_{r,\mathfrak{m},\mathfrak{m}}(R_{\mathfrak{m}})$.

First let $I \in \mathfrak{A}_{r,\mathfrak{m}}(R)$ and $J', L' \in \mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}})$ be such that $I_{\mathfrak{m}} = (J'L')_{r_{\mathfrak{m}}}$. Assume to the contrary that $J', L' \neq R_{\mathfrak{m}}$. Then $\sqrt{J'} = \sqrt{L'} = \mathfrak{m}_{\mathfrak{m}}$. Set $J = J' \cap R$ and $L = L' \cap R$. Then $J, L \in \mathcal{I}_r(R)$ and $\sqrt{J} = \sqrt{L} = \mathfrak{m}$, and hence $\sqrt{(JL)_r} = \mathfrak{m}$. Observe that $I_{\mathfrak{m}} = (J'L')_{r_{\mathfrak{m}}} = (J_{\mathfrak{m}}L_{\mathfrak{m}})_{r_{\mathfrak{m}}} = ((JL)_r)_{\mathfrak{m}}$. We infer that $I = (JL)_r$, and thus $J = R$ or $L = R$. Therefore, $J' = R_{\mathfrak{m}}$ or $L' = R_{\mathfrak{m}}$, a contradiction.

Next let $J \in \mathfrak{A}_{r_{\mathfrak{m}}, \mathfrak{m}_{\mathfrak{m}}}(R_{\mathfrak{m}})$ and set $I = J \cap R$. Let $A, B \in \mathcal{I}_r(R)$ be such that $I = (AB)_r$. Then $J = I_{\mathfrak{m}} = (A_{\mathfrak{m}}B_{\mathfrak{m}})_{r_{\mathfrak{m}}}$, and hence $A_{\mathfrak{m}} = R_{\mathfrak{m}}$ or $B_{\mathfrak{m}} = R_{\mathfrak{m}}$. It follows that $A = R$ or $B = R$ (since $\sqrt{A}, \sqrt{B} \supset \mathfrak{m}$). \square

Theorem 4.13. *Let R be a generalized Cohen-Kaplansky domain, and r an ideal system on R . Let $\mathcal{P} = \{\mathfrak{p} \in \mathfrak{X}(R) \mid \mathfrak{p} \not\supset (R : \widehat{R})\}$, $\mathcal{P}^* = \mathfrak{X}(R) \setminus \mathcal{P}$, and let T be the subsemigroup of $\mathcal{I}_r(R)$ generated by $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$. Suppose that one of the following conditions is satisfied:*

- (a) r is finitary and $r\text{-max}(R) = \mathfrak{X}(R)$.
- (b) $r = v$.
- (c) R is a G-domain.

Then R is r -Noetherian, $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^\}$ is finite, T is a finitely generated monoid, and $\mathcal{I}_r(R) \cong \mathcal{F}(\mathcal{P}) \times T$.*

Proof. CASE 1: r is finitary and $r\text{-max}(R) = \mathfrak{X}(R)$.

By Proposition 4.11.4 we have R is r -Noetherian. Clearly, $(R : \widehat{R}) \neq \{0\}$. Consequently, T is a monoid and $\mathcal{I}_r(R) \cong \mathcal{F}(\mathcal{P}) \times T$ by Theorem 4.8.1. Let $\mathfrak{m} \in \mathcal{P}^*$. Then $R_{\mathfrak{m}}$ is a Cohen-Kaplansky domain by Proposition 4.11.3. We infer by Proposition 4.5 that $\mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}})$ is finitely generated, hence $\mathcal{A}(\mathcal{I}_{r_{\mathfrak{m}}}(R_{\mathfrak{m}}))$ is finite. It follows by Lemma 4.12 that $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} = \mathfrak{m}\}$ is finite. Since \mathcal{P}^* is finite, we have $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \in \mathcal{P}^*\}$ is finite, and thus T is finitely generated by Theorem 4.8.2.

CASE 2: $r = v$.

By Proposition 4.11.1 we have R is a Mori domain and $v\text{-max}(R) = \mathfrak{X}(R)$. This implies that v is finitary, and hence the assertion follows from CASE 1.

CASE 3: R is a G-domain.

By Proposition 4.11.3 it follows that R is a Cohen-Kaplansky domain. Without restriction we may suppose that R is not a field. By [5, Theorem 4.3], R is one-dimensional Noetherian whence $r\text{-max}(R) = \mathfrak{X}(R)$ and R is r -Noetherian. Thus we are back to CASE 1. \square

We continue with an example of a domain R for which $\mathcal{I}_v(R)$ is not unit-cancellative (whence the statements of Theorem 4.13 do not hold) but R satisfies three of the four conditions in Lemma 4.9.(b) characterizing generalized Cohen-Kaplansky domains.

Example 4.14. There is a two-dimensional Noetherian domain R satisfying the following properties:

- \overline{R} is local Noetherian and factorial.
- $R_{\mathfrak{p}}$ is a discrete valuation domain for all $\mathfrak{p} \in \mathfrak{X}(R)$.
- $(R : \overline{R})$ is a nontrivial idempotent of $\mathcal{I}_v(R)$ and $\mathcal{I}_v(R)$ is not unit-cancellative.
- $\overline{R}/(R : \overline{R})$ is finite and $R \subset \overline{R}$ is a root extension.

Proof. Let S be a discrete valuation domain, and $d \in S^{\bullet} \setminus S^{\times}$ such that S/dS is finite. Let X be an indeterminate over S and $R = \{f \in S[[X]] \mid \text{the linear coefficient of } f \text{ is divisible by } d\}$. As shown in [31, Example 5.5.1], R is a two-dimensional Noetherian domain, \overline{R} is local Noetherian and factorial, $R_{\mathfrak{p}}$ is a discrete valuation domain for all $\mathfrak{p} \in \mathfrak{X}(R)$, and $(R : \overline{R})$ is a nontrivial idempotent of $\mathcal{I}_v(R)$. Clearly, $\mathcal{I}_v(R)$ is not unit-cancellative. Along the same lines as in the proof of [31, Lemma 5.3.3] it follows that $\overline{R}/(R : \overline{R})$ is finite. Set $k = |S/dS|$. It is straightforward to prove that $f^k \in R$ for all $f \in S[[X]] = \overline{R}$, and thus $R \subset \overline{R}$ is a root extension. \square

We end this section with our main arithmetical result. It combines the semigroup theoretical work in Section 3 with the ideal theoretic results of the present section.

Theorem 4.15. *Let R be a domain with $(R : \widehat{R}) \neq \{0\}$, r an r -Noetherian ideal system on R with $r\text{-max}(R) = \mathfrak{X}(R)$, and suppose that $\{I \in \mathcal{A}(\mathcal{I}_r(R)) \mid \sqrt{I} \supset (R : \widehat{R})\}$ is finite.*

1. $\mathcal{I}_r(R)$ has finite monotone catenary degree and finite successive distance. In particular, the catenary degree and the set of distances are finite.
2. $\mathcal{I}_r(R)$ satisfies the Structure Theorem for Unions.

Proof. We use all notation as in Theorem 4.8 and that $\mathcal{I}_r(R) \cong \mathcal{F}(\mathcal{P}) \times T$ where T is a finitely generated monoid.

1. Since $c_{\text{mon}}(\mathcal{F}(\mathcal{P}) \times T) = c_{\text{mon}}(T)$ and $\delta(\mathcal{F}(\mathcal{P}) \times T) = \delta(T)$, it suffices to prove the assertion for the monoid T . Theorem 3.1 implies that $\delta(T) < \infty$ and that $c_{\text{mon}}(T) < \infty$. Since $1 + \sup \Delta(T) \leq c(T) \leq c_{\text{mon}}(T)$, the remaining assertions follow.

2. It can be seen from the definitions that

$$\Delta(\mathcal{F}(\mathcal{P}) \times T) = \Delta(T) \quad \text{and} \quad \rho_k(\mathcal{F}(\mathcal{P}) \times T) = \rho_k(T)$$

for every $k \in \mathbb{N}$. By [10, Proposition 3.4], there exists a constant $M \in \mathbb{N}$ such that $\rho_{k+1}(T) \leq \rho_k(T) + M$ for all $k \in \mathbb{N}$ whence $\rho_{k+1}(\mathcal{F}(\mathcal{P}) \times T) \leq \rho_k(\mathcal{F}(\mathcal{P}) \times T) + M$ for all $k \in \mathbb{N}$. Since $\Delta(T)$ is finite by 1, it follows that $\Delta(\mathcal{F}(\mathcal{P}) \times T)$ is finite. Therefore, all assumptions of [10, Theorem 2.2] are satisfied whence $\mathcal{F}(\mathcal{P}) \times T$ satisfies the Structure Theorem for Unions. \square

5. MONOIDS OF v -INVERTIBLE v -IDEALS IN WEAKLY KRULL MONOIDS

Weakly Krull domains were introduced by Anderson, Anderson, Mott, and Zafrullah [2, 4]. Halter-Koch gave a divisor-theoretic characterization and showed that a domain is weakly Krull if and only if its multiplicative monoid of non-zero elements is a weakly Krull monoid ([25]). We will restrict to the setting of v -Noetherian monoids and domains and recall that a (commutative cancellative) v -Noetherian monoid H is weakly Krull if and only if $v\text{-max}(H) = \mathfrak{X}(H)$ ([26, Theorem 24.5]). Thus one-dimensional Mori domains are weakly Krull, and by Proposition 4.11 generalized Cohen-Kaplansky domains are weakly Krull Mori domains.

In this section we study the monotone catenary degree of the monoid $\mathcal{I}_v^*(H)$ of v -invertible v -ideals, where H is a weakly Krull Mori monoid with nonempty conductor $(H : \widehat{H})$. This monoid is a direct product of a free abelian part and of finitely many finitely primary monoids (see (5.2)). The seminormal case has already been studied in detail. Indeed, if H is a seminormal v -Noetherian weakly Krull monoid with proper nonempty conductor such that $H_{\mathfrak{p}}$ is finitely primary for all $\mathfrak{p} \in \mathfrak{X}(H)$, then the monotone catenary degree of $\mathcal{I}_v^*(H)$ is either 2, 3, or 5, and it is well-understood which case occurs ([21, Theorem 5.8]).

However, in general and even in the local case (thus for finitely primary monoids), the monotone catenary degree may be infinite (Remark 5.3). We study a special class of finitely primary monoids, called strongly ring-like, which was introduced by Hassler in [28]. Strongly ring-like monoids of rank at most two (the restriction on the rank is essential, as outlined in Remark 5.3) have finite monotone catenary degree (Proposition 5.12) and the same is true for $\mathcal{I}_v^*(H)$ provided that the localizations $H_{\mathfrak{p}}$ are strongly ring-like of rank at most two (Theorem 5.13).

We begin with the local case. A monoid H is said to be *finitely primary* if there are $s, \alpha \in \mathbb{N}$ and a factorial monoid $F = F^\times \times \mathcal{F}(\{p_1, \dots, p_s\})$ such that $H \subset F$ with

$$(5.1) \quad H \setminus H^\times \subset p_1 \cdots p_s F \quad \text{and} \quad (p_1 \cdots p_s)^\alpha F \subset H.$$

Let $H \subset F$ be finitely primary as above. Then s is called the rank of H , α is called an exponent of H , and $\mathbf{v}: H \rightarrow \mathbb{N}_0^s$, defined by $(a \mapsto (\mathbf{v}_{p_1}(a), \dots, \mathbf{v}_{p_s}(a)))$ for all $a \in H$, denotes the map from H to its value semigroup $\mathbf{v}(H)$. It is well-known ([19, Theorems 2.9.2 and 3.1.5]) that H is primary, that F is the complete integral closure of H , and that $s = |\mathfrak{X}(\widehat{H})|$. Clearly, every finitely primary monoid is strongly primary.

The following two lemmas gather the main arithmetical properties of finitely primary monoids and the connection to ring theory.

Lemma 5.1. *Let H be a finitely primary monoid. Then H has finite catenary degree, finite set of distances, and it satisfies the Structure Theorem for Sets of Lengths. Moreover, if the rank of H is greater than one, then H is not half-factorial.*

Proof. By [19, Corollary 4.5.5] we have H satisfies the Structure Theorem for Sets of Lengths. It follows from [19, Theorems 2.9.2.4, 3.1.5.2, 3.1.1.2] that $\Delta(H)$ is finite. \square

Lemma 5.2. *Let R be a domain.*

1. R^\bullet is finitely primary if and only if R is one-dimensional local, $(R:\widehat{R}) \neq \{0\}$, and \widehat{R} is a semilocal principal ideal domain.
2. If R is a one-dimensional local Mori domain such that $(R:\widehat{R}) \neq \{0\}$, then R^\bullet is finitely primary of rank $|\mathfrak{X}(\widehat{R})|$.

Proof. See [19, Proposition 2.10.7]. \square

Remark 5.3.

1. Let $H \subset F$ be a finitely primary monoid as in (5.1). By [19, Corollary 2.9.8], H is a C-monoid if and only if the following two conditions are fulfilled:

- (a) There exists a subgroup $V \subset \widehat{H}^\times$ of finite index such that $V(H \setminus H^\times) \subset H$.
- (b) There exists some $\alpha \in \mathbb{N}$ such that, for every $j \in [1, s]$ and $a \in p_j^\alpha \widehat{H}$, we have $a \in H$ if and only if $p_j^\alpha a \in H$.

In general, finitely primary monoids need neither be v -Noetherian nor C-monoids (see [27, 32]).

2. Let $H \subset F$ be a finitely primary monoid as above. Then H_{red} is finitely generated if and only if $s = 1$ and $(F^\times : H^\times) < \infty$ ([19, Theorem 2.9.2]). If H_{red} is finitely generated, then $\delta_w(H) \leq \delta(H) < \infty$ and $c_{\text{mon}}(H) < \infty$ by Theorem 3.1.

3. In contrast to Lemma 5.1, there is a finitely primary monoid of rank one with $c_{\text{mon}}(H) = \delta(H) = \infty$. Furthermore, there are finitely primary monoids of rank two and exponent two having infinite monotone catenary degree (see [13, Remark 4.6, Examples 4.5 and 4.16]).

4. There are one-dimensional local Noetherian domains R with maximal ideal \mathfrak{m} such that:

- \overline{R} is a finitely generated R -module and R/\mathfrak{m} is finite.
- $(\overline{R}$ has 2 maximal ideals and $\delta(R^\bullet) = \infty$) or $(\overline{R}$ has 3 maximal ideals and $c_{\text{mon}}(R^\bullet) = \infty)$.

([28, Examples 6.3 and 6.5]). By Lemma 5.2, R^\bullet is finitely primary of rank two or three.

The examples discussed in Remark 5.3 show that finitely primary monoids need to satisfy further structural properties if we want their monotone catenary degree to be finite. Such properties were introduced by Hassler in [28], and we recall the definition.

Definition 5.4. Let H be a finitely primary monoid of rank $s \in \mathbb{N}$ such that there exist some exponent $\alpha \in \mathbb{N}$ of H and some system $\{p_i \mid i \in [1, s]\}$ of representatives of the prime elements of \widehat{H} with the following property: for all $i \in [1, s]$ and for all $a \in \widehat{H}$ with $v_{p_i}(a) \geq \alpha$ we have $p_i a \in H$ if and only if $a \in H$. Then H is said to be

- *ring-like* if $\widehat{H}^\times/H^\times$ is finite or $\{(v_{p_i}(a))_{i=1}^s \mid a \in H \setminus H^\times\} \subset \mathbb{N}^s$ has a smallest element with respect to the partial order.
- *strongly ring-like* if $\widehat{H}^\times/H^\times$ is finite and $\{(v_{p_i}(a))_{i=1}^s \mid a \in H \setminus H^\times\} \subset \mathbb{N}^s$ has a smallest element with respect to the partial order.

Let H be a ring-like monoid of rank s and exponent α and $\{p_i \mid i \in [1, s]\}$ a system of representatives of the prime elements of \widehat{H} . We say that α and $\{p_i \mid i \in [1, s]\}$ are suitably chosen if for all $i \in [1, s]$ and for all $a \in \widehat{H}$ with $v_{p_i}(a) \geq \alpha$ we have $p_i a \in H$ if and only if $a \in H$.

We continue with a ring theoretical analysis which one-dimensional local domains are strongly ring-like (Propositions 5.6 and 5.9). The characterization mentioned in Remark 5.3.1 shows that strongly ring-like monoids are C-monoids and hence in particular they are v -Noetherian.

Lemma 5.5. *Let S be a commutative ring, $a, b \in S$, and $\mathcal{R} \subset S$ such that $x - y \in S^\times$ for all distinct $x, y \in \mathcal{R}$. If \mathfrak{J} is a finite set of ideals of S such that $|\mathfrak{J}| < |\mathcal{R}|$ and $b \notin I$ for all $I \in \mathfrak{J}$, then there is some $\eta \in \mathcal{R}$ such that $a + \eta b \notin \bigcup_{I \in \mathfrak{J}} I$.*

Proof. It is sufficient to show by induction that for all $k \in \mathbb{N}_0$ and all sets of ideals \mathfrak{J} of S such that $k = |\mathfrak{J}| < |\mathcal{R}|$ and $b \notin I$ for all $I \in \mathfrak{J}$ it follows that there is some $\eta \in \mathcal{R}$ such that $a + \eta b \notin \bigcup_{I \in \mathfrak{J}} I$.

The assertion is clear for $k = 0$. Now let $k \in \mathbb{N}_0$ and \mathfrak{J} a set of ideals of S be such that $k+1 = |\mathfrak{J}| < |\mathcal{R}|$ and $b \notin I$ for all $I \in \mathfrak{J}$. By the induction hypothesis there is some $(\eta_I)_{I \in \mathfrak{J}} \in \mathcal{R}^{\mathfrak{J}}$ such that for all $I \in \mathfrak{J}$ it follows that $a + \eta_I b \notin \bigcup_{J \in \mathfrak{J} \setminus \{I\}} J$.

CASE 1: $a + \eta_J b \notin J$ for some $J \in \mathfrak{J}$.

Then $a + \eta_J b \notin \bigcup_{I \in \mathfrak{J}} I$.

CASE 2: $a + \eta_I b \in I$ for all $I \in \mathfrak{J}$.

There is some $\eta \in \mathcal{R} \setminus \{\eta_I \mid I \in \mathfrak{J}\}$. Assume that $a + \eta b \in J$ for some $J \in \mathfrak{J}$. Then $(\eta - \eta_J)b = a + \eta b - (a + \eta_J b) \in J$, hence $b \in J$, a contradiction. Therefore, $a + \eta b \notin \bigcup_{I \in \mathfrak{J}} I$. \square

If R is a domain such that R^\bullet is finitely primary and $\mathfrak{m} = R \setminus R^\times$, then set $V(\mathfrak{m}^\bullet) = \{(v_{\mathfrak{q}}(a))_{\mathfrak{q} \in \max(\widehat{R})} \mid a \in \mathfrak{m}^\bullet\}$.

Proposition 5.6. *Let R be a domain such that R^\bullet is finitely primary and $\mathfrak{m} = R \setminus R^\times$.*

1. *If $|\max(\widehat{R})| \leq |R/\mathfrak{m}|$, then $V(\mathfrak{m}^\bullet)$ has a smallest element.*
2. *The following statements are equivalent:*
 - (a) $\widehat{R}^\times/R^\times$ is finite.
 - (b) R is a discrete valuation domain or (R is Noetherian and $|R/\mathfrak{m}| < \infty$).
 - (c) R is a discrete valuation domain or (\widehat{R} is a finitely generated R -module and $|R/\mathfrak{m}| < \infty$).
 - (d) R is an FF-domain.
3. *There are some $\alpha, s \in \mathbb{N}$ and some system $\{p_i \mid i \in [1, s]\}$ of representatives of prime elements of \widehat{R} such that for all $i \in [1, s]$ and $a \in \widehat{R}$ with $v_{p_i}(a) \geq \alpha$ it follows that $p_i a \in R$ if and only if $a \in R$.*

Proof. 1. Let $|\max(\widehat{R})| \leq |R/\mathfrak{m}|$. By [19, Theorem 1.5.3] it suffices to show that $|\text{Min}(V(\mathfrak{m}^\bullet))| \leq 1$. Assume to the contrary that there are distinct $x, y \in \text{Min}(V(\mathfrak{m}^\bullet))$. There are some $e, f \in \mathfrak{m}^\bullet$ such that $x = (v_{\mathfrak{q}}(e))_{\mathfrak{q} \in \max(\widehat{R})}$ and $y = (v_{\mathfrak{q}}(f))_{\mathfrak{q} \in \max(\widehat{R})}$. Since \widehat{R} is a principal ideal domain, there is some $d \in \text{GCD}_{\widehat{R}}(e, f)$. Set $a = \frac{e}{d}$ and $b = \frac{f}{d}$. There is some $\mathcal{R} \subset R^\times$ such that $|\mathcal{R}| = |R/\mathfrak{m}| - 1$ and $v - w \in R^\times \subset \widehat{R}^\times$ for all distinct $v, w \in \mathcal{R}$. Set $\mathfrak{J} = \{\mathfrak{q} \in \max(\widehat{R}) \mid v_{\mathfrak{q}}(e) = v_{\mathfrak{q}}(f)\}$. We have $\mathfrak{J} \subset \{\mathfrak{q} \in \max(\widehat{R}) \mid b \notin \mathfrak{q}\}$. Since $x, y \in \text{Min}(V(\mathfrak{m}^\bullet))$ are distinct it follows that $|\mathfrak{J}| < |\max(\widehat{R})| - 1 \leq |R/\mathfrak{m}| - 1 = |\mathcal{R}|$. By Lemma 5.5 there is some $\eta \in \mathcal{R}$ such that $a + \eta b \notin \bigcup_{\mathfrak{q} \in \mathfrak{J}} \mathfrak{q}$.

Next we show that $v_{\mathfrak{n}}(e + \eta f) = \min\{v_{\mathfrak{n}}(e), v_{\mathfrak{n}}(f)\}$ for all $\mathfrak{n} \in \max(\widehat{R})$. Let $\mathfrak{n} \in \max(\widehat{R})$.

CASE 1: $v_{\mathfrak{n}}(e) \neq v_{\mathfrak{n}}(f)$.

Since $v_{\mathfrak{n}}(e) \neq v_{\mathfrak{n}}(\eta f)$ it follows that $v_{\mathfrak{n}}(e + \eta f) = \min\{v_{\mathfrak{n}}(e), v_{\mathfrak{n}}(\eta f)\} = \min\{v_{\mathfrak{n}}(e), v_{\mathfrak{n}}(f)\}$.

CASE 2: $v_{\mathfrak{n}}(e) = v_{\mathfrak{n}}(f)$.

Since $\mathfrak{n} \in \mathfrak{J}$ we obtain that $a + \eta b \notin \mathfrak{n}$. Consequently, $v_{\mathfrak{n}}(e + \eta f) = v_{\mathfrak{n}}(d(a + \eta b)) = v_{\mathfrak{n}}(d) + v_{\mathfrak{n}}(a + \eta b) = v_{\mathfrak{n}}(d) = \min\{v_{\mathfrak{n}}(e), v_{\mathfrak{n}}(f)\}$.

Since $(v_{\mathfrak{q}}(e + \eta f))_{\mathfrak{q} \in \max(\widehat{R})} \in V(\mathfrak{m}^\bullet)$ we infer that $x = (v_{\mathfrak{q}}(e + \eta f))_{\mathfrak{q} \in \max(\widehat{R})} = y$, a contradiction.

2. (a) \Rightarrow (b) Let $\widehat{R}^\times/R^\times$ be finite and R not a discrete valuation domain. It follows from [32, Theorems 4.2 and 4.3] that R is Noetherian and $\widehat{R}/(R : \widehat{R})$ is finite. Since R is not a discrete valuation domain we have $(R : \widehat{R}) \subset \mathfrak{m}$, and thus $|R/\mathfrak{m}| < \infty$.

(b) \Rightarrow (c) This is clear, since $(R : \widehat{R}) \neq \{0\}$.

(c) \Rightarrow (a) The assertion is clear if R is a discrete valuation domain. Now let \widehat{R} be a finitely generated R -module and $|R/\mathfrak{m}| < \infty$. We have \widehat{R}/\mathfrak{q} is a finite-dimensional R/\mathfrak{m} -vector space for all $\mathfrak{q} \in \max(\widehat{R})$, hence \widehat{R}/\mathfrak{q} is finite for all $\mathfrak{q} \in \max(\widehat{R})$. Since \widehat{R} is a principal ideal domain we infer that \widehat{R}/I is finite for every nonzero ideal I of \widehat{R} . Consequently, $\widehat{R}/(R : \widehat{R})$ is finite, and thus \widehat{R}/R is finite. Since R is local it follows from [32, Proposition 3.5] that $\widehat{R}^\times/R^\times$ is finite.

(a) \Leftrightarrow (d) This is an immediate consequence of [19, Theorem 2.9.2.4].

3. This follows from [27, Theorem 2.7]. \square

Corollary 5.7. *Let (R, \mathfrak{m}) be a one-dimensional local domain with $(R : \widehat{R}) \neq \{0\}$.*

1. *If \widehat{R} is a semilocal principal ideal domain and $|\max(\widehat{R})| \leq |R/\mathfrak{m}|$, then R^\bullet is ring-like.*
2. *If R is Noetherian, then R^\bullet is ring-like.*
3. *If R is Noetherian and $|\max(\widehat{R})| \leq |R/\mathfrak{m}| < \infty$, then R^\bullet is strongly ring-like.*

Proof. 1. This follows from Proposition 5.6.

2. By Lemma 5.2, R^\bullet is a finitely primary monoid. By Proposition 5.6.3 it is sufficient to show that $\widehat{R}^\times/R^\times$ is finite or $V(\mathfrak{m}^\bullet)$ has a smallest element. Let $\widehat{R}^\times/R^\times$ be infinite. It follows from Proposition 5.6.2 that R/\mathfrak{m} is infinite. Since \widehat{R} is semilocal we have $|\max(\widehat{R})| \leq |R/\mathfrak{m}|$, and thus $V(\mathfrak{m}^\bullet)$ has a smallest element by Proposition 5.6.1.

3. It is an immediate consequence of Proposition 5.6 that R^\bullet is strongly ring-like. \square

The next example shows that Corollary 5.7.3 does not hold true for Mori domains.

Example 5.8. There are one-dimensional local seminormal Mori domains R with $(R : \widehat{R}) \neq \{0\}$ satisfying the following properties:

- \widehat{R} is a discrete valuation domain,
- $|\max(\widehat{R})| \leq |R/\mathfrak{m}| < \infty$ and $\widehat{R}^\times/R^\times$ is infinite,
- R is integrally closed or $\overline{R} = \widehat{R}$,
- R^\bullet is finitely primary but not strongly ring-like.

Proof. Let K be a finite field and L an infinite extension field of K . Let X be an indeterminate over L . Set $R = K + XL[[X]]$ and $\mathfrak{m} = XL[[X]]$. Then \mathfrak{m} is the unique nonzero prime ideal of R . In particular, R is one-dimensional and local. Clearly, $\widehat{R} = L[[X]]$ is a discrete valuation domain. Observe that R is seminormal, hence R is a Mori domain and $(R : \widehat{R}) \neq \{0\}$. We obtain that R^\bullet is finitely primary. Observe that $R/\mathfrak{m} \cong K$, and thus $|\max(\widehat{R})| = 1 \leq |R/\mathfrak{m}| = |K| < \infty$. Since K is finite and L is infinite, it follows that $\widehat{R}^\times/R^\times \cong L^\times/K^\times$ is infinite. Therefore, R^\bullet is not strongly ring-like.

CASE 1: L/K is purely transcendental (e.g., $L = K(Y)$ for some indeterminate Y over K). It is easy to prove that R is integrally closed.

CASE 2: L/K is an algebraic field extension (e.g., L is an algebraic closure of K). Observe that $\overline{R} = \widehat{R}$. \square

Proposition 5.9. *Let R be a Mori domain such that $(R : \widehat{R}) \neq \{0\}$ and $\mathfrak{p} \in \mathfrak{X}(R)$.*

1. $R_{\mathfrak{p}}^\bullet$ is a finitely primary monoid.

2. For all $\mathfrak{m} \in \text{spec}(\widehat{R})$ with $\mathfrak{m} \cap R = \mathfrak{p}$ it follows that $\mathfrak{m} \in \mathfrak{X}(\widehat{R})$. In particular, $\{\mathfrak{m} \in \text{spec}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\} = \{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}$.
3. If $\mathfrak{p} \supset (R : \widehat{R})$, then $R_{\mathfrak{p}}^{\bullet}$ is strongly ring-like of rank at most two if and only if $R_{\mathfrak{p}}$ is Noetherian, $|R/\mathfrak{p}| < \infty$ and $|\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}| \leq 2$.

Proof. Clearly, $R_{\mathfrak{p}}$ is a one-dimensional local Mori domain, $\widehat{R}_{\mathfrak{p}} = \widehat{R}_{\mathfrak{p}}$ (since \widehat{R} is a Krull domain) and $\{0\} \neq (R : \widehat{R}) \subset (R_{\mathfrak{p}} : \widehat{R}_{\mathfrak{p}})$. If $\mathfrak{m} \in \text{spec}(\widehat{R})$, then set $\mathfrak{m}_{\mathfrak{p}} = (R \setminus \mathfrak{p})^{-1}\mathfrak{m}$.

1. It follows from Lemma 5.2.1 that $R_{\mathfrak{p}}^{\bullet}$ is finitely primary.

2. Let $\mathfrak{m} \in \text{spec}(\widehat{R})$ with $\mathfrak{m} \cap R = \mathfrak{p}$. Since \widehat{R} is a Krull domain, there is some $\mathfrak{q} \in \mathfrak{X}(\widehat{R})$ such that $\mathfrak{q} \subset \mathfrak{m}$. This implies that $\{0\} \neq \mathfrak{q}_{\mathfrak{p}} \subset \mathfrak{m}_{\mathfrak{p}}$. Since $\widehat{R}_{\mathfrak{p}}$ is a semilocal principal ideal domain it follows that $\mathfrak{q}_{\mathfrak{p}} = \mathfrak{m}_{\mathfrak{p}}$, and thus $\mathfrak{q} = \mathfrak{m}$. We infer that $\mathfrak{m} \in \mathfrak{X}(\widehat{R})$.

3. Let $\mathfrak{p} \supset (R : \widehat{R})$. Note that $\widehat{R}_{\mathfrak{p}}$ is a semilocal principal ideal domain. We infer by 2 that $\mathfrak{X}(\widehat{R}_{\mathfrak{p}}) = \text{spec}(\widehat{R}_{\mathfrak{p}}) \setminus \{(0)\} = \text{spec}(\widehat{R}_{\mathfrak{p}}) \setminus \{(0)\} = \{\mathfrak{m}_{\mathfrak{p}} \mid \mathfrak{m} \in \text{spec}(\widehat{R}) \setminus \{(0)\}, \mathfrak{m} \cap R \subset \mathfrak{p}\} = \{\mathfrak{m}_{\mathfrak{p}} \mid \mathfrak{m} \in \text{spec}(\widehat{R}), \mathfrak{m} \cap R = \mathfrak{p}\} = \{\mathfrak{m}_{\mathfrak{p}} \mid \mathfrak{m} \in \mathfrak{X}(\widehat{R}), \mathfrak{m} \cap R = \mathfrak{p}\}$. Note that $\varphi : \{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\} \rightarrow \{\mathfrak{m}_{\mathfrak{p}} \mid \mathfrak{m} \in \mathfrak{X}(\widehat{R}), \mathfrak{m} \cap R = \mathfrak{p}\}$ defined by $\varphi(\mathfrak{m}) = \mathfrak{m}_{\mathfrak{p}}$ for all $\mathfrak{m} \in \mathfrak{X}(\widehat{R})$ with $\mathfrak{m} \cap R = \mathfrak{p}$ is a bijection. Consequently, $|\mathfrak{X}(\widehat{R}_{\mathfrak{p}})| = |\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}|$.

First suppose that $R_{\mathfrak{p}}^{\bullet}$ is strongly ring-like of rank at most two. Then $\widehat{R}_{\mathfrak{p}}^{\times}/R_{\mathfrak{p}}^{\times}$ is finite. It follows by [19, Theorem 2.6.5.3] that $R_{\mathfrak{p}}$ is not a discrete valuation domain. Therefore, we obtain by Proposition 5.6 that $R_{\mathfrak{p}}$ is Noetherian and $|R_{\mathfrak{p}}/\mathfrak{p}_{\mathfrak{p}}| < \infty$. Since $f : R/\mathfrak{p} \rightarrow R_{\mathfrak{p}}/\mathfrak{p}_{\mathfrak{p}}$ defined by $f(x + \mathfrak{p}) = x + \mathfrak{p}_{\mathfrak{p}}$ for all $x \in R$ is a ring monomorphism, we have $|R/\mathfrak{p}| < \infty$. We infer that $|\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}| = |\mathfrak{X}(\widehat{R}_{\mathfrak{p}})| = |\mathfrak{X}(\widehat{R}_{\mathfrak{p}}^{\bullet})| \leq 2$.

Conversely suppose that $R_{\mathfrak{p}}$ is Noetherian, $|R/\mathfrak{p}| < \infty$ and $|\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}| \leq 2$. Then \mathfrak{p} is a maximal ideal of R whence $R/\mathfrak{p} \cong R_{\mathfrak{p}}/\mathfrak{p}_{\mathfrak{p}}$. Thus we obtain that have

$$|\max(\widehat{R}_{\mathfrak{p}})| = |\mathfrak{X}(\widehat{R}_{\mathfrak{p}}^{\bullet})| = |\mathfrak{X}(\widehat{R}_{\mathfrak{p}})| = |\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}| \leq 2 \leq |R_{\mathfrak{p}}/\mathfrak{p}_{\mathfrak{p}}| = |R/\mathfrak{p}| < \infty.$$

Therefore, it follows from Corollary 5.7 that $R_{\mathfrak{p}}^{\bullet}$ is strongly ring-like. Clearly, $R_{\mathfrak{p}}^{\bullet}$ is of rank at most two. \square

Our next goal is to show that for strongly ring-like monoids of rank at most two the weak successive distance $\delta_w(\cdot)$ is finite (see Proposition 5.12).

Lemma 5.10. *Let H be a ring-like monoid of rank s and exponent α and $\{p_i \mid i \in [1, s]\}$ a system of representatives of the prime elements of \widehat{H} such that α and $\{p_i \mid i \in [1, s]\}$ are suitably chosen and such that either $s \geq 2$ or $\widehat{H}^{\times}/H^{\times}$ is finite.*

1. If $u \in H$, $i \in [1, s]$ and $\nu_{p_i}(u) \geq 2\alpha$, then $u \in \mathcal{A}(H)$ if and only if $p_i u \in \mathcal{A}(H)$.
2. There are some $M_1, C_1 \in \mathbb{N}$ such that for each $a \in H$ and all adjacent $k, \ell \in \mathbb{L}(a)$ such that $\max\{k, \ell\} + M_1 \leq \max \mathbb{L}(a)$ it follows that $\text{Dist}(Z_k(a), Z_{\ell}(a)) \leq C_1$.

Proof. 1. This follows from [28, Proposition 4.4].

2. We distinguish two cases.

CASE 1: $s = 1$.

Then H_{red} is finitely generated by Remark 5.3.2. Thus Theorem 3.1 implies that $\delta(H) < \infty$ whence the assertion follows.

CASE 2: $s \geq 2$.

We infer by [19, Theorem 3.1.5] that $\min \mathbb{L}(a) \leq 2\alpha$ for all $a \in H$. It follows by [28, Theorem 4.14] that there are some $M_1, C_2 \in \mathbb{N}$ such that for all $a \in H$ and all adjacent $k, \ell \in \mathbb{L}(a)$ for which $\min \mathbb{L}(a) + M_1 \leq \min\{k, \ell\} \leq \max\{k, \ell\} \leq \max \mathbb{L}(a) - M_1$ it follows that $\text{Dist}(Z_k(a), Z_{\ell}(a)) \leq C_2$. By

Lemma 5.1, $\Delta(H)$ is finite and nonempty. Set $C_1 = \max\{C_2, M_1 + 2\alpha + \max\Delta(H)\}$. Let $a \in H$ and $k, \ell \in \mathbf{L}(a)$ be adjacent such that $\max\{k, \ell\} + M_1 \leq \max\mathbf{L}(a)$.

CASE 2.1: $\min\{k, \ell\} \geq M_1 + 2\alpha$. Obviously, $\min\mathbf{L}(a) + M_1 \leq \min\{k, \ell\}$, and thus $\text{Dist}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq C_2 \leq C_1$.

CASE 2.2: $\min\{k, \ell\} < M_1 + 2\alpha$. It is easy to see that $\text{Dist}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq \max\{k, \ell\} \leq \min\{k, \ell\} + \max\Delta(H) \leq M_1 + 2\alpha + \max\Delta(H) \leq C_1$. \square

Lemma 5.11. *Let H be a strongly ring-like monoid of rank 2 and exponent α , $\{p_1, p_2\}$ a system of representatives of the prime elements of \hat{H} such that α and $\{p_1, p_2\}$ are suitably chosen and (μ_1, μ_2) is the smallest element of $\{(\mathbf{v}_{p_1}(a), \mathbf{v}_{p_2}(a)) \mid a \in H \setminus H^\times\}$. Set $\mathbb{A}(H) = \{q \in \mathcal{A}(H) \mid \mathbf{v}_{p_1}(q) \leq 2\alpha, \mathbf{v}_{p_2}(q) \leq 2\alpha\}$.*

1. For every $a \in H$ we have $\min\{\mathbf{v}_{p_i}(a) - \mu_i \max\mathbf{L}(a) \mid i \in \{1, 2\}\} < \alpha$.
2. There are $L, E \in \mathbb{N}$ such that for all $u \in \mathbb{A}(H)$, $r \in \mathbb{N}$ and $(u_i)_{i=1}^r \in \mathcal{A}(H)^r$ with $\mathbf{v}_{p_1}(u) = \mu_1$, $|\{i \in [1, r] \mid \mathbf{v}_{p_1}(u_i) = \mu_1\}| > L$ and $\max\{\mathbf{v}_{p_2}(u_i) \mid i \in [1, r]\} > E$ it follows that $u \mid_H \prod_{i=1}^r u_i$ and $r - 1 \in \mathbf{L}(u^{-1} \prod_{i=1}^r u_i)$.
3. For every $N \in \mathbb{N}$ there is some $D \in \mathbb{N}$ such that for all $a \in H$ and $k, \ell \in \mathbf{L}(a)$ for which $\min\{k, \ell\} + N \geq \max\mathbf{L}(a)$ it follows that $\text{d}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq D|\ell - k|$.

Proof. 1. This follows from [28, Lemma 5.1].

2. It follows from [28, Lemma 5.2] that there are some $L, E \in \mathbb{N}$ such that for all $u \in \mathbb{A}(H)$, $(u_i)_{i=1}^L \in \mathcal{A}(H)^L$ with $\mathbf{v}_{p_1}(u) = \mu_1$, $\mathbf{v}_{p_1}(u_i) = \mu_1$ for all $i \in [1, L]$ and $v \in \mathcal{A}(H)$ with $\mathbf{v}_{p_2}(v) \geq E$ it follows that $u \prod_{i=1}^L w_i = v \prod_{i=1}^L u_i$ for some $(w_i)_{i=1}^L \in \mathcal{A}(H)^L$. Now let $u \in \mathbb{A}(H)$, $r \in \mathbb{N}$ and $(u_i)_{i=1}^r \in \mathcal{A}(H)^r$ be such that $\mathbf{v}_{p_1}(u) = \mu_1$, $|\{i \in [1, r] \mid \mathbf{v}_{p_1}(u_i) = \mu_1\}| > L$ and $\max\{\mathbf{v}_{p_2}(u_i) \mid i \in [1, r]\} > E$. There are some $j \in [1, r]$ such that $\mathbf{v}_{p_2}(u_j) = \max\{\mathbf{v}_{p_2}(u_i) \mid i \in [1, r]\}$ and $I \subset [1, r] \setminus \{j\}$ such that $|I| = L$ and $\mathbf{v}_{p_1}(u_i) = \mu_1$ for all $i \in I$. Consequently, there is some $(w_i)_{i=1}^L \in \mathcal{A}(H)^L$ such that $u \prod_{i=1}^L w_i = u_j \prod_{i \in I} u_i$. This implies that $\prod_{i=1}^r u_i = u \prod_{i=1}^L w_i \prod_{i \in [1, r] \setminus (I \cup \{j\})} u_i$, and thus $u \mid_H \prod_{i=1}^r u_i$ and $r - 1 = L + r - (L + 1) \in \mathbf{L}(u^{-1} \prod_{i=1}^r u_i)$.

3. Let $N \in \mathbb{N}$. Without restriction let H be reduced. By 2 there are some $L_1, E_1 \in \mathbb{N}$ such that for all $u \in \mathbb{A}(H)$, $r \in \mathbb{N}$ and $(u_i)_{i=1}^r \in \mathcal{A}(H)^r$ with $\mathbf{v}_{p_1}(u) = \mu_1$, $|\{i \in [1, r] \mid \mathbf{v}_{p_1}(u_i) = \mu_1\}| > L_1$ and $\max\{\mathbf{v}_{p_2}(u_i) \mid i \in [1, r]\} > E_1$ it follows that $u \mid_H \prod_{i=1}^r u_i$ and $r - 1 \in \mathbf{L}(u^{-1} \prod_{i=1}^r u_i)$.

Analogously, it follows by 2 that there are some $L_2, E_2 \in \mathbb{N}$ such that for all $u \in \mathbb{A}(H)$, $r \in \mathbb{N}$ and $(u_i)_{i=1}^r \in \mathcal{A}(H)^r$ with $\mathbf{v}_{p_2}(u) = \mu_2$, $|\{i \in [1, r] \mid \mathbf{v}_{p_2}(u_i) = \mu_2\}| > L_2$ and $\max\{\mathbf{v}_{p_1}(u_i) \mid i \in [1, r]\} > E_2$ it follows that $u \mid_H \prod_{i=1}^r u_i$ and $r - 1 \in \mathbf{L}(u^{-1} \prod_{i=1}^r u_i)$.

Set $E = \max\{E_1, E_2, \mu_1(N + 1) + \alpha, \mu_2(N + 1) + \alpha\}$ and $\mathbb{B} = \{q \in \mathcal{A}(H) \mid \mathbf{v}_{p_1}(q) \leq E, \mathbf{v}_{p_2}(q) \leq E\}$. Let $H(\mathbb{B})$ be the submonoid of H generated by \mathbb{B} . Note that $\mathcal{A}(H(\mathbb{B})) = \mathbb{B}$. Since \mathbb{B} is finite we have $H(\mathbb{B})$ is (quasi) finitely generated. By Theorem 3.1 and Lemma 3.5, there is some $C \in \mathbb{N}$ such that for all $a \in H(\mathbb{B})$ and all $k, \ell \in \mathbf{L}_{H(\mathbb{B})}(a)$ it follows that $\text{d}(\mathbf{Z}_{H(\mathbb{B}), k}(a), \mathbf{Z}_{H(\mathbb{B}), \ell}(a)) \leq C|\ell - k|$. Set $D = \max\{C, N(\mu_1 + 1) + L_1 + \alpha, N(\mu_2 + 1) + L_2 + \alpha\}$. It suffices to show by induction that for all $r \in \mathbb{N}_0$ and $a \in H$ such that $\max\mathbf{L}(a) = r$ we have for all $k, \ell \in \mathbf{L}(a)$ for which $\min\{k, \ell\} + N \geq \max\mathbf{L}(a)$ it follows that $\text{d}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq D|\ell - k|$. Let $r \in \mathbb{N}_0$, $a \in H$ and $k, \ell \in \mathbf{L}(a)$ be such that $\max\mathbf{L}(a) = r$ and $\min\{k, \ell\} + N \geq \max\mathbf{L}(a)$. We have to show that $\text{d}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq D|\ell - k|$. Without restriction let $k < \ell$. By 1 we can assume without restriction that $\mathbf{v}_{p_1}(a) - \mu_1 \max\mathbf{L}(a) < \alpha$. If $\max\mathbf{L}(a) \leq D$, then $\text{d}(\mathbf{Z}_k(a), \mathbf{Z}_\ell(a)) \leq \max\{k, \ell\} \leq \max\mathbf{L}(a) \leq D \leq D|\ell - k|$. Now let $\max\mathbf{L}(a) > D$.

Next we show that for all $s \in \mathbb{N}_0$ and $(w_i)_{i=1}^s \in \mathcal{A}(H)^s$ with $a = \prod_{i=1}^s w_i$ and $s + N \geq \max\mathbf{L}(a)$ it follows that $|\{i \in [1, s] \mid \mathbf{v}_{p_1}(w_i) = \mu_1\}| > L_1$ and $\mathbf{v}_{p_1}(w_j) \leq \mu_1(N + 1) + \alpha$ for all $j \in [1, s]$. Let $s \in \mathbb{N}_0$ and $(w_i)_{i=1}^s \in \mathcal{A}(H)^s$ be such that $a = \prod_{i=1}^s w_i$ and $s + N \geq \max\mathbf{L}(a)$. We have $|\{i \in [1, s] \mid \mathbf{v}_{p_1}(w_i) = \mu_1\}| = s - |\{i \in [1, s] \mid \mathbf{v}_{p_1}(w_i) \neq \mu_1\}| \geq s - \sum_{i=1}^s (\mathbf{v}_{p_1}(w_i) - \mu_1) = s - (\mathbf{v}_{p_1}(a) - s\mu_1) = s(\mu_1 + 1) - \mathbf{v}_{p_1}(a) > s(\mu_1 + 1) - \alpha - \mu_1 \max\mathbf{L}(a) \geq (\max\mathbf{L}(a) - N)(\mu_1 + 1) - \alpha - \mu_1 \max\mathbf{L}(a) = \max\mathbf{L}(a) - N(\mu_1 + 1) - \alpha >$

$D - N(\mu_1 + 1) - \alpha \geq L_1$. Let $j \in [1, s]$. Then $\mathbf{v}_{p_1}(w_j) = \mathbf{v}_{p_1}(a) - \sum_{i=1, i \neq j}^s \mathbf{v}_{p_1}(w_i) \leq \mathbf{v}_{p_1}(a) - (s-1)\mu_1 \leq \alpha + \mu_1 \max \mathbf{L}(a) - (s-1)\mu_1 = \mu_1(\max \mathbf{L}(a) - s + 1) + \alpha \leq \mu_1(N+1) + \alpha$.

We continue with the following two assertions.

A1. There are some $u_1 \in \mathbb{A}(H)$ and $(u_i)_{i=2}^k \in \mathcal{A}(H)^{k-1}$ such that $\mathbf{v}_{p_1}(u_1) = \mu_1$ and $a = \prod_{i=1}^k u_i$.

A2. There are some $v_1 \in \mathbb{A}(H)$ and $(v_j)_{j=2}^\ell \in \mathcal{A}(H)^{\ell-1}$ such that $\mathbf{v}_{p_1}(v_1) = \mu_1$ and $a = \prod_{j=1}^\ell v_j$.

Proof of A1. and A2. By symmetry it is sufficient to prove **A1**. Since $k + N \geq \max \mathbf{L}(a)$ there is some $(w_i)_{i=1}^k \in \mathcal{A}(H)^k$ such that $\mathbf{v}_{p_1}(w_1) = \mathbf{v}_{p_1}(w_2) = \mu_1$ and $a = \prod_{i=1}^k w_i$. It is obvious that $\mu_1 \leq 2\alpha$. If $\mathbf{v}_{p_2}(w_1) \leq 2\alpha$ or $\mathbf{v}_{p_2}(w_2) \leq 2\alpha$, then $w_1 \in \mathbb{A}(H)$ or $w_2 \in \mathbb{A}(H)$ and we are done. Now suppose that $\mathbf{v}_{p_2}(w_1) > 2\alpha$ and $\mathbf{v}_{p_2}(w_2) > 2\alpha$. We have $w_1 = \beta p_1^{\mu_1} p_2^t$ and $w_2 = \gamma p_1^{\mu_1} p_2^s$ for some $\beta, \gamma \in \widehat{H}^\times$ with $t = \mathbf{v}_{p_2}(w_1)$ and $s = \mathbf{v}_{p_2}(w_2)$. Set $u_1 = \beta p_1^{\mu_1} p_2^{2\alpha}$, $u_2 = \gamma p_1^{\mu_1} p_2^{t+s-2\alpha}$ and $u_i = w_i$ for all $i \in [3, k]$. We infer by Lemma 5.10.1 that $u_1, u_2 \in \mathcal{A}(H)$. It is clear that $u_1 \in \mathbb{A}(H)$ and $a = \prod_{i=1}^k u_i$. \square (**A1** and **A2**)

CASE 1: $\max\{\mathbf{v}_{p_2}(u_i) \mid i \in [1, k]\} \cup \{\mathbf{v}_{p_2}(v_i) \mid i \in [1, \ell]\} \leq E_1$.

Observe that $\{u_i \mid i \in [1, k]\} \cup \{v_j \mid j \in [1, \ell]\} \subset \mathbb{B}$. Therefore, we have $k, \ell \in \mathbf{L}_{H(\mathbb{B})}(a)$, and thus $\mathbf{d}(Z_k(a), Z_\ell(a)) \leq \mathbf{d}(Z_{H(\mathbb{B}), k}(a), Z_{H(\mathbb{B}), \ell}(a)) \leq C|\ell - k| \leq D|\ell - k|$.

CASE 2: $\max\{\mathbf{v}_{p_2}(u_i) \mid i \in [1, k]\} > E_1$.

We have $k-1, \ell-1 \in \mathbf{L}(v_1^{-1}a)$, $\max \mathbf{L}(v_1^{-1}a) < \max \mathbf{L}(a)$ and $\min\{k-1, \ell-1\} + N \geq \max \mathbf{L}(v_1^{-1}a)$. Therefore, it follows by the induction hypothesis that $\mathbf{d}(Z_{k-1}(v_1^{-1}a), Z_{\ell-1}(v_1^{-1}a)) \leq D|(k-1) - (\ell-1)| = D|\ell - k|$. There are some $x \in Z_{k-1}(v_1^{-1}a)$ and $y \in Z_{\ell-1}(v_1^{-1}a)$ such that $\mathbf{d}(Z_{k-1}(v_1^{-1}a), Z_{\ell-1}(v_1^{-1}a)) = \mathbf{d}(x, y)$. Clearly, $v_1 x \in Z_k(a)$ and $v_1 y \in Z_\ell(a)$. Consequently, $\mathbf{d}(Z_k(a), Z_\ell(a)) \leq \mathbf{d}(v_1 x, v_1 y) = \mathbf{d}(x, y) \leq D|\ell - k|$.

CASE 3: $\max\{\mathbf{v}_{p_2}(v_i) \mid i \in [1, \ell]\} > E_1$.

This follows by analogy with CASE 2. \square

Proposition 5.12. *Let H be a strongly ring-like monoid of rank $s \leq 2$. Then $\mathbf{c}_{\text{mon}}(H) < \infty$ and $\delta_w(H) < \infty$.*

Proof. If $s = 1$, then the assertion follows from Remark 5.3.2. Now suppose that $s = 2$. We infer by [28, Theorem 5.3] that $\mathbf{c}_{\text{mon}}(H) < \infty$. Both Lemmas, 5.10.2 and 5.11.3, together with Lemma 5.1 show that the assumptions of Lemma 3.6 are satisfied whence we obtain that $\delta_w(H) < \infty$. \square

Now we formulate the main result of the present section. The crucial point in its proof is that the finiteness of the weak successive distance (achieved in Proposition 5.12) and the validity of the Structure Theorem for Sets of Lengths preserve the finiteness of the monotone catenary degree when passing to direct products (see Theorem 3.8).

Theorem 5.13. *Let H be a v -Noetherian weakly Krull monoid with $\emptyset \neq \mathfrak{f} = (H : \widehat{H})$ such that $H_{\mathfrak{p}}$ is strongly ring-like of rank $s_{\mathfrak{p}} \leq 2$ for each $\mathfrak{p} \in \mathfrak{X}(H)$ with $\mathfrak{p} \supset \mathfrak{f}$.*

1. $\mathcal{I}_v^*(H)$ is a C-monoid and if $\mathcal{C}_v(H)$ is finite, then H is a C-monoid.
2. $\mathcal{I}_v^*(H)$ satisfies the Structure Theorem for Sets of Lengths and the Structure Theorem for Unions.
3. $\delta_w(\mathcal{I}_v^*(H)) < \infty$, and $\mathbf{c}_{\text{mon}}(\mathcal{I}_v^*(H)) < \infty$.

Proof. By [21, Theorem 5.5], there is a monoid isomorphism

$$(5.2) \quad \chi: \mathcal{I}_v^*(H) \rightarrow \mathcal{F}(\mathcal{P}) \times \prod_{\mathfrak{p} \in \mathcal{P}^*} (H_{\mathfrak{p}})_{\text{red}} \quad \text{satisfying} \quad \chi|_{\mathcal{P}} = \text{id}_{\mathcal{P}}.$$

where $\mathcal{P} = \{\mathfrak{p} \in \mathfrak{X}(R) \mid \mathfrak{p} \not\supset \mathfrak{f}\}$ and $\mathcal{P}^* = \mathfrak{X}(H) \setminus \mathcal{P}$. Observe that \mathcal{P}^* is finite.

1. Clearly, $\mathcal{F}(\mathcal{P})$ is a C-monoid. It follows from [19, Corollary 2.9.8] that all $H_{\mathfrak{p}}$ (and thus all $(H_{\mathfrak{p}})_{\text{red}}$) with $\mathfrak{p} \in \mathcal{P}^*$ are C-monoids. Since $\widehat{H}_{\mathfrak{p}}^\times / H_{\mathfrak{p}}^\times$ is finite for all $\mathfrak{p} \in \mathcal{P}^*$ we infer by [19, Theorem 2.9.16] that

$\mathcal{F}(\mathcal{P}) \times \prod_{\mathfrak{p} \in \mathcal{P}^*} (H_{\mathfrak{p}})_{\text{red}} \cong (\mathcal{F}(\mathcal{P}) \times \prod_{\mathfrak{p} \in \mathcal{P}^*} H_{\mathfrak{p}})_{\text{red}}$ is a C-monoid. Therefore, $\mathcal{I}_v^*(H)$ is a C-monoid. Now let $\mathcal{C}_v(H)$ be finite. We have $H_{\text{red}} \cong \{aH \mid a \in H\}$, $\{aH \mid a \in H\} \subset \mathcal{I}_v^*(H)$ is a saturated submonoid and $\mathcal{I}_v^*(H)/\{aH \mid a \in H\} \subset \mathcal{C}_v(H)$ is finite. Therefore, it follows by [19, Theorems 2.9.10 and 2.9.16] that H is a C-monoid.

2. and 3. Note that all $H_{\mathfrak{p}}$ (and thus all $(H_{\mathfrak{p}})_{\text{red}}$) with $\mathfrak{p} \in \mathcal{P}^*$ satisfy the Structure Theorem for Sets of Lengths by Lemma 5.1, and they have finite weak successive distance and finite equal catenary degree by Proposition 5.12. By (5.2) and Theorem 3.8, $\mathcal{I}_v^*(H)$ satisfies the Structure Theorem for Sets of Lengths, $\delta_w(\mathcal{I}_v^*(H)) < \infty$, and $c_{\text{eq}}(\mathcal{I}_v^*(H)) < \infty$. We infer by Lemma 3.5 that $c_{\text{mon}}(\mathcal{I}_v^*(H)) < \infty$. Since $\mathcal{I}_v^*(H)$ is a C-monoid by 1, and C-monoids satisfy the Structure Theorem for Unions by [17, Theorems 3.10 and 4.2], $\mathcal{I}_v^*(H)$ satisfies the Structure Theorem for Unions. \square

Corollary 5.14. *Let R be a weakly Krull Mori domain with $(R : \widehat{R}) \neq \{0\}$ such that $R_{\mathfrak{p}}$ is Noetherian, $|R/\mathfrak{p}| < \infty$, and $|\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}| \leq 2$ for each $\mathfrak{p} \in \mathfrak{X}(R)$ with $\mathfrak{p} \supset (R : \widehat{R})$. Then the monoid $\mathcal{I}_v^*(R)$ has finite weak successive distance and finite monotone catenary degree.*

In particular, if R is an order in a quadratic number field, then $\mathcal{I}^(R)$ has finite weak successive distance and finite monotone catenary degree.*

Comment. The example given in Remark 5.3.4 shows that the assumption, that $|\{\mathfrak{m} \in \mathfrak{X}(\widehat{R}) \mid \mathfrak{m} \cap R = \mathfrak{p}\}| \leq 2$ for each $\mathfrak{p} \in \mathfrak{X}(R)$, is crucial.

Proof. This follows from Theorem 5.13 and Proposition 5.9.3. Suppose that R is an order in a quadratic number field. Then R is one-dimensional Noetherian whence $\mathcal{I}^*(R) = \mathcal{I}_v^*(R)$, and $\mathcal{I}_v^*(R)$ has finite weak successive distance and finite monotone catenary degree by the main statement. \square

REFERENCES

- [1] D.D. Anderson and O.A. Al-Mallah, *Commutative group rings that are présimplifiable or domainlike*, J. Algebra Appl. **16** (2017).
- [2] D.D. Anderson, D.F. Anderson, and M. Zafrullah, *Atomic domains in which almost all atoms are prime*, Commun. Algebra **20** (1992), 1447 – 1462.
- [3] D.D. Anderson and S. Chun, *Commutative rings with finitely generated monoids of fractional ideals*, J. Algebra **320** (2008), 3006 – 3021.
- [4] D.D. Anderson, J. Mott, and M. Zafrullah, *Finite character representations for integral domains*, Boll. Unione Mat. Ital. **6** (1992), 613 – 630.
- [5] D.D. Anderson and J.L. Mott, *Cohen-Kaplansky domains: integral domains with a finite number of irreducible elements*, J. Algebra **148** (1992), 17 – 41.
- [6] D.D. Anderson and S. Valdes-Leon, *Factorization in commutative rings with zero divisors*, Rocky Mt. J. Math. **26** (1996), 439 – 480.
- [7] V. Blanco, P. A. García-Sánchez, and A. Geroldinger, *Semigroup-theoretical characterizations of arithmetical invariants with applications to numerical monoids and Krull monoids*, Illinois J. Math. **55** (2011), 1385 – 1414.
- [8] A. Bouvier, *Résultats nouveaux sur les anneaux présimplifiables*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A955–A957.
- [9] J. Coykendall and W.W. Smith, *On unique factorization domains*, J. Algebra **332** (2011), 62 – 70.
- [10] Y. Fan, A. Geroldinger, F. Kainrath, and S. Tringali, *Arithmetic of commutative semigroups with a focus on semigroups of ideals and modules*, J. Algebra Appl. **11** (2017), 1750234 (42 pages).
- [11] Y. Fan and S. Tringali, *Power monoids: A bridge between factorization theory and arithmetic combinatorics*, J. Algebra **512** (2018), 252 – 294.
- [12] M. Fontana, E. Houston, and T. Lucas, *Factoring Ideals in Integral Domains*, Lecture Notes of the Unione Matematica Italiana, vol. 14, Springer, 2013.
- [13] A. Foroutan, *Monotone chains of factorizations*, Focus on commutative rings research (A. Badawi, ed.), Nova Sci. Publ., New York, 2006, pp. 107 – 130.
- [14] A. Foroutan and A. Geroldinger, *Monotone chains of factorizations in C-monoids*, Arithmetical Properties of Commutative Rings and Monoids, Lect. Notes Pure Appl. Math., vol. 241, Chapman & Hall/CRC, 2005, pp. 99 – 113.
- [15] A. Foroutan and W. Hassler, *Chains of factorizations and factorizations with successive lengths*, Commun. Algebra **34** (2006), 939 – 972.

- [16] C. Frei and S. Frisch, *Non-unique factorization of polynomials over residue class rings of the integers*, Commun. Algebra **39** (2011), 1482 – 1490.
- [17] W. Gao and A. Geroldinger, *On products of k atoms*, Monatsh. Math. **156** (2009), 141 – 157.
- [18] A. Geroldinger, D.J. Gryniewicz, G.J. Schaeffer, and W.A. Schmid, *On the arithmetic of Krull monoids with infinite cyclic class group*, J. Pure Appl. Algebra **214** (2010), 2219 – 2250.
- [19] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.
- [20] A. Geroldinger, W. Hassler, and G. Lettl, *On the arithmetic of strongly primary monoids*, Semigroup Forum **75** (2007), 567 – 587.
- [21] A. Geroldinger, F. Kainrath, and A. Reinhart, *Arithmetic of seminormal weakly Krull monoids and domains*, J. Algebra **444** (2015), 201 – 245.
- [22] A. Geroldinger, S. Ramacher, and A. Reinhart, *On v -Marot Mori rings and C -rings*, J. Korean Math. Soc. **52** (2015), 1 – 21.
- [23] A. Geroldinger and P. Yuan, *The monotone catenary degree of Krull monoids*, Result. Math. **63** (2013), 999 – 1031.
- [24] K. R. Goodearl, *von Neumann regular rings*, Monographs and Studies in Mathematics, vol. 4, Pitman (Advanced Publishing Program), Boston, Mass.-London, 1979.
- [25] F. Halter-Koch, *Divisor theories with primary elements and weakly Krull domains*, Boll. Un. Mat. Ital. B **9** (1995), 417 – 441.
- [26] ———, *Ideal Systems. An Introduction to Multiplicative Ideal Theory*, Marcel Dekker, 1998.
- [27] F. Halter-Koch, W. Hassler, and F. Kainrath, *Remarks on the multiplicative structure of certain one-dimensional integral domains*, Rings, Modules, Algebras, and Abelian Groups, Lect. Notes Pure Appl. Math., vol. 236, Marcel Dekker, 2004, pp. 321 – 331.
- [28] W. Hassler, *Properties of factorizations with successive lengths in one-dimensional local domains*, J. Commut. Algebra **1** (2009), 237 – 268.
- [29] A. Philipp, *A precise result on the arithmetic of non-principal orders in algebraic number fields*, J. Algebra Appl. **11**, 1250087, 42pp.
- [30] ———, *A characterization of arithmetical invariants by the monoid of relations*, Semigroup Forum **81** (2010), 424 – 434.
- [31] A. Reinhart, *Structure of general ideal semigroups of monoids and domains*, J. Commut. Algebra **4** (2012), 413 – 444.
- [32] ———, *On integral domains that are C -monoids*, Houston J. Math. **39** (2013), 1095 – 1116.
- [33] S. Tringali, *Structural properties of subadditive families with applications to factorization theory*, Israel J. Math., to appear, <https://arxiv.org/abs/1706.03525>.

INSTITUT FÜR MATHEMATIK UND WISSENSCHAFTLICHES RECHNEN, KARL-FRANZENS-UNIVERSITÄT GRAZ, NAWI GRAZ,
HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA

Email address: alfred.geroldinger@uni-graz.at, andreas.reinhart@uni-graz.at

URL: <http://imsc.uni-graz.at/geroldinger>, <http://imsc.uni-graz.at/reinhart>